



UWL REPOSITORY

repository.uwl.ac.uk

The impact of novel coronavirus SARS-CoV-2 among healthcare workers in hospitals: An aerial overview

Xiang, Boqi, Li, Peining, Yang, Xinhui, Zhong, Shuyi, Manyande, Anne ORCID: <https://orcid.org/0000-0002-8257-0722> and Feng, Maohui (2020) The impact of novel coronavirus SARS-CoV-2 among healthcare workers in hospitals: An aerial overview. *American Journal of Infection Control*, 48 (8). pp. 915-917. ISSN 0196-6553

<http://dx.doi.org/10.1016/j.ajic.2020.05.020>

This is the Accepted Version of the final output.

UWL repository link: <https://repository.uwl.ac.uk/id/eprint/7214/>

Alternative formats: If you require this document in an alternative format, please contact: open.research@uwl.ac.uk

Copyright: Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy: If you believe that this document breaches copyright, please contact us at open.research@uwl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

1 **The impact of novel coronavirus SARS-CoV-2 among healthcare workers in hospitals: An**
2 **aerial overview**

3

4 Boqi Xiang¹, Peining Li¹, Xinhui Yang², Shuyi Zhong³, Anne Manyande⁴, *Maohui Feng⁵,

5

6 ¹University of California Davis, Davis, CA, USA;

7 ²School of Public Health, Brown University, Providence, RI, USA;

8 ³Nebula Care inc, San Francisco, CA, USA.

9 ⁴School of Human and Social Sciences, University of West London, London, UK

10 ⁵Department of Gastrointestinal Surgery, Wuhan Peritoneal Cancer Clinical Medical Research Center, Zhongnan
11 Hospital of Wuhan University, Hubei Key Laboratory of Tumor Biological Behaviors and Hubei Cancer Clinical
12 Study Center, Wuhan 430071, Hubei, China.

13

14

15 *** Correspondence:**

16 **Maohui Feng:** E-mail: Fengmh@whu.edu.cn

17

18

19

20

21

22

23

24

25

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

Abstract:

The ongoing outbreak of COVID-19, caused by the novel coronavirus SARS-CoV-2, places healthcare workers at an increased risk of infection as they are in close contact with patients. In this article, we report an overview of cases of infected healthcare workers in China and Italy during the early periods of the COVID-19 epidemic. China’s coronavirus response highlights the importance of implementing effective public health strategies. The authorities worldwide, therefore, need to be extremely cautious when they implement stringent protective measures that safeguard healthcare workers in hospitals and counteract the threats created by the pandemic.

Keywords:

Coronavirus SARS-CoV-2, COVID-19 disease, Healthcare workers, Protective measures.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

Introduction

Daily, the outbreak of the Coronavirus disease, COVID-19, seems to reach a new and tragic milestone. Current epidemiological data shows that person-to-person transmission of coronavirus SARS-CoV-2 is more likely to be occurring [1-4]. As of March 22, 2020, the World Health Organization (WHO) reported 267,013 confirmed cases, including 11,201 deaths, in 184 countries and regions. The virus spreads mainly between people who are in close contact (6 feet) with an infected person through respiratory pathways, though it can also be transmitted through contact with contaminated surfaces or objects. In this case, every individual is potentially susceptible to the virus as no vaccine is currently available. Data from the Chinese Center for Disease Control and Prevention indicates that leading up to February 11, 2020, a total of 1716 healthcare workers became infected and five had died (0.3%) [5]. In addition, according to a more recent report from the Italian Institute of Higher Health and National Federation of surgeons and Dentists, from 1st to 23rd March 2020 4826 healthcare workers were infected with COVID-19 (9% of confirmed cases) and 18 had died.

It is therefore, imperative to prevent further transmission of COVID-19 in medical facilities to protect healthcare workers. In this article, we report the findings of infected cases in hospitals during the early outbreak period of COVID-19. The purpose of this paper is for readers to better understand the epidemiological characteristics and protective measures by which the coronavirus SARS-CoV-2 caused the COVID-19 disease in the tens and thousands of patients in Wuhan and worldwide.

1 **Methods**

2 We collected publicly available data from the National Health Commission, Hubei Health
3 Commission, Wuhan Health Commission National, China Center for Disease Control and
4 Prevention (CDC), Italy, and the World Health Organization (WHO). We also searched literature
5 reporting healthcare workers infected with CoVID-19 in hospitals using the PubMed database and
6 Cochrane Library from January 1, 2020 to March 24, 2020. Search terms included ‘healthcare
7 workers’ or ‘medical staff’ and ‘novel coronavirus’ or ‘COVID-19’ or ‘2019-nCoV’.

8

9 **Results**

10 **Geographical epidemiological characteristics of 11 hospitals with infected healthcare**
11 **workers**

12 By the end of January 2020, 11 hospitals in Wuhan had reported over 15 confirmed cases among
13 healthcare workers. Geographical distribution of these hospitals is shown in Fig.1A. According to
14 the date of the first confirmed case with COVID-19 detected using viral nucleic acid testing, these
15 hospitals were arranged in sequence and coded 1 to 11. Hospital-2, -3, and -11 are closer to Huanan
16 Seafood Wholesale Market in comparison to the rest, and reported 63, 60, and 115 COVID-19-
17 infected cases (including confirmed and suspected cases) of healthcare workers respectively
18 (Fig.1B). Three hospitals farther from Huanan Seafood Wholesale Market are located in the
19 southern region of the Yangtze River and reported 194, 66, and 44 COVID-19-infected cases
20 respectively (Fig.1B).

21

22 **Characteristics of Chinese medical staff infected during the early period of the COVID-19**
23 **outbreak**

1 Data from hospital-6 showed that of the 138 patients with COVID-19, 40 (29%) were medical
2 staff who were infected in the hospital setting. Moreover, over 10 healthcare workers in the
3 surgical department were presumed to have been infected by a confirmed patient presenting with
4 abdominal symptoms. Among those infected healthcare workers, 31(77.5%) worked on general
5 outpatient clinics and wards, 7 (17.5%) were in the emergency department, and 2 (5%) were in the
6 intensive care unit (ICU) (Fig.2). In addition, there were 230 diagnosed cases (130 inpatient cases
7 and 100 home-quarantine cases) in hospital-11 by the end of February 2020.

8
9 The Chinese National Health Commission is the authoritative organization responsible for
10 publishing data on medical staff infections. According to the National Health Commission, on 11th
11 February 2020, among the 44672 confirmed cases, a total of 1716 cases were healthcare workers
12 (3.8%). Specifically, 1502 were in Hubei (87.5%) and 1102 in Wuhan (64.2%) (Fig.2). After
13 implementing the clinical diagnostic criteria, the number of clinically diagnosed cases had surged
14 by 13th February 2020. On 24th February 2020, the WHO-China Expert Group conference stated
15 that a total of 3387 infected cases (2055 confirmed cases, 1070 diagnosed cases, and 157 suspected
16 cases) were healthcare workers in 476 health-care settings, of whom 3062 were in Hubei (90.4%).
17 These results suggest that the number of infected healthcare workers were mostly in Hubei
18 province which is the epicenter of the outbreak in China.

19 20 **Characteristics of Italian medical staff infected during the early period of the COVID-19** 21 **outbreak**

22 COVID-19 has spread rapidly with confirmed cases currently reported in more than 180 countries.
23 The WHO declared COVID-19 a pandemic on 11th March 2020. To provide an understanding of

1 the epidemic for medical staff, the case of Italy is described in detail. As of 26th March 2020, a
2 total of 74386 confirmed cases had been reported in Italy with 7503 deaths. To date (17th March
3 2020), according to the Italian Institute of Higher Health, a total of 2629 healthcare workers were
4 infected with COVID-19 (8.3% of confirmed cases). Only six days later, there were 4826
5 healthcare workers infected with COVID-19 (9 % of confirmed cases) with 18 deaths (Fig.2).

6

7 **Discussion**

8 Since COVID-19 is a novel disease with a high transmission rate [6-8], it presents a greater risk
9 of infection for healthcare workers who are in close contact with patients. Huang et al[2] first
10 reported 41 COVID-19 cases known to have had a history of exposure to the Huanan Seafood
11 Wholesale Market. Thus, we described geographical epidemiological characteristics of 11
12 hospitals with infected healthcare workers illustrated in Fig.1. To date (24th March 2020), there
13 were 230 healthcare workers infected with COVID-19 in hospital-11 which is closer to Huanan
14 Seafood Wholesale Market and had 4 deaths. The data suggest that several cases among the
15 infected healthcare workers may be closely linked to the geographical proximity to the Market
16 during the early phase of the COVID-19 outbreak, especially the beginning of January 2020.

17

18 We analyzed the data of 12 deceased healthcare workers and found that their dates of symptom
19 onset were at the beginning and middle of January 2020 (Fig.2). Because of the 14-day latency, it
20 was speculated that they were infected with the novel coronavirus SARS-CoV-2 between late
21 December 2019 and early January 2020. Notably, at that time, many important questions involving
22 the SARS-CoV-2 infections remained unanswered, including its ability and duration of
23 transmission in humans and clinical consequences of human infections. Unfortunately, a

1 substantial number of healthcare workers were infected in healthcare settings before the end of
2 January 2020. Although SARS-CoV-2 itself is significantly infectious, it should not be held fully
3 responsible for the extensive spread of the infection. In fact, the main reasons for earlier virus
4 transmission to healthcare workers were due to lack of awareness of adequate protection of
5 healthcare workers and leaders in health-care settings, and later transmissions were as a result of
6 insufficient supply of protective devices (Fig.2). Moreover, to date, there is zero infection among
7 the 42, 322 medical staff from across China who voluntarily came to Wuhan to support local
8 medical healthcare professionals from 25th January to 26th March 2020. This suggests that high-
9 grade protection and social distancing are crucial strategies to prevent transmission in hospitals.

10

11 In an article from *The Lancet* published online on January 24, 2020, Wang and colleagues[3] put
12 forward a proposition that we have to be aware of the challenges SARS-CoV-2 poses to our worlds.
13 Specifically, we need to be wary of the current COVID-19 outbreak turning into a sustained
14 epidemic or even a pandemic associated with substantial morbidity and mortality. Though SARS-
15 CoV-2 put all health organizations in China on high alert [9], other countries and regions seemed
16 to underestimate the transmission rate of the highly contagious virus. Mounting evidence shows
17 that the viral shedding of asymptomatic or mild cases of SARS-CoV-2 spread to infect others[10-
18 13]. This may explain why the virus SARS-CoV-2 spread so quickly in Europe and is now swiftly
19 circulating around the world. Before the WHO declared COVID-19 a pandemic on 11th March,
20 2020, the Italian government did not implement strong measures through public health
21 interventions, including case detection, wearing protective masks, closing public spaces,
22 cancelling public gatherings, and self-isolation at home. Moreover, insufficient protective supplies
23 fueled the infection of healthcare workers in hospitals. All these factors contributed to the current

1 pandemic in Italy and other hardest hit countries such as Spain, France, Iran, United Kingdom and
2 the United States of America.

3

4 **Conclusion**

5 The ongoing outbreak of COVID-19 worldwide highlights the importance of implementing
6 effective public health strategies. What can China’s coronavirus response teach the rest of the
7 world? The answer is that authorities need to be extremely cautious and implement stringent
8 protective measures to safeguard our healthcare workers in order to counteract the threats brought
9 by the pandemic.

10

11 **Acknowledgement**

12 This work was supported by the National Natural Science Foundation of China (No. 81873467,
13 81670240, 81770283) and the Clinical Medical Research Center of Peritoneal Cancer of Wuhan
14 (No. 2015060911020462) and the Natural Science Foundation of Hubei Province (No. 2015-
15 CFA027), the Research Foundation of Health and Family Planning Commission of Hubei Province
16 (No. WJ2015MA010).

17

18

19

20

21

22

23


24

25

1 **References**

- 2
- 3 [1] Chan JF, Yuan S, Kok KH, To KK, Chu H, Yang J, Xing F, Liu J, Yip CC, Poon RW,
4 Tsoi HW, Lo SK, Chan KH, Poon VK, Chan WM, Ip JD, Cai JP, Cheng VC, Chen H,
5 Hui CK and Yuen KY. A familial cluster of pneumonia associated with the 2019 novel
6 coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet*
7 2020;
- 8 [2] Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X, Cheng Z,
9 Yu T, Xia J, Wei Y, Wu W, Xie X, Yin W, Li H, Liu M, Xiao Y, Gao H, Guo L, Xie J,
10 Wang G, Jiang R, Gao Z, Jin Q, Wang J and Cao B. Clinical features of patients infected
11 with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;
- 12 [3] Wang C, Horby PW, Hayden FG and Gao GF. A novel coronavirus outbreak of global
13 health concern. *Lancet* 2020;
- 14 [4] Phan LT, Nguyen TV, Luong QC, Nguyen TV, Nguyen HT, Le HQ, Nguyen TT, Cao
15 TM and Pham QD. Importation and Human-to-Human Transmission of a Novel
16 Coronavirus in Vietnam. *N Engl J Med* 2020;
- 17 [5] Prevention CCfDca. The epidemiological characteristics of an outbreak of 2019 novel
18 coronavirus Disease (COVID-19) in China. *Chin J Epidemiol* 2020; 41:
- 19 [6] Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, Wang B, Xiang H, Cheng Z, Xiong Y,
20 Zhao Y, Li Y, Wang X and Peng Z. Clinical Characteristics of 138 Hospitalized Patients
21 With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA* 2020;
- 22 [7] Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, Qiu Y, Wang J, Liu Y, Wei Y, Xia J,
23 Yu T, Zhang X and Zhang L. Epidemiological and clinical characteristics of 99 cases of
24 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020;
- 25 [8] Habibzadeh P and Stoneman EK. The Novel Coronavirus: A Bird's Eye View. *Int J*
26 *Occup Environ Med* 2020; 11: 65-71.
- 27 [9] Khan S, Ali A, Siddique R and Nabi G. Novel coronavirus is putting the whole world on
28 alert. *J Hosp Infect* 2020;
- 29 [10] Rothe C, Schunk M, Sothmann P, Bretzel G, Froeschl G, Wallrauch C, Zimmer T, Thiel
30 V, Janke C, Guggemos W, Seilmaier M, Drosten C, Vollmar P, Zwirgmaier K, Zange S,
31 Wolfel R and Hoelscher M. Transmission of 2019-nCoV Infection from an
32 Asymptomatic Contact in Germany. *N Engl J Med* 2020;
- 33
34
35
36
37
38
39
40
41
42


1 Fig.1 Geographical epidemiological characteristics of 11 hospitals with infected healthcare
 2 workers. (A) Geographical distribution of 11 hospitals. (B) According to the date of the first
 3 confirmed case with COVID-19 detected using viral nucleic acid testing, 11 hospitals were
 4 arranged in sequence from code 1 to 11. Confirmed and suspected cases of every hospital were
 5 identified by the end of January 2020.



6
 7 Fig.2 The graph's x-axis (dates from 1st January to 26th March 2020) is used as a timeline of the
 8 key events and dynamic profile of infected healthcare workers in the different phases of the
 9 COVID-19 outbreak. The key events include the international public health emergency and the
 10 declaration of COVID-19 as a pandemic by the WHO. The profiles of infected healthcare workers
 11 are shown in blue boxes;
 12 The data of 12 deceased healthcare workers are shown in the yellow box. Italian information is
 13 shown in the red box.

1

Figure 2 Dynamic Profile of COVID-19-infected healthcare workers



2

3