

UWL REPOSITORY

repository.uwl.ac.uk

Environmental heterogeneity of Staphylococcus species from alkaline fermented foods and associated toxins and antimicrobial resistance genetic elements

Ouoba, L. Irène I., Voudibio Mbozo, Alain Brice, Anyogu, Amarachukwu ORCID: <https://orcid.org/0000-0001-9652-7728>, Obioha, Promiselynda I., Lingani-Sawadogo, Hagrétou, Sutherland, Jane P., Jespersen, Lene and Ghoddusi, Hamid B. (2019) Environmental heterogeneity of Staphylococcus species from alkaline fermented foods and associated toxins and antimicrobial resistance genetic elements. International Journal of Food Microbiology, 311. p. 108356. ISSN 0168-1605

<http://dx.doi.org/10.1016/j.ijfoodmicro.2019.108356>

This is the Accepted Version of the final output.

UWL repository link: <https://repository.uwl.ac.uk/id/eprint/8242/>

Alternative formats: If you require this document in an alternative format, please contact: open.research@uwl.ac.uk

Copyright: Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy: If you believe that this document breaches copyright, please contact us at open.research@uwl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Environmental heterogeneity of *Staphylococcus* species from alkaline fermented foods and associated toxins and antimicrobial resistance genetic elements

L. Irène I. Ouoba^{a,b*}, Alain Brice Voudibio-Mbozo^c, Amarachukwu Anyogu^d, Promiselynda I. Obioha^a, Hagrétou Lingani-Sawadogo^e, Jane P. Sutherland^a, Lene Jespersen^f, and Hamid B. Ghoddusi^a

^a Microbiology Research Unit, School of Human Sciences, London Metropolitan University, UK

^b Ouoba-Consulting, London, UK

^c Faculté des Sciences, Université Marien N’Gouabi, Brazzaville, Congo

^d Department of Life Sciences, Faculty of Science and Technology, University of Westminster, UK

^e Département Technologie Alimentaire/IRSAT/CNRST, Ouagadougou, Burkina Faso

^f Microbial Ecology and Food Fermentation, Faculty of Sciences, University of Copenhagen

Corresponding author: Dr L. Irène I. Ouoba, Microbiology Research Unit, School of Human Sciences, London Metropolitan University London N7 8DB, UK

Tel.: +44 (0) 20 71334154, Email: ouobairene@hotmail.com

Abstract

Different samples of three products including Bikalga and Soumbala from Burkina Faso (West Africa) and Ntoba Mbodi from Congo-Brazzaville (Central Africa) were evaluated. The bacteria (400) were phenotyped and genotypically characterized by Rep-PCR, PFGE, 16S rRNA and *rpoB* gene sequencing and *spa* typing. Their PFGE profiles were compared with those of 12,000 isolates in the Center for Disease Control (CDC, USA) database. They were screened for the production of enterotoxins, susceptibility to 19 antimicrobials, presence of 12 staphylococcal toxin and 38 AMR genes and the ability to transfer erythromycin and tetracycline resistance genes to *Enterococcus faecalis* JH2-2.

Fifteen coagulase negative (CoNS) and positive (CoPS) species characterised by 25 Rep-PCR/PFGE clusters were identified: *Staphylococcus arlettae*, *S. aureus*, *S. cohnii*, *S. epidermidis*, *S. gallinarum*, *S. haemolyticus*, *S. hominis*, *S. pasteurii*, *S. condimenti*, *S. piscifermentans*, *S. saprophyticus*, *S. sciuri*, *S. simulans*, *S. warneri* and *Macrococcus caseolyticus*. Five species were specific to Soumbala, four to Bikalga and four to Ntoba Mbodi. Two clusters of *S. gallinarum* and three of *S. sciuri* were particular to Burkina Faso. The *S. aureus* isolates exhibited a *spa* type t355 and their PFGE profiles did not match any in the CDC database. Bacteria from the same cluster displayed similar AMR and toxin phenotypes and genotypes, whereas clusters peculiar to a product or a location generated distinct profiles. The toxin genes screened were not detected and the bacteria did not produce the staphylococcal enterotoxins A, B, C and D. AMR genes including *bla_zA*, *cat501*, *dfr(A)*, *dfr(G)*, *mecA*, *mecA1*, *msr(A)* and *tet(K)* were identified in CoNS and CoPS. Conjugation experiments produced JH2-2 isolates that acquired resistance to erythromycin and tetracycline, but no gene transfer was revealed by PCR.

The investigation of the heterogeneity of *Staphylococcus* species from alkaline fermented foods, their relationship with clinical and environmental isolates and their safety in relation to antimicrobial resistance (AMR) and toxin production is anticipated to contribute to determining the importance of staphylococci in alkaline fermented foods, especially in relation to the safety of the consumers.

Key words: Alkaline fermented foods; *Staphylococcus* spp.; diversity; distribution; safety; toxins; AMR

1. Introduction

Alkaline fermented foods are products that are generally rich in proteins (30-50%) and their production involves a fermentation process associated with an increase of pH to values as high as 10.0 (Parkouda et al., 2009; Voudibio Mbozo et al., 2017). They are mainly known in Africa and Asia, where they constitute low cost sources of protein foods and play an important part in the diet of people (Parkouda et al., 2009). It is generally recognized that the pH rise is the result of accrued degradation of raw material proteins that leads to the release of peptides, essential amino acids as well as alkalinizing substances such as ammonia (Kiers et al., 2000). In some cases, alkali-treatments during the production are responsible for the pH augmentation (Parkouda et al., 2008).

Bikalga, Soumbala, and Ntoba Mbodi are alkaline fermented foods which are produced through the traditional fermentation of roselle seeds (*Hibiscus sabdariffa*), African locust bean seeds (*Parkia biglobosa*) and cassava leaves (*Manihot esculenta* Crantz) respectively. Production of Bikalga and Soumbala comprises several processing stages of the seeds: cleaning, boiling (about 95°C for 24-40 h), dehulling (Soumbala), fermentation (48-72 h), steaming (Bikalga) and drying (Ouoba et al., 2008a; Parkouda et al., 2008; Parkouda et al., 2009). They are the most popular traditional fermented foods used as condiments in Burkina Faso (West Africa) and are also produced in other African countries. Ntoba Mbodi, which is popular in the Republic of the Congo (Central Africa), is made of cassava leaves and the production process does not involve heating steps. After being cleaned and cut, the leaves are distributed into small portions in larger leaves, wrapped and left to ferment for 2-4 days (Voudibio Mbozo et al., 2017). The fermented cassava leaves are then used to prepare sauces, which are eaten with starch-based foods. In Bikalga, Soumbala and Ntoba Mbodi, pH

values up to 8.5, 9 and 10 have been respectively reported (Ouoba, 2015; Ouoba, 2017; Parkouda et al., 2009; Voudibio Mbozo et al., 2017).

The key microorganisms involved in alkaline fermentations are *Bacillus* species such as *B. subtilis*, *B. licheniformis*, *B. amyloliquefaciens*, *B. pumilis*, and *B. xylanilyticus* that can degrade proteins, lipids, and carbohydrates of the raw materials and bring about the nutritional quality and safety of the product (Parkouda et al., 2009). However, secondary microorganisms such as lactic acid bacteria, *Staphylococcus* spp. and various Gram-positive and Gram-negative bacteria are often isolated with no clear indication of their role in the fermentation. The production techniques of most alkaline fermented foods are traditional, leading to variability of microbial population, nutritional quality, product stability and potential safety issues. It is believed that some of the secondary microorganisms are in fact contaminants from the producers, processing materials and environments that have poor hygienic quality in most traditional fermentation settings. Some secondary microorganisms are part of the original microbiota of the raw materials that are suppressed by products released during the fermentation. For example, those that are alkali sensitive will be eliminated or reduced during the fermentation.

Staphylococci are Gram-positive non-motile cocci ubiquitously spread in nature, and are commonly isolated from humans, animals, food and environmental sources. More than 70 species and subspecies of the *Staphylococcus* genus have been characterized so far (LPSN, 2018) with *Staphylococcus aureus* being one of the most studied species due to its clinical importance. *Staphylococcus* species are usually divided into two groups i.e. coagulase-positive (CoPS) and coagulase-negative (CoNS) according to their ability to coagulate rabbit plasma. Various *Staphylococcus* species have been recurrently isolated in many alkaline and non-

alkaline vegetable based fermented foods, but their role in such fermentation is unknown. However, their importance in the fermentation of meat-based products has been well documented. They occur in the production of various types of sausages where they constitute a part of the autochthonous microbiota or are added as starter cultures in combination with lactic acid bacteria (Fonseca et al., 2013; Milicevic et al., 2014). *Staphylococcus equorum*, *S. saprophyticus*, *S. xylosus* and *S. carnosus* are the most common species found in fermented sausages, but other species such as *S. warneri*, *S. vitulinus*, *S. pasteurii*, *S. epidermidis*, *S. lentus*, *S. haemolyticus*, *S. intermedius*, and *S. hominis* can also be occasionally isolated (Milicevic et al., 2014). They have been reported to be involved in biochemical activities during meat fermentation such as the degradation of proteins, lipids and peroxide as well as the reduction of nitrite (Casaburi et al., 2008; Milicevic et al., 2014). These activities generate compounds such as peptides, amino acids, carbonyls and volatile compounds that contribute to the nutritional quality, specific organoleptic properties and safety of the fermented meat products (Milicevic et al., 2014).

Although beneficial properties of *Staphylococcus* species in food have been documented, many species are most known worldwide for causing food poisoning and other types of clinical infections in humans and animals including e.g. bacteraemia, wound infections, pyogenic lesions, and mastitis. Moreover, they present detrimental antimicrobial resistance (AMR) profiles that make some of their infections hard to treat. With regard to toxin production in food, staphylococcal food poisoning is one of the most common food-borne diseases. It results from the ingestion of heat stable staphylococcal enterotoxins (SEs) produced in food by enterotoxigenic strains of mainly *S. aureus* (Hennekinne et al., 2012). Symptoms related to ingestion of such toxins are fast-acting, occurring generally within 30 min to 8 h and lasting for about 24

hours. Individuals generally experience vomiting, nausea, stomach cramps, and diarrhoea. Severe illnesses related to staphylococcal food poisoning are rare but occasionally, it can be deadly, especially in children, elderly and patients weakened by a long-term illness. *Staphylococcus aureus* is considered one of the world's leading causes of disease outbreaks associated with food consumption. Other CoPS such as *S. intermedius* and CoNS such as *S. epidermidis*, *S. sciuri*, *S. simulans*, *S. xylosus*, *S. equorum*, *S. lentus* and *S. capitis* have been reported to harbour genes coding for SEs and to produce some SEs (Casaes Nunes et al., 2015; Crass and Bergdoll, 1986). However, information on their real implication in staphylococcal food poisoning outbreaks is scarce (Breckinridge and Bergdoll, 1971; Podkowik et al., 2013). Significantly, the incidence of AMR has continually increased in staphylococci, as well as in other types of bacteria. This is attributed to factors such as the frequent antibiotic administration to livestock, or misuse and overuse of antimicrobials. Therefore, the food chain constitutes one possible route of AMR transfer (Angulo et al., 2004). Both CoPS and CoNS from food origin exhibit a wide range of AMR genetic elements coding for resistance to different antimicrobials such as methicillin, tetracycline, penicillin, and vancomycin (Gundogan et al., 2005; Simeoni et al., 2008). The most frequently encountered resistant staphylococcus is methicillin-resistant *Staphylococcus aureus* (MRSA), which is resistant to multiple antimicrobials and is responsible for hard-to-treat infections. Thus, it constitutes a serious public health concern, as it causes thousands of deaths per year worldwide and constitutes a global economic burden.

The study aimed to screen the environmental heterogeneity of *Staphylococcus* species from alkaline fermented foods, their relationship with clinical isolates worldwide and their safety with regards to genes encoding antimicrobial resistance

(AMR) and toxin production. This is anticipated to elucidate their distribution and importance in such foods, especially in relation to the safety of the consumers.

2. Materials and Methods

2.1 Identification of the bacteria

2.1.1 Phenotypic characterisation

The bacteria (400) were isolated from different samples of Bikalga (9), Ntoba Mbodi (8) and Soumbala (10) collected at different markets and production sites in Burkina Faso (Bikalga and Soumbala) and the Republic of the Congo (Ntoba Mbodi), from areas constituting major consumers of these food products. Each sample (10 g) was homogenised for 1 min in 90 ml of maximum recovery diluent (MRD, Oxoid) and ten-fold dilutions (10^{-2} - 10^{10}) prepared. The dilutions (0.1 ml) were spread on nutrient agar (NA; Oxoid CM0003, Basingstoke, UK) for the enumeration of total aerobic mesophilic bacteria. The plates were incubated aerobically at 37°C and the count recorded after 2 days. For enumeration and isolation of the presumptive *Staphylococcus* population, the dilutions were spread on Baird Parker (BPA; Oxoid) and Mannitol Salt (MSA; Oxoid) agars and the plates incubated aerobically at 37°C for 2 days. Bacteria isolated from the two the latter media were purified by repeated streaking on NA, maintained in nutrient broth (NB; Oxoid CM001) containing 20% (v/v) glycerol (Sigma G8773, Gillingham, UK) as well as on cryobeads and stored at -20°C for further analyses.

For the phenotypic characterization, the isolates were streaked on NA and examined for colony and cell morphology, as well as tested for Gram, catalase, oxidase and coagulase reactions. The Gram reaction was evaluated using the KOH method (Gregersen, 1978) and the oxidase test using *Pseudomas fragi* as positive control and *Brochothrix thermosphacta* as negative control. The coagulase test was

performed using the Staphylase test kit (Oxoid DR0595) and following the manufacturer's instructions. The cell morphology was determined by light microscopy (Nikon Model Eclipse, E400, Japan). The phenotypic screening resulted in the elimination from the study of 310 non-staphylococcal isolates which were able to grow on BPA and MSA and whose colony morphologies initially suggested they were potential *Staphylococcus* spp. Thus, 90 isolates were retained for further characterization.

2.1.2 Molecular typing

2.1.2.1 DNA extraction

The bacteria were grown on Brain Heart Infusion Agar (BHI-A) prepared by mixing BHI (Oxoid CM1135) and technical agar (Oxoid LP0013, 15g/L). After incubation at 37°C for 48 h, the DNA was extracted from purified cultures following the manufacturer's instructions using InstaGene Matrix (Bio-Rad 732-6030, Hemel Hempstead, UK).

2.1.2.2 Repetitive element palindromic-polymerase chain reaction (rep-PCR)

The isolates were characterized by rep-PCR using the GTG5 primer as described by Ouoba et al. (2008a). Bacteria showing the same DNA profile were clustered in the same group by visual screening and cluster analysis using the Bionumerics system (2.50, Dice's Coefficient of similarity, Unweighted pair group method with arithmetic: UPGMA; Applied Maths, Saint-Martens-Latem, Belgium).

2.1.2.3 Pulse Field Gel Electrophoresis (PFGE)

In order to assess the worldwide distribution of the types of *S. aureus* found in the fermented foods, PFGE of SmaI-digested genomic DNA was performed according to the method proposed by the Center for Disease Control and Prevention/Pulse Net (CDC, USA) as described by Van Balen et al. (2013). The *Salmonella* serotype

Branderup strain H9812 digested with XbaI was used as a molecular size marker. The PFGE profiles of the *Staphylococcus* isolates were compared with those in a CDC database that contains worldwide profiles of about 12,000 *S. aureus* strains recovered from outbreaks, surveillance, reference and the environment, and submitted to the database.

2.1.2.4 Sequencing of the 16S rRNA and rpoB genes

The bacteria were identified by the amplification and sequencing of the 16S rRNA and *rpoB* genes. For the 16S rRNA gene amplification, the method and the pA and pE primers described by Ouoba et al. (2008a) were used. For the amplification of the *rpoB* gene, primers 2491F and 3554R described by Drancourt and Raoult (2002) were used with the following PCR mixture: 5 µl of 10 X PCR buffer containing 15 mM of MgCl₂ (Applied Biosystems N8080160), 0.5 µl of dNTP (1.25 mM), 0.5 µl of each primer (21 pmol/µl), 0.2 µl of AmpliTaq polymerase (5 U; Applied Biosystems N808-0160) and 41.3 µl of sterile high purity water. The PCR conditions were as follows: initial denaturation at 94°C for 2 min, 35 cycles of 94°C for 30 s, 52°C for 30 s, 72°C for 1 min and a final extension step at 72°C for 5 min.

After the amplification of both genes, positive amplicons were checked by electrophoresis and purified using the QIAquick PCR purification kit (Qiagen, Crawley, UK). The 16S rRNA and *rpoB* genes were sequenced (SourceBioscience, Cambridge, UK) using primers pD (3.2 pmol/µl) and 2491F/ 3554R (3.2 pmol/µl) respectively. The 16S rRNA gene sequences were analysed in the ezbiocloud server (Yoon et al., 2017) whereas those of the *rpoB* gene were screened in the GenBank/EMBL/DDBJ database (National Center for Biotechnology, MD, USA).

2.1.2.4 Spa typing

A multiplex PCR method and primers recommended by the European Reference Laboratory for antimicrobial resistance (EURL-AR, 2012), and that allows at the same time the detection of the *spa* gene and the *S. aureus* methicillin resistance genes *mecA*, *mecC* was used. The method includes the use of a PCR mixture containing 2.5 µl of 2xGreen PCR Master Mix (Thermo Scientific K1081), 2 µl (10 µM) of a mixture of the forward primers, 2 µl (10 µM) of a mixture of the reverse primers and 6.5 µl of high purity water. The PCR conditions described by Stegger et al. (2012) were applied.

2.2 Screening haemolysis and toxin production potential of the bacteria

2.2.1 Haemolytic activity on blood agar

The *Staphylococcus* isolates were screened for their haemolytic activity on blood agar as follows: Columbia agar base (Oxoid CM003) was autoclaved at 121 °C for 15 min and sheep blood (5%; Oxoid SR0051B) added after cooling to 50 °C, before distribution into Petri dishes. The isolates were then streaked on the agar and incubated at 37 °C for 48 h. Haemolysis was identified by the appearance of a zone of clearing around the colonies.

2.2.2 Production of staphylococcal enterotoxins A, B, C and D

The ability of the bacteria to produce the staphylococcal enterotoxin A, B, C and D was investigated using the staphylococcal enterotoxin reversed passive latex agglutination (SET-RPLA) toxin detection kit (Oxoid TD0900). Each isolate was grown at 37°C for 24h in 10 ml of Tryptone Soya broth (TSB; Oxoid CM0129) and 1ml of the culture centrifuged using a benchtop centrifuge at 3000 rpm for 20 min. The presence of toxins was screened in the supernatant according to the manufacturer's instructions.

2.2.3 Detection of staphylococcal enterotoxin, exfoliative and toxic shock syndrome toxin genes

The isolates were tested for the presence of the staphylococcal enterotoxin genes *sea*, *seb*, *sec*, *sed*, *see*, *seg*, *seh*, *sei*, *sej*, exfoliative toxin genes *eta* and *etb*, and toxic shock syndrome toxin gene *tsst-1*. Primers described by Johnson et al. (1991), Omoe et al. (2002) and Nashev et al. (2004) were used. The PCR mixture was that used by Ouoba et al. (2008b). The PCR conditions were as follows: 94°C for 2 min followed by 35 cycles of denaturation at 94°C for 1 min, annealing at the corresponding temperature for 1 min and elongation at 72°C for 1 min. The PCR ended with a final extension at 72°C for 5 min.

2.3 Screening the AMR phenotypic and genotypic profiles of the bacteria

2.3.1 Inoculum preparation

The isolates were grown on BHI-A and pure colonies re-suspended in 1 ml of Maximum Recovery Diluent (MRD; Oxoid CM0733). A quantity of the latter suspension was transferred to 10 ml of MRD to obtain a turbidity of 0.5 (10^7 - 10^8 CFU/ml) on the McFarland scale using a nephelometer (TREK Diagnostic systems, West Sussex, UK).

2.3.2 Determination of the susceptibility of the bacteria to antimicrobials

The Minimal Inhibitory Concentration (MIC) for 19 antimicrobials (Table 1) was determined using a 96 well sensititre™ plate (EUST, TREK Diagnostic Systems Ltd, East Grinstead, UK) containing varying amounts of each antimicrobial. The method recommended by EURL-AR (DTU, Denmark) was applied (EURL-AR, 2009). Briefly, 50 µl of the inoculum was mixed with 10 ml Mueller Hinton broth (MHB; Oxoid CM0405) and 50 µl of this mixture were dispensed in each well. Incubation was carried out aerobically at 37°C for 24 h and visible growth recorded. The susceptibility of the isolates to each antimicrobial was established using the cut-off values recorded in Table 1.

2.3.3 Detection of AMR genes

The presence of 38 genes encoding resistance to chloramphenicol, erythromycin, gentamycin, methicillin, penicillin, quinupristin/dalfopristin, tetracycline, trimethoprim, streptomycin and vancomycin were investigated (Table 1).

For the detection of the genes, the PCR mixtures described by Naghizadeh Raeisi et al. (2018) were used. However, for the detection of the *S. aureus* methicillin resistance genes *mecA* and *mecC*, the multiplex PCR (described in 2.1.2.4) and primers recommended by EURL-AR (2012) were used. For the *mecA1* gene pertaining to *S. sciuri*, the same mixture (except the primers) as that used for the detection of erythromycin (Naghizadeh Raeisi et al., 2018) was applied. The references of the primers are depicted in Table 1. Positive control isolates (EURL-AR) for most genes screened were included. Moreover, all positive amplicons were sequenced to confirm the identity of the genes.

All PCR amplifications were performed in a thermocycler (GeneAmp PCR 2700 system) using the following program: initial denaturation at 94°C for 3 min, followed by 25 - 35 cycles of 94°C for 1 min, 45-68°C for 1 min depending on the annealing temperature for the individual set of primers, 72 °C for 1 min and a final extension at 72 °C for 10 min. After purification, the presumptive positive amplicons were sequenced using the amplification primers at a concentration of 3.2 pmol/μl and the similarity of the sequence tested against published sequences in GenBank.

2.3.4 Screening the transferability of AMR

The ability of *S. warneri* 105 to transfer the *msr(A)* gene to *Enterococcus faecalis* JH2-2 and *S. saprophyticus* A17M105 to transfer the *tet(K)* gene to the same recipient was investigated by conjugation experiments according to Ouoba *et al.* (2008c) and Naghizadeh Raeisi et al. (2018). The donors are resistant to erythromycin

(*S. warneri*) and tetracycline (*S. saprophyticus*), but sensitive to rifampicin and fusidic acid. The recipient *E. faecalis* JH2-2 is resistant to rifampicin and fusidic acid and sensitive erythromycin and tetracycline.

Inocula were prepared as described above and 10 ml of the donor suspension and 1 ml of that of the recipient mixed and filtered through a sterile membrane filter (0.45 µm) (Whatman Laboratory Division, Maidstone, UK) using a vacuum pump (Welch Thomas 2522C-02, Skokie, Illinois, USA). The filters were incubated aerobically on BHI-A at 37 °C for 48h. The growth was washed off the filter with 3 ml of MRD, diluted (up to 10⁻⁵) and 100 µl of each dilution spread on agar plates containing different antimicrobials. For the *msr(A)* gene, the inoculation was made on BHI-RFE [rifampicin (25µg/ml), fusidic acid (25 µg/ml) and erythromycin (16 µg/ml)], BHI-RF [rifampicin (25 µg/ml) and fusidic acid (25 µg/ ml)], BHI-E [Erythromycin (16 µg/ml)] and BHI agar plates. For the tet(K) gene transfer, the mating mixtures were spread on BHI-RFT [rifampicin (25µg/ml), fusidic acid (25 µg/ml) and tetracycline (10 µg/ml)], BHI-RF [rifampicin (25 µg/ml) and fusidic acid (25 µg/ ml)], BHI-T [(tetracycline (10 µg/ml)] and BHI agar plates. The plates were incubated at 37 °C for 7-21 days. Cells of the donors and recipient that had not undergone a conjugation procedure were all spread on the same type of media as those used to inoculate the mating mixtures.

Only recipients that had acquired resistance to erythromycin and tetracycline are able to grow on BHI-RFE and BHI-RFT agar plates, respectively. Potential *E. faecalis* JH2-2 transconjugant (PET) DNA fingerprints were compared with those of the recipients and donors by rep-PCR using the GTG5 primer (Ouoba et al., 2008a). In addition, their MICs for erythromycin and tetracycline were determined.

2.3.5 Determination of the genetic background of acquired erythromycin and tetracycline resistance in the PETs

2.3.5.1 Extraction of DNA

Total DNA was extracted as described above. Plasmid DNA was extracted as follows: the donors, recipient and PETs were cultured for 48 h on BHI-A. A pure colony was then transferred into 10 ml BHI and incubated in a shaking water bath at 37°C for 12 h. Plasmid DNA was extracted using the QIAGEN Plasmid miniprep Kit (QIAGEN GmbH, Hilden, Germany) according to the manufacturer's instructions.

2.3.5.2 Amplification of the AMR genes in the PETs

PCR using total and plasmid DNA were performed to detect the presence of the *msr(A)* and *tet(K)* genes in the PETs. Since the *S. warneri* isolate contained also the *cat501*, *blazA* and *dfr(A)* genes, these were also screened in the PETs. The use of plasmid DNA aimed to determine the location of genes in the bacteria and to increase the chance to detect a potential transfer in the PETs.

3. Results

3.1 Identification of the bacteria

The pH values of the products were 5-6 in the unfermented samples and 8-9 in the fermented products. The bacterial numbers recovered (CFU/g) for each product were as follows: Soumbala: 10^6 - 10^{13} CFU/g on NA, 10^3 - 10^{10} on MSA, 10^1 - 10^8 on BP; Bikalga: 10^8 - 10^{11} on NA, 10^7 - 10^8 on MSA, 10^4 - 10^8 on BP; Ntoba Mbodi: 10^9 - 10^{11} on NA, 10^8 - 10^{10} on MSA, 10^5 - 10^8 on BP. Out of 27 samples analysed, *Staphylococcus* species were recovered from 18 samples. The products where *Staphylococcus* isolates were not detected included mainly samples from the production steps of Bikalga and Soumbala that had undergone heat treatment, but also one sample of each of the three foods obtained from different markets. As mentioned earlier, many bacteria recovered on MSA and BP did not possess typical staphylococcal characteristics and were not included in the advanced screening. Most of those

bacteria were Gram-positive (many of which were spore-forming) and Gram-negative rods that were able to grow on the selective media (BPA and MSA). The 90 isolates that were further screened showed variable colony and cell morphology. All were cocci, arranged differently (single, pairs, clusters) according to the isolate, catalase and Gram positive and oxidase negative. Two isolates from Bikalga were coagulase positive and the remaining 88 bacteria were coagulase negative.

The microbial profile was variable according to the fermented product, the production site and country (Table 2, Table 3). Genotyping by rep-PCR and PFGE both generated 25 clusters representing 15 species (Table 2, Fig 1). The comparison of the PFGE profiles of the *S. aureus* isolates did not generate a similarity with any of the 12,000 profiles in the CDC database. By sequencing the 16S rRNA and *rpoB* genes, the bacteria of eight rep-PCR/PFGE clusters were identified as *S. sciuri* (45.6 % of the total isolates), three as *S. gallinarum* (7.8 %), two as *S. cohnii* (4.4%), two as *S. warneri* (5.6%) and one cluster each as *S. arlettae* (2.2%), *S. aureus* (2.2%), *S. epidermidis* (1.1%), *S. haemolyticus* (1.1%), *S. hominis* (1.1%), *S. pasteurii* (1.1%), *S. condiment* (1.1%), *S. piscifermentans* (1.1%), *S. saprophyticus* (1.1%), *S. simulans* (18.9%), and *Macrococcus caseolyticus* (previously designated as *S. caseolyticus*, 5.5%). In general, *S. sciuri* was the most frequently isolated species and was dominant in the samples of Bikalga (42.9%) and particularly Ntoba Mbodi (71.1%) where it occurred from the beginning of the fermentation and persisted throughout the process. In Soumbala, *S. simulans* (54.8%) was the dominant species. The isolate of *S. haemolyticus* was clearly identified by 16S rRNA gene sequencing/EZtaxon analysis with 100 % similarity, but the *rpoB* gene sequencing generated a 96.5% (30 base pairs difference) similarity with *S. hominis* and 93.5% (30 base pairs difference) with *S. haemolyticus*. The *spa* type of the *S. aureus* isolate was identified as t355.

The species of *S. sciuri* and *S. gallinarum* were common to the two countries. One cluster (5) of *S. sciuri* was detected in all three types of food although this particular cluster was not present in all samples (Table 3). Clusters 4 and 6 of the same species were detected in samples from both countries but not in all three products. *Staphylococcus gallinarum* was also recovered from both countries but not all products, with one cluster (15) observed in Soumbala and Ntoba Mbodi. Nine species were specific to Burkina Faso including five isolated solely from Soumbala (*S. epidermidis*, *S. pasteurii*, *S. condimenti*, *S. piscifermentans*, *S. simulans*) and four from Bikalga (*S. aureus*, *S. cohnii*, *S. haemolyticus*, *M. caseolyticus*). On the other hand, four species were peculiar to Congo/ Ntoba Mbodi (*S. arlettae*, *S. hominis*, *S. saprophyticus*, *S. warneri*). Moreover, two clusters of *S. gallinarum* (1, 23) and two of *S. sciuri* (2, 7) were peculiar to Soumbala while one cluster of *S. sciuri* was particular to Bikalga.

3.2 Haemolysis on blood agar and toxin production

Haemolytic activity of the isolates on sheep blood agar was variable as seen in Table 4. Positive haemolysis was observed mainly in isolates of *S. aureus* and *S. simulans*. *Staphylococcus epidermidis*, *S. haemolyticus*, *S. pasteurii* and *S. warneri* exhibited partial haemolysis, whereas the rest of the species did not show haemolytic activity within the 48 h of incubation. The different species clusters showed similar haemolysis potential.

None of the bacteria screened produced the staphylococcal enterotoxins A, B, C and D. Furthermore, the genes coding for the pre-cited enterotoxins as well as the other eight genes were not detected in the isolates.

3.3 Antimicrobial resistance

All bacteria were susceptible to gentamycin, kanamycin, streptomycin, and vancomycin. For the rest of the antimicrobials, susceptibility was variable according to the species, the cluster, and the antimicrobial screened (Table 4). Most of the staphylococci were resistant to trimethoprim with 87.8 % of the isolates, representing 11 species, exhibiting phenotypic resistance to that antimicrobial. Both *S. cohnii* clusters exhibited the highest phenotypic resistance profiles involving 11 antimicrobials. The bacteria with the least phenotypic resistance were *S. condimentii* and *S. pasteurii*, which showed reduced susceptibility toward sulfamethoxazole only. Within the same species, the different clusters exhibited different resistance patterns, but all isolates from the same cluster showed a similar pattern irrespective of the origin. For example, isolates of cluster 1 of *S. gallinarum* were resistant to ceftiofur, clindamycin, quinupristin/dalfopristin and tiamulin, whereas those of cluster 15 were resistant to clindamycin, fusidate, penicillin, quinupristin/dalfopristin, tiamulin, sulfamethoxazole, and trimethoprim, resulting in phenotypic resistance profiles which differ by five antimicrobials (Table 4).

None of the 38 AMR genes screened was detected in the isolates of *S. condimentii*, *S. pasteurii*, *S. piscifermentans*, *S. simulans*, and *M. caseolyticus* (Table 4). For the other species, different genes encoding resistance to multiple antimicrobials including penicillin: *bla*_zA, chloramphenicol: *cat*501, trimethoprim: *dfr*(A), *dfr*(G), erythromycin: *msr*(A), methicillin: *mecA*, *mecA1* and tetracycline: *tet* (K) were detected, with a degree of AMR genotypic profile variability observed between some species and clusters of the same species (Table 4). The *mecA1* gene was detected in 45.6% of the isolates, *mecA* in 20%, *tet* (K) in 11.1 %, *bla*_zA in 10 %, *dfr*(A) in 10%, *msr*(A) in 6.7%, *cat*501 in 1.1%, and *dfr*(G) in 1.1 %. Isolates of all six clusters of *S. sciuri* showed *mecA1*, but bacteria in only three clusters displayed *mecA*. None

of the *S. aureus* isolates contained *mecA*, *mecA1* or *mecC*. The isolates of *S. epidermidis* and *S. warneri* showed the highest numbers (4) of AMR genes, including *blazA*, *dfr(A)*, *dfr(G)* and *tet(K)* for *S. epidermidis* and *blazA*, *Cat501*, *dfr(A)* and *msr(A)* for cluster 13 of *S. warneri*. The *blazA*, *mecA*, *mecA1* and *tet (K)* genes were found in isolates from all products and all countries, *cat501* in Ntoba Mbodi /Congo, *dfr(A)* in Ntoba Mbodi/Congo and Soumbala/Burkina Faso, *dfr(G)* in Soumbala/Burkina Faso, and *msr(A)* in Ntoba Mbodi /Congo and Bikalga/Burkina Faso

Conjugation experiments produced *E. faecalis* JH2-2 PETs with increased resistance to erythromycin and tetracycline. The MIC of some PETs increased up to 6 and 4 fold for erythromycin and tetracycline respectively (Table 5). All PETs showed the same rep-PCR profiles as that of the recipient *E. faecalis* JH2-2 (results not shown). No bacteria were recovered on the antimicrobial selective agar plates inoculated with cells which did not undergo a conjugation process. Positive amplicons were obtained for the genes screened for transfer using both total and plasmid DNA samples from the donors. However, no positive amplicon was observed with DNA samples from the recipient and the PETs under the experimental conditions used.

4. Discussion

The specificity in the bacterial profile of the products can be attributed to differences in the raw materials and production processes. Also, environmental factors can affect bacterial populations in a given geographical area and its associated plants and thus can affect the bacterial population of the products derived from such plants. However, since a similar type of fermentation occurs during production of the fermented foods, similarities can be anticipated. High bacterial counts as observed in our study have been reported in other studies describing the microbial composition of alkaline fermented products (Ahaotu et al., 2013; Azokpota et al., 2006; Dakwa et al.,

2005; Ouoba et al., 2010). Furthermore, the increase of pH constitutes a selection of analogous alkali resistant microorganisms in the foods. The presence of the same clusters of *S. sciuri* and *S. gallinarum* in both countries suggests a wide distribution of those strains in the environment. A peculiarity of the *S. sciuri* isolates recovered from the three products is that they were all oxidase negative. This seems unusual because the *S. sciuri* group of species including *S. sciuri*, *S. fleuritti*, *S. lentus*, *S. stepanovicii*, and *S. vitulinus* are known as the only oxidase positive species of the *Staphylococcus* genus (Becker et al., 2014). The discrepancy could be related to the origin specificity and to a lesser extent, the screening methods. The fact that the *S. aureus* PFGE profile did not match that of any clinical and environmental isolates in the CDC database suggests that the cluster is uncommon and not globally distributed.

The species of *S. sciuri*, *S. gallinarum*, *S. epidermidis*, *S. aureus* and *M. caseolyticus* have been reported in other African alkaline fermented products such as maari, dawadawa, and soydawadawa (Parkouda et al., 2009). However, most of the other species, as well as the high species and strain diversity in the *Staphylococcus* profiles observed in the current study, seem to have not been reported before. The reason could be that *Staphylococcus* populations in such foods are usually screened as part of the general microbial population and are not addressed specifically and thoroughly using both phenotypic and genotypic methods. The species of *S. condimenti* and *S. piscifermentans* have been reported to be typically associated with fermented foods such as fermented fish, sausages and their starter cultures (Becker et al., 2014). It is believed that *Staphylococcus* spp. from fermented foods are contaminants from the raw materials, equipment, environment and possibly the producers. To produce Ntoba Mbodi, no heating step is included in the process, thus the original *Staphylococcus* isolates, derived from the cassava and wrapping leaves,

that can withstand alkaline conditions probably constitute the major part of the staphylococcal population. However, production of Soumbala and Bikalga includes long cooking steps (24-48h), so it can be assumed that there is a recontamination of the products from the processing and storage materials, the producers and the environment, especially during the drying process. The limited recovery of *Staphylococcus* isolates (90/400 presumptive isolates) supports their status as a minor component of the microflora of the foods. They have often been isolated from alkaline fermented foods as secondary microorganisms, but a clear indication of their role in the fermentation is not established. The main microorganisms responsible for alkaline fermentation, and which bring about desired changes to the foods, have been demonstrated to be *Bacillus* species (Odunfa and Adewuyi, 1985; Ouoba et al., 2017). Fermentation trials conducted by Odunfa and Adewuyi (1985) showed that *S. saprophyticus* was unable to ferment African locust bean seeds to generate a product with the typical organoleptic and nutritional properties. Nevertheless, it can be suggested that the presence of *Staphylococcus* species contributes to some biochemical activities such as those related to the degradation of proteins and lipids that some species are able to perform during meat fermentation to produce sausages (Casaburi et al., 2008; Milicevic et al., 2014).

Staphylococcus spp., especially CoPS, are known to cause various illnesses including food intoxication due to their capacity to produce heat resistant enterotoxins. Interestingly, none of the isolates investigated, including the two CoP *S. aureus*, exhibited a potential toxin-producing phenotype and genotype. It cannot be ruled out that they may contain other toxin genes that were not screened, but since the most common genes were not detected, it can be assumed that these bacteria do not represent a serious threat for consumers. In Burkina Faso and the Republic of the

Congo, there are no official reports of foodborne illnesses resulting from the consumption of alkaline traditional fermented foods, although this may be related to non-availability of data and lack of statistics. If unrecorded cases had occurred, these would probably have been attributed to toxins elaborated by the fermenting *B. cereus* strains that commonly occur in the products (Ouoba et al., 2008 b; Voudibio-Mbozo et al., 2017). In the study by Fowoyo and Ogunbanwo (2017a), the *sea*, *seb*, *sec* and *sed*, and *tsst-1* genes were detected in CoNS, including *S. epidermidis*, *S. simulans*, *S. xylosus*, *S. kloosii* and *S. caprae*, isolated from six Nigerian traditional fermented foods including iru, an alkaline fermented African locust bean product that is similar to Soumbala. The positive amplicons were not sequenced to confirm the identity of the gene, but the ability of some isolates to produce the staphylococcal toxins A, B, C, and D suggest that these bacteria may pose a potential threat to consumers.

Antimicrobial resistances that are not associated with the presence of a gene are of concern, but those involving a gene represent an enhanced threat, due to the potential for transfer to other bacteria by mechanisms such as horizontal transfer, and therefore can be spread. The *tet* (K), *blazA*, *dfr*(A), *msr*(A), *cat501*, and *dfr*(G) genes detected can be associated with transferable mobile genetic elements such as plasmids and transposons, and can therefore potentially be transferred to other bacteria including pathogenic microorganisms. In the current research, *tet* (K), *blazA*, *msr*(A), *cat501* and *dfr*(A) genes were amplified using plasmid DNA, suggesting that they are located on plasmids, at least in the isolates screened. The AMR threat is greater with bacteria with multi-drug genotypic profiles, such as one of the isolates of *S. warneri* which exhibited AMR genes for four antimicrobials: erythromycin, chloramphenicol, penicillin and trimethoprim, and those of *S. epidermidis* and *S.*

hominis which showed respectively four and three genes encoding resistance to penicillin, tetracycline and chloramphenicol.

Methicillin resistance is an important factor when screening AMR in *Staphylococcus*. This is because MRSA is responsible for a serious and persistent problem in treatment of infections caused by such bacteria. The most virulent MRSA are also resistant to other beta-lactam antimicrobials. Methicillin resistance is usually caused by the presence of the *mecA* gene coding for a penicillin-binding protein (PBP2a) associated with a significantly lower affinity for beta-lactams. Thus, cell-wall biosynthesis by the bacteria continues, even when a typically inhibitory concentration of antibiotic is administered (Peacock et al., 2015). Variants of *mecA* include genes such as *mecA* and *mecC* (*mecA*_{ALGA251}) originally found in *S. aureus* isolates, *mecA1*, *mecA2* and *mecB* (*mecAm*) initially detected in the *S. sciuri* group, *S. vitulinus* and *M. caseolyticus* respectively (Ito et al., 2012). Isolates of three clusters of *S. sciuri* from both countries exhibited both *mecA1* and *mecA*, whereas the isolates of the three other clusters also found in both countries exhibited only *mecA1*. The presence of the *mecA1* gene in *S. sciuri* is a common trait of this species and does not generally imply methicillin resistance in the bacteria (Couto et al., 2000). However, the presence of *mecA* in the isolates implicates methicillin resistance and represents a threat, although a phenotypic resistance was not observed for cefoxitin. Couto et al. (2000) showed that only *S. sciuri* isolates that possess a copy of *mecA* in addition to *mecA1* exhibit significant levels of resistance to beta-lactams. In our study, such isolates displayed phenotypic resistance to penicillin. Interestingly, the *S. aureus* did not display any of the methicillin resistance genes screened and were susceptible to cefoxitin. However, the strains were resistant to penicillin and contained the *blaZ* gene.

The prevalence of AMR genes in various CoNS and CoPS from processed and unprocessed foods such as meat, fish and fermented dairy and meat products from different origins has been widely documented (Argudín et al., 2011; Kürekci, 2016; Perreten et al., 1998; Zdolec et al., 2012). However, such information on CoNS and CoPS from alkaline fermented products and especially those from Africa are, to the best of our knowledge, non-existent. This can be due to the limited importance attributed to these bacteria in alkaline fermentation. The study by Fowoyo and Ogunbanwo (2017b) on CoNS from different Nigerian fermented foods, reported phenotypic resistance to antimicrobials such as trimethoprim-sulfamethoxazole, ciprofloxacin, erythromycin, gentamicin, and cefotaxime, but no AMR gene was reported, probably because only *mecA* was screened in the isolates with no positive results.

The significant increase of the MIC above the cut-off values in the PETs for erythromycin and tetracycline indicates that they have acquired resistance to the antimicrobials. Since the genes screened were not detected in the PETs, it can be assumed that they were not transferred, but that the isolates acquired resistance through other mechanisms, such as mutations. The occurrence of resistances was triggered by the presence of the *Staphylococcus* donors because PETs were not recovered from recipient cells which did not undergo a conjugation process. Mutations can occur as a result of a mechanism by which the *Staphylococcus* spp. stimulate an evolution of *E. faecalis* JH2-2 by increasing, for instance, the expression of error-prone DNA repair mechanisms (Naghizadeh Raeisi, 2018). Alternatively, the growth of the recipient could be stimulated by the donor leading to a larger recipient population in which mutations can arise (Naghizadeh Raeisi, 2018). However, it is worth noting that when a low number of plasmid copies which carry a gene is transferred, it may not be

detected using conventional PCR. Moreover, other genes coding for resistance to the antimicrobials and which were not screened, may be present in the donors and transferred to the recipients. Transfer potential varies according to different factors related to the environment, type of donor and recipient, and the location of the AMR genes. Thus, the genes screened may have not been transferred into *E. faecalis* JH2-2 and therefore the threat with regards to the potential of the donors to spread AMR is limited. However, the possibility of gene transfer to other types of bacteria cannot be ruled out since the genes are located on plasmids which are mobile genetic elements that can be transferred in appropriate conditions.

The current study demonstrates ecological similarities and differences in the *Staphylococcus* population of different types of alkaline fermented foods. The main microorganisms identified are primarily CoNS with a very limited number of CoPS, as would be expected in non-clinical samples. The bacteria did not suggest a safety issue with regards to the potential to produce toxins responsible for foodborne diseases. Due to the presence of multiple AMR genes in some isolates, they are reservoirs of AMR determinants which may be spread under particular conditions. However, in general, the results of the study herein reported suggest little safety importance of the staphylococcus isolates in the fermented products screened.

Acknowledgements

The authors are grateful to the following people for their technical and scientific support in the realization of the research project: Professor Simon C. Kobawila, and Professor Delphin Louembe (Marien N’Gouabi University, Brazzaville, Congo), Dr Brigitte Awamaria (London Metropolitan University, London, UK), Dr Rene S. Hendriksen, Susanne Karlslose Pedersen, Ana Rita Rebelo, and Dr. Lars Bøgo Jensen

(Technical University of Denmark, Lyngby, Denmark), Janne M. Benjaminsen (University of Copenhagen, Copenhagen, Denmark), Dr Eva Møller Nielsen, and Dr Jesper Larsen (Statens Serum Institut, Copenhagen, Denmark), Dr Molly Freeman, Dr Jeniffer Concepcion Acevedo, and Valerie S. Albrecht (Centers for Disease Control and Prevention, Atlanta, GA, USA)

Declaration of interest: None

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors

References

- Ahaotu, I., Anyogu, A., Njoku, O.H., Odu, N.N., Sutherland, J.P., Ouoba, L.I.I., 2013. Molecular identification and safety of *Bacillus* species involved in the fermentation of African oil beans (*Pentaclethra macrophylla* Benth) for production of Ugba. *Int. J. Food Microbiol.* 162, 95-104.
- Angulo, F., Nargund, V., Chiller, T., 2004. Evidence of an Association Between Use of Anti-microbial Agents in Food Animals and Anti-microbial Resistance Among Bacteria Isolated from Humans and the Human Health Consequences of Such Resistance. *J. Vet. Med. B.* 51, 374-379.
- Argudín, M., Tenhagen, B., Fetsch, A., Sachsenröder, J., Käsbohrer, A., Schroeter, A., Hammerl, J.A., Hertwig, S., Helmuth, R., Bräunig, J., Mendoza, M.C., Appel, B., Rodicio, M.R., Guerra, B., 2011. Virulence and Resistance Determinants of German *Staphylococcus aureus* ST398 Isolates from Non-human Sources. *Appl. Environ. Microbiol.* 77, 3052-3060.

623 Azokpota, P., Hounhouigan, D.J., Nago, M.C., 2006. Microbiological and chemical
 624 changes during the fermentation of African locust bean (*Parkia biglobosa*) to produce
 625 *afitin*, *iru* and *sonru*, three traditional condiments produced in Benin. *Int. J. Food*
 626 *Microbiol.* 107, 304 – 309.

627 Becker, K., Heilmann, C., Peters, G., 2014 Coagulase-Negative Staphylococci. *Clin.*
 628 *Microbiol. Rev.* 27, 870-926.

629 Breckinridge, J., Bergdoll, M., 1971. Outbreak of Food-Borne Gastroenteritis Due to a
 630 Coagulase-Negative Enterotoxin-Producing *Staphylococcus*. *N. Engl. J. Med.* 284,
 631 541-543.

632 Casaburi, A., Di Monaco, R., Cavella, S., Toldrá, F., Ercolini, D., Villani, F., 2008.
 633 Proteolytic and lipolytic starter cultures and their effect on traditional fermented
 634 sausages ripening and sensory traits. *Food Microbiol.* 25, 335-347.

635 Couto, I., Santos Sanches, I., Sá-Leão, R., de Lencastre, H., 2000. Molecular
 636 characterization of *Staphylococcus sciuri* strains isolated from humans. *J. Clin.*
 637 *Microbiol.* 38, 136-143.

638 Crass, B., Bergdoll, M., 1986. Involvement of coagulase-negative staphylococci in
 639 toxic shock syndrome. *J. Clin. Microbiol.* 23, 43-45.

640 Dakwa, S., Sakyi-Dawson, E., Diako, C., Annan, N.T., Amoa-Awua, W.K., (2005)
 641 Effect of boiling and roasting on the fermentation of soybeans into dawadawa (soy-
 642 dawadawa). *Int. J. Food Microbiol.* 104, 69 – 82.

643 Drancourt, M., Raoult, D., 2002. *rpoB* Gene Sequence-Based Identification of
 644 *Staphylococcus* Species. *J. Clin. Microbiol.* 40, 1333-1338.

645 EUCAST., 2018. Breakpoint tables for interpretation of MICs and zone diameters.
646 [http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/v_](http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/v_8.1_Breakpoint_Tables.pdf)
647 [8.1_Breakpoint_Tables.pdf](http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/v_8.1_Breakpoint_Tables.pdf)

648 EURL-AR., 2009. Protocol for MIC-testing with sensititre of *Staphylococcus* spp.
649 [https://www.eurl-ar.eu/CustomerData/Files/Folders/21-protocols/285_1-protocol-for-](https://www.eurl-ar.eu/CustomerData/Files/Folders/21-protocols/285_1-protocol-for-mic.pdf)
650 [mic.pdf](https://www.eurl-ar.eu/CustomerData/Files/Folders/21-protocols/285_1-protocol-for-mic.pdf)

651 EURL-AR., 2013. List of primers for detection of antimicrobial resistance genes.
652 [https://www.eurl-ar.eu/CustomerData/Files/Folders/25-resourcer/333_primerliste-til-](https://www.eurl-ar.eu/CustomerData/Files/Folders/25-resourcer/333_primerliste-til-web-07-11-2013-9.pdf)
653 [web-07-11-2013-9.pdf](https://www.eurl-ar.eu/CustomerData/Files/Folders/25-resourcer/333_primerliste-til-web-07-11-2013-9.pdf)

654 EURL-AR., 2017. PROTOCOL For antimicrobial susceptibility testing of *Escherichia*
655 *coli*, enterococci and staphylococci. [https://www.eurl-](https://www.eurl-ar.eu/CustomerData/Files/Folders/23-eqas/356_protocol-eurl-ar-eqas-ec-ent-staph-2017.pdf)
656 [ar.eu/CustomerData/Files/Folders/23-eqas/356_protocol-eurl-ar-eqas-ec-ent-staph-](https://www.eurl-ar.eu/CustomerData/Files/Folders/23-eqas/356_protocol-eurl-ar-eqas-ec-ent-staph-2017.pdf)
657 [2017.pdf](https://www.eurl-ar.eu/CustomerData/Files/Folders/23-eqas/356_protocol-eurl-ar-eqas-ec-ent-staph-2017.pdf).

658 EURL-AR., 2012. Protocol for amplification of *MecA*, *MecC* (MecALGA251), *spa* and
659 *PVL*. [https://www.eurl-ar.eu/CustomerData/Files/Folders/21-protocols/279_pcr-spa-](https://www.eurl-ar.eu/CustomerData/Files/Folders/21-protocols/279_pcr-spa-pvl-meca-mecc-sept12.pdf)
660 [pvl-meca-mecc-sept12.pdf](https://www.eurl-ar.eu/CustomerData/Files/Folders/21-protocols/279_pcr-spa-pvl-meca-mecc-sept12.pdf)

661 EFSA., 2012. Technical specifications on the harmonised monitoring and reporting of
662 antimicrobial resistance in methicillin-resistant *Staphylococcus aureus* in food-
663 producing animals and food. EFSA J. 10, 2897.

664 Fonseca, S., Cachaldora, A., Gómez, M., Franco, I., Carballo, J., 2013. Monitoring the
665 bacterial population dynamics during the ripening of Galician chorizo, a traditional dry
666 fermented Spanish sausage. Food Microbiol. 33, 77-84.

667 Fowoyo, P., Ogunbanwo, S., 2017a. Virulence and toxigenicity of coagulase-negative
 668 staphylococci in Nigerian traditional fermented foods. Can. J. Microbiol. 62, 572-578.

669 Fowoyo, P., Ogunbanwo, S., 2017b. Antimicrobial resistance in coagulase-negative
 670 staphylococci from Nigerian traditional fermented foods. Ann. Clin. Microbiol.
 671 Antimicrob. 16, 4.

672 Gregersen, T., 1978. Rapid method for distinction of gram-negative from gram-positive
 673 bacteria. Eu. J. Appl. Microbiol and Biotechnol. 5, 123-127.

674 Gundogan, N., Citak, S., Yucel, N., Devren, A., 2005. A note on the incidence and
 675 antibiotic resistance of *Staphylococcus aureus* isolated from meat and chicken
 676 samples. Meat Sci. 69, 807-810.

677 Hennekinne, J., De Buyser, M., Dragacci, S., 2012. *Staphylococcus aureus* and its
 678 food poisoning toxins: characterization and outbreak investigation. FEMS Microbiol.
 679 Rev. 36, 815-836.

680 Yoon, S.H., Ha, S.M., Kwon, S., Lim, J., Kim, Y., Seo, H., Chun, J., 2017. Introducing
 681 EzBioCloud: a taxonomically united database of 16S rRNA gene sequences and
 682 whole-genome assemblies. Int. J. Syst. Evol. Microbiol. 67, 1613-1617.

683 Ito, T., Hiramatsu, K., Tomasz, A., de Lencastre, H., Perreten, V., Holden, M.,
 684 Coleman, D.C., Goering, R., Giffard, P.M., Skov, R.L., Zhang, K., Westh, H., O'Brien,
 685 F., Tenover, F.C., Oliveira, D.C., Boyle-Vavra, S., Laurent, F., Kearns, A.M.,
 686 Kreiswirth, B., Ko, K.S., Grundmann, H., Sollid, J.E., John, J.F. Jr., Daum, R.,
 687 Soderquist, B., Buist, G., IWG-SCC., 2012. Guidelines for Reporting Novel *mecA*
 688 Gene Homologues. Antimicrob. Agents Chemother. 56, 4997-4999.

689 Johnson, W., Tyler, S., Ewan, E., Ashton, F., Pollard, D., Rozee, K., 1991. Detection
 690 of Genes for Enterotoxins, Exfoliative Toxins, and Toxic Shock Syndrome Toxin 1 in
 691 *Staphylococcus aureus* by the Polymerase Chain Reaction. J. Clin. Microbiol. 29, 426-
 692 430.

693 Kiers, J.L, Van laeken, A.E., Rombouts, F.M., Nout, M.J., 2000. In vitro digestibility of
 694 Bacillus fermented soya bean. In vitro digestibility of Bacillus fermented soya bean.
 695 Int. J. Food Microbiol., 60,163-169.

696 Kurekci, C., Onen, S., Yipel, M., Aslantas, O., Gundogdu, A., 2016. Characterisation
 697 of phenotypic and genotypic antibiotic resistance profile of enterococci from cheeses
 698 in Turkey. Korean J. Food Sci. Anim. Resour. 36, 352-358.

699 LPSN., 2018 - List of prokaryotic names with standing in nomenclature [Internet].
 700 Bacterio.net. <http://www.bacterio.net/>

701 Milicevi, B., Danilovic, B., Zdolec, N., Kozachinski, L., Dobranic, V., Savic, D., 2014.
 702 Microbiota of the fermented sausages: influence to product quality and safety. Bulg. J.
 703 Agric Sci. 20, 1061-1078.

704 Naghizadeh Raeisi, S., Ghoddusi, H., Juncker Boll, E., Farahmand, N., Stuer-
 705 Lauridsen, B., Johansen, E., Sutherland, J.P., Ouoba, L.I.I., 2018. Antimicrobial
 706 susceptibility of bifidobacteria from probiotic milk products and determination of the
 707 genetic basis of tetracycline resistance in *Enterococcus* species after in vitro
 708 conjugation with *Bifidobacterium animalis subsp. lactis*. Food Control. 2018; 94: 205-
 709 211.

710 Nashev, D., Toshkova, K., Isrina, S., Salasia, S., Hassan, A., Lammler, C., Zschock,
 711 M., 2004. Distribution of virulence genes of *Staphylococcus aureus* isolated from
 712 stable nasal carriers. FEMS Microbiol. Lett. 233, 45-52.

713 Odunfa, S., Adewuyi, E., 1985. Optimization of process conditions for the fermentation
 714 of African locust bean (*Parkia biglobosa*) (Effect of starter cultures). Chem. Mikrobio.
 715 Technol. Lebensm. 9, 6-10.

716 Omoe, K., Ishikawa, M., Shimoda, Y., Hu, D., Ueda, S., Shinagawa, K., 2002.
 717 Detection of seg, seh, and sei genes in *Staphylococcus aureus* isolates and
 718 determination of the enterotoxin productivities of *S. aureus* isolates harboring seg, seh,
 719 or sei genes. J. Clin. Microbiol. 40: 857-862.

720 Ouoba, L., Parkouda, C., Diawara, B., Scotti, C., Varnam, A., 2008a. Identification of
 721 *Bacillus* spp. from Bikalga, fermented seeds of *Hibiscus sabdariffa*: phenotypic and
 722 genotypic characterization. J. Appl. Microbiol. 104, 122-131.

723 Ouoba, L., Thorsen, L., Varnam, A., 2008b. Enterotoxins and emetic toxins production
 724 by *Bacillus cereus* and other species of *Bacillus* isolated from Soumbala and Bikalga,
 725 African alkaline fermented food condiments. Int. J. Food Microbiol. 124, 224-230.

726 Ouoba, L., Lei, V., Jensen, L., 2008c. Resistance of potential probiotic lactic acid
 727 bacteria and bifidobacteria of African and European origin to antimicrobials:
 728 Determination and transferability of the resistance genes to other bacteria. Int. J. Food
 729 Microbiol. 121, 217-224.

730 Ouoba, L., Nyanga-Koumou, C.A., Parkouda, C., Sawadogo, H., Kobawila, S.C.,
 731 Keleke, S., Diawara, B., Louembe, D., Sutherland, J.P., 2010. Genotypic diversity of

732 lactic acid bacteria isolated from African traditional alkaline-fermented foods. Int. J.
 733 Food Microbiol. 108, 2019 – 2029.

734 Ouoba, L., 2015. Bikalga/Furundu/Mbuja. In: Sarkar, P., Nout, M.J.R., (Eds.),
 735 Handbook of indigenous foods involving alkaline fermentation. CRC Press, Boca
 736 Raton, pp. 129-136.

737 Ouoba, L., 2017. Traditional alkaline fermented foods: Selection of functional *Bacillus*
 738 starter cultures for Soumbala production. In: Speranza, B., Bevilacqua, A., Corbo, M.,
 739 Sinigaglia, M., (Eds), Starter Cultures in Food Production. 1st ed. Wiley Blackwell,
 740 Chichester, pp. 370-383.

741 Parkouda, C., Diawara, B., Ouoba, L., 2008. Technology and physico-chemical
 742 characteristics of Bikalga, alkaline fermented seeds of *Hibiscus sabdariffa*. Afr. J.
 743 Biotechnol. 7, 916-922.

744 Parkouda, C., Nielsen, D., Azokpota, P., Ouoba, L.I.I., Amoa-Awua, W., Thorsen, L.,
 745 Hounhouigan., J.D., Jensen, J.S, Tano-Debrah, K., Diawara, B., Jakobsen, M., 2009.
 746 The microbiology of alkaline-fermentation of indigenous seeds used as food
 747 condiments in Africa and Asia. Crit. Rev. Microbiol. 35,139-156.

748 Peacock, S., Paterson, G., 2015. Mechanisms of methicillin resistance in
 749 *Staphylococcus aureus*. Annu. Rev. Biochem. 84, 577-601.

750 Perreten, V., Giampà, N., Schuler-Schmid, U., Teuber, M., 1998. Antibiotic resistance
 751 genes in coagulase-negative staphylococci isolated from food. Syst. Appl. Microbiol.
 752 21,113-120.

753 Podkowik, M., Park, J., Seo, K., Bystróż, J., Bania, J., 2013. Enterotoxigenic potential
 754 of coagulase-negative staphylococci. Int. J. Food Microbiol. 163, 34-40.

755 Simeoni, D., Rizzotti, L., Cocconcelli, P., Gazzola, S., Dellaglio, F., Torriani, S., 2008.
 756 Antibiotic resistance genes and identification of staphylococci collected from the
 757 production chain of swine meat commodities. Food Microbiol. 25,196-201.

758 Soares Casaes Nunes, R., Mere Del Aguila, E., Paschoalin, V., 2015. Safety
 759 Evaluation of the coagulase-negative staphylococci microbiota of salami:
 760 superantigenic toxin production and antimicrobial resistance. BioMed Res. Int. 2015,
 761 1-17.

762 Stegger, M., Andersen, P., Kearns, A., Pichon, B., Holmes, M., Edwards, G., Laurent,
 763 F., Teale, C., Skov, R., Larsen, A.R., 2012. Rapid detection, differentiation and typing
 764 of methicillin-resistant *Staphylococcus aureus* harbouring either *mecA* or the new
 765 *mecA* homologue *mecALGA251*. Clin. Microbiol. Infect. 18, 395-400.

766 Sutcliffe, J., Grebe, T., Tait-Kamradt, A., Wondrack, L., 1996. Detection of
 767 erythromycin-resistant determinants by PCR. Antimicrob. Agents Chemother. 40,
 768 2562-2566.

769 Tsubakishita, S., Kuwahara-Arai, K., Sasaki, T., Hiramatsu, K., 2010. Origin and
 770 Molecular evolution of the determinant of methicillin resistance in staphylococci.
 771 Antimicrob. Agents Chemother. 10, 4352-4359

772 van Balen, J., Kelley, C., Nava-Hoet, R., Bateman, S., Hillier, A., Dyce, J., Wittum,
 773 T.E., Hoet, A. E. 2013. Presence, Distribution, and Molecular Epidemiology of
 774 Methicillin-Resistant *Staphylococcus aureus* in a Small Animal Teaching Hospital: A
 775 Year-Long Active Surveillance Targeting Dogs and Their Environment. Vector Borne
 776 Zoonotic Dis. 5, 299-311.

777 Viridis, S., Scarano, C., Cossu, F., Spanu, V., Spanu, C., De Santi, E. 2010. Antibiotic
778 resistance in *Staphylococcus aureus* and Coagulase Negative Staphylococci Isolated
779 from Goats with Subclinical Mastitis. Vet. Med. Int., 2010, 517060.

780 Voudibio Mbozo, B. A., Kobawila, S., Anyogu, A., Awamaria, B., Louembe, D.,
781 Sutherland, J., Ouoba, L.I.I., 2017. Investigation of the diversity and safety of the
782 predominant *Bacillus pumilus sensu lato* and other *Bacillus* species involved in the
783 alkaline fermentation of cassava leaves for the production of Ntoba Mbodi. Food
784 Control. 82, 154-162.

785 Zdolec, N., Dobranić, V., Zdolec, G., Đuričić, D. 2013. Antimicrobial resistance of
786 coagulase-negative staphylococci and lactic acid bacteria from industrially produced
787 dairy products. Mljekarstvo. 63, 30-35.

788

Figure caption

Figure 1: Cluster analysis (Bionumerics, Dice's Coefficient of similarity, Unweighted pair group method with arithmetic: UPGMA) of the different rep-PCR and PFGE fingerprints of *Staphylococcus species* isolated from Bikalga, Soumbala and Ntoba Mbodi

810 Table 1: Antimicrobials studied and their effect on the *Staphylococcus* isolates
811 screened

Antimicrobials	Concentration Range screened (mg/L)	Cut off values(mg/L)	Genes screened ^e	<i>Staphylococcus</i> isolates		
				MIC Range (mg/L)	Resistant (%)	Susceptible (%)
Cefoxitin	0.5 - 16	4 ^{a,b}	<i>mecA</i> , <i>mecA1</i> and <i>mecC</i> for methicillin resistance	1 - 8	8.9	91.1
Chloramphenicol	4 - 64	8 ^c	<i>cat501</i> , <i>catA1</i> , <i>cmlA</i>	<4 - 64	52.2	47.8
Ciprofloxacin	0.25 - 8	1 ^a		<0.25 - 2	1.1	98.9
Clindamycin	0.12 - 4	0.25 ^a		<0.12 - 4	41.1	58.9
Erythromycin	0.25 - 8	1 ^a	<i>erm(A)</i> , <i>erm(B)</i> , <i>erm(C)</i> , <i>erA</i> , <i>erB</i> , <i>msrA</i> , <i>mphA</i> , <i>mefA/mefE</i>	<0.25 - >8	11.1	88.9
Fusidate	0.5 - 4	1 ^c		<0.5 - >4	43.3	56.7
Gentamicin	1 - 16	2 ^a	<i>aac(6')aph(2'')</i> , <i>aac(3'')II</i> , <i>aac(3'')IV</i> , <i>ant(2'')-I</i>	<1	0	100
Kanamycin	4 - 64	16 ^d		<4	0	100
Linezolid	1 - 8	4 ^a		<1 - 8R	1.1	98.9
Mupirocin	0.5 - 256	1 ^a		<0.5 - > 256	1.1	98.9
Penicillin	0.12 - 2	0.125 ^b	<i>BlaZ</i>	<0.12 - >2	43.3	56.7
Quinupristin- dalfopristin	0.5 - 4	1 ^a	<i>VatD</i> , <i>VatE</i>	<0.5 - 4	64.4	35.6
Rifampicin	0.016 - 0.5	0.5 ^c		<0.016 - >0.5	4.4	95.6
Streptomycin	4 - 32	16 ^b	<i>StrA</i> , <i>StrB</i> , <i>aadA</i> , <i>aadE</i>	<4 - 16	0	100
Sulfamethoxazole	64 - 512	128 ^a		<64 - >512	61.1	38.9
Tetracycline	0.5 - 16	1 ^a	<i>tet(M)</i> , <i>tet(L)</i> , <i>tet(S)</i> , <i>tet(Q)</i> , <i>tet(K)</i> , <i>tet(O)</i> , <i>tet(W)</i>	<0.5 - >16	6.7	93.76
Tiamulin	0.5 - 4	2 ^a		<0.5 - > 4	48.9	51.1
Trimethoprim	2 - 32	2 ^a	<i>dfr(A)</i> , <i>dfr(G)</i> , <i>dfr(K)</i>	<2 - >32	87.8	12.2
Vancomycin	1 - 16	2 ^a	<i>Van A</i> , <i>VanB</i> , <i>VanX</i>	<1	0	100

812

813 Cut-off values according to a) EFSA (2012), b) *EUURL-AR* (2017), c) EUCAST (2018) and d) Virdis et
814 al. (2010)
815 e) Most primer sequences are from *EUURL-AR* (2013) and Ouoba et al. (2008), except for *mecA1*
816 (*Tsubakishita* et al., 2010), *erA*, *erB*, *msrA*, *mphA*, *mefA/mefE* (*Sutcliffe* et al., 1996), *dfr(G)*, *dfr(K)*
817 (*Argudín* et al., 2011)

818 f) MIC: Minimal Inhibitory Concentration

819

820

821

822

823

824 Table 2: Identity and origin of the *Staphylococcus* species recovered from Soumbala,
825 Bikalga, and Ntoba Mbodi

Samples/Origin	Fermentation time/ sampling site ^a	Isolates	^b Rep-PCR/ PFGE clusters	Identification by 16S RNA and <i>rpoB</i> gene sequencing
Soumbala				
Burkina Faso				
S3	Site 1/ 0h	S35M (47)	14	<i>Staphylococcus epidermidis</i>
S5	Site 1/ 48h	S51B (33)	23	<i>Staphylococcus gallinarum</i>
		S54a M (34)	23	<i>Staphylococcus gallinarum</i>
		S54b M (35)	23	<i>Staphylococcus gallinarum</i>
		S63aM (41)	3	<i>Staphylococcus simulans</i>
S6	Site 1/ Dried	SF1M (1)	1	<i>Staphylococcus gallinarum</i>
SF	Site 2/ ≥72h	SF3M (2)	3	<i>Staphylococcus simulans</i>
		SF2M (3)	5	<i>Staphylococcus sciuri</i>
		SF2bB (6)	3	<i>Staphylococcus simulans</i>
		SF2aB (7)	3	<i>Staphylococcus simulans</i>
		SF1B (8)	3	<i>Staphylococcus simulans</i>
		SF8M (21)	2	<i>Staphylococcus sciuri</i>
		SF7M (22)	3	<i>Staphylococcus simulans</i>
		SF6M (23)	7	<i>Staphylococcus sciuri</i>
		SP6M (26)	10	<i>Staphylococcus piscifermentans</i>
		SP5M (27)	3	<i>Staphylococcus simulans</i>
		SP3M (28)	3	<i>Staphylococcus simulans</i>
		SP1M (29)	11	<i>Staphylococcus pasteurii</i>
SP	Site 3/ ≥72h	SP4B (30)	3	<i>Staphylococcus simulans</i>
		SP2B (31)	3	<i>Staphylococcus simulans</i>
		SP1B (32)	3	<i>Staphylococcus simulans</i>
		SZ1B (10)	3	<i>Staphylococcus simulans</i>
SZ	Site 4/ ≥72h	SZ2B (11)	3	<i>Staphylococcus simulans</i>
		SZ6B (13)	3	<i>Staphylococcus simulans</i>
		SZ1M (14)	5	<i>Staphylococcus sciuri</i>
		SZ3M (15)	3	<i>Staphylococcus simulans</i>
		SZ4M (16)	5	<i>Staphylococcus sciuri</i>
		SZ5M (17)	9	<i>Staphylococcus condimentii</i>
		SZ6M (18)	15	<i>Staphylococcus gallinarum</i>
		SZ7M (24)	3	<i>Staphylococcus simulans</i>
		SZ9M (25)	3	<i>Staphylococcus simulans</i>
		Bikalga		
Burkina Faso				
B2	Site 5/ 48h	B21M (50)	6	<i>Staphylococcus sciuri</i>
		B22M (51)	4	<i>Staphylococcus sciuri</i>
		B27aM (53)	16	<i>Staphylococcus cohnii</i>
		B27bM (54)	16	<i>Staphylococcus cohnii</i>
		B25M (56)	16	<i>Staphylococcus cohnii</i>
		B25B (57)	8	<i>Staphylococcus sciuri</i>
		B24B (58)	24	<i>Staphylococcus cohnii</i>
		B35M (60)	5	<i>Staphylococcus sciuri</i>
B3	Site 5/ 72h	B36B (61)	5	<i>Staphylococcus sciuri</i>
		B34M (62)	5	<i>Staphylococcus sciuri</i>
		B32B (64)	5	<i>Staphylococcus sciuri</i>
		B31M (65)	5	<i>Staphylococcus sciuri</i>
		B31B (66)	8	<i>Staphylococcus sciuri</i>

826 ^aSite 1: Ouagadougou (Samandin production site), Site 2: Fada market, Site 3: Ouagadougou (Patte
827 d'Oie market), Site 4: Ouagadougou (Zogona market), Site 5: Ouagadougou (Zone Pilote production

828 site) ^b Rep-PCR : Repetitive element palindromic-polymerase chain reaction ; PFGE : Pulse Field Gel
829 Electrophoresis

Accepted Manuscript

Table 2 *continued*: Identity of the *Staphylococcus* species recovered from Soumbala, Bikalga, and Ntoba Mbodi

Samples/Origin	Fermentation time/ sampling site ^a	Isolates	Rep-PCR/ PFGE clusters	Identification 16S RNA and <i>rpoB</i> gene sequencing
B5	Site 5/ Dried	B57B (69)	18	<i>Staphylococcus aureus</i>
		B51B (70)	18	<i>Staphylococcus aureus</i>
BP	Site 6/ Dried	BP2B (73)	19	<i>Macrococcus caseolyticus</i>
		BP2M (74)	19	<i>Macrococcus caseolyticus</i>
		BP3B (75)	19	<i>Macrococcus caseolyticus</i>
		BP6B (76)	19	<i>Macrococcus caseolyticus</i>
		BP5B (77)	19	<i>Macrococcus caseolyticus</i>
BB Ntoba Mbodi	Site 7/ Dried	BB3B (84)	20	<i>S. haemolyticus</i>
Congo AU0	Site 8/ 0h	A04M (93)	25	<i>Staphylococcus sciuri</i>
		A05M (94)	12	<i>Staphylococcus warneri</i>
		A08M (95)	12	<i>Staphylococcus warneri</i>
		A07M (185)	12	<i>Staphylococcus warneri</i>
		A06B (193)	12	<i>Staphylococcus warneri</i>
AU1	Site 8/ 24h	A12B (96)	6	<i>Staphylococcus sciuri</i>
		A15M (104)	25	<i>Staphylococcus sciuri</i>
		A17M (105)	13	<i>Staphylococcus warneri</i>
		A18M (106)	17	<i>Staphylococcus arlettae</i>
		A15M (176)	4	<i>Staphylococcus sciuri</i>
		A16M (177)	6	<i>Staphylococcus sciuri</i>
		A111B (186)	5	<i>Staphylococcus sciuri</i>
		A110B (194)	5	<i>Staphylococcus sciuri</i>
AU2	Site 8/ 48h	A25B (111)	17	<i>Staphylococcus arlettae</i>
		A215B (113)	15	<i>Staphylococcus gallinarum</i>
		A21M (115)	4	<i>Staphylococcus sciuri</i>
		A27B (178)	5	<i>Staphylococcus sciuri</i>
		A29B (179)	6	<i>Staphylococcus sciuri</i>
		A26M (180)	6	<i>Staphylococcus sciuri</i>
		A23M (191)	4	<i>Staphylococcus sciuri</i>
		A24M (192)	4	<i>Staphylococcus sciuri</i>
AU3	Site 8/ 72h	A37bB (123)	6	<i>Staphylococcus sciuri</i>
		A39B (124)	21	<i>Staphylococcus hominis</i>
		A34M (128)	6	<i>Staphylococcus sciuri</i>
		A31M (188)	6	<i>Staphylococcus sciuri</i>
		A33M (187)	6	<i>Staphylococcus sciuri</i>
JO0	Site 9/ 0h	J03B (131)	6	<i>Staphylococcus sciuri</i>
		J04B (132)	6	<i>Staphylococcus sciuri</i>
		J01B (189)	4	<i>Staphylococcus sciuri</i>
		J02B (190)	6	<i>Staphylococcus sciuri</i>
JO3	Site 9/ 72h	J03M (138)	22	<i>Staphylococcus saprophyticus</i>
		J311B (148)	15	<i>Staphylococcus gallinarum</i>
		J33M (150)	6	<i>Staphylococcus sciuri</i>
		J34M (151)	6	<i>Staphylococcus sciuri</i>
		J31M (181)	6	<i>Staphylococcus sciuri</i>
		J32M (182)	6	<i>Staphylococcus sciuri</i>
MB	Site 10/ 72-96h	MB1M (156)	5	<i>Staphylococcus sciuri</i>
		MB11M (184)	5	<i>Staphylococcus sciuri</i>

^aSite 5: Ouagadougou (Zone Pilote production site), Site 6: Ouagadougou (Patte d'Oie market), Site 7: Bobo Dioulasso (market), Site 8: Brazzaville (Production site A), Site 9: Brazzaville (Production site B), Site 10: Brazzaville (market)

^b Rep-PCR : Repetitive element palindromic-polymerase chain reaction ; PFGE : Pulse Field Gel Electrophoresis

857 Table 3 Geographical distribution of the *Staphylococcus* species and associated clusters
858

Species	Rep-PCR/ PFGE Clusters	Burkina Faso/ Soumbala				Burkina Faso/ Bicalga			Congo/ Ntoba Mbodi		
		Site 1 ^b	Site 2	Site 3	Site 4	Site 5	Site 6	Site 7	Site 8	Site 9	Site 10
<i>S. arlettae</i>	17	- ^c	-	-	-	-	-	-	+	-	-
<i>S. aureus</i>	18	-	-	-	-	+	-	-	-	-	-
<i>S. cohnii</i>	16	-	-	-	-	-	+	-	-	-	-
	24	-	-	-	-	-	+	-	-	-	-
<i>S. epidermidis</i>	14	+	-	-	-	-	-	-	-	-	-
<i>S. gallinarum</i>	1	+	-	-	-	-	-	-	-	-	-
	15	-	-	-	+	-	-	-	+	+	-
	23	+	-	-	-	-	-	-	-	-	-
<i>S. haemolyticus</i>	20	-	-	-	-	-	-	+	-	-	-
<i>S. hominis</i>	21	-	-	-	-	-	-	-	+	-	-
<i>S. pasteurii</i>	11	-	-	+	-	-	-	-	-	-	-
<i>S. condimentii</i>	9	-	-	-	+	-	-	-	-	-	-
<i>S. piscifermentans</i>	10	-	+	-	-	-	-	-	-	-	-
<i>S. saprophyticus</i>	22	-	-	-	-	-	-	-	-	+	-
<i>S. sciuri</i>	2	-	+	-	-	-	-	-	-	-	-
	4	-	-	-	-	+	-	-	+	+	-
	5	-	+	-	+	+	-	-	-	-	+
	6	-	-	-	-	+	-	-	+	+	-
	7	-	+	-	-	-	-	-	-	-	-
	8	-	-	-	-	+	-	-	-	-	-
	25	-	-	-	-	-	-	-	+	-	-
<i>S. simulans</i>	3	+	+	+	+	-	-	-	-	-	-
<i>S. warneri</i>	12	-	-	-	-	-	-	-	+	-	-
	13	-	-	-	-	-	-	-	+	-	-
<i>M. caseolyticus</i>	19	-	-	-	-	-	+	-	-	-	-

859
860 ^a Rep-PCR : Repetitive element palindromic-polymerase chain reaction ; PFGE : Pulse Field Gel Electrophoresis

861 ^bSite 1: Ouagadougou (Samandin production site), Site 2: Fada market, Site 3: Ouagadougou (Patte d'Oie market), Site 4: Ouagadougou (Zogona market),
862 Site 5: Ouagadougou (Zone Pilote production site), Site 6: Ouagadougou (Patte d'Oie market), Site 7: Bobo Dioulasso (market), Site 8: Brazzaville
863 (Production site A), Site 9: Brazzaville (Production site B), Site 10: Brazzaville (market)

864 ^c - : not detected, + : detected

865 Table 4 Profiles of antimicrobial resistance (AMR) and toxin production potential of the species and clusters of *Staphylococcus* from
866 Bikalga, Soumbala, and Ntoba Mbodi

Origin ^a	Rep-PCR/ PFGE Cluster	Species	Number of Isolates	AMR phenotype ^b	AMR genes	Toxin genes/ Toxin A, B, C, D	Haemolysis on blood agar
Ntoba Mbodi (Co)	17	<i>S. arlettae</i>	2	CHL, CLI, FUS, SYN, SMX, TIA, TMP	dfr(A)	- ^c	-
Bikalga (BF)	18	<i>S. aureus</i>	2	CHL, ERY, PEN, SYN, SUL	blazA	-	+
Bikalga (BF)	16	<i>S. cohnii</i>	3	FOX, CHL, CLI, ERY, FUS, PEN, RIF, SMX, SYN, TIA, TMP	msr(A)	-	-
Bikalga (BF)	24	<i>S. cohnii</i>	1	FOX, CLI, ERY, FUS, PEN, RIF, SMX., SYN, TET, TIA, TMP	msr(A)	-	-
Soumbala (BF)	14	<i>S. epidermidis</i>	1	CHL, PEN, SMX, TET, TMP	blazA, dfr(A), dfr(G) tet(K)	-	±
Soumbala (BF)	1	<i>S. gallinarum</i>	1	FOX, CLI, SYN, TIA	tet(K)	-	-
Soumbala, Ntoba Mbodi (BF, Co)	15	<i>S. gallinarum</i>	3	CLI, FUS, PEN, SYN, TIA, SMX, TMP	tet(K)	-	-
Soumbala (BF)	23	<i>S. gallinarum</i>	3	FOX, CLI, FUS, PEN, SYN, TIA, TMP	tet(K)	-	-
Bikalga (BF)	20	<i>S. haemolyticus</i>	1	CLI, ERY, PEN, SYN, TIA, TMP	tet(K), msr(A)	-	±
Ntoba Mbodi (Co)	21	<i>S. hominis</i>	1	CHL, PEN, SMX, TET, TMP	blazA, tet(K), dfr(A)	-	-
Soumbala (BF)	11	<i>S. pasteurii</i>	1	SMX		-	±
Soumbala (BF)	10	<i>S. piscifermentans</i>	1	CHL, ERY, SMX, TIA		-	-
Soumbala (BF)	9	<i>S. condimentii</i>	1	SMX		-	-
Ntoba Mbodi (Co)	22	<i>S. saprophyticus</i>	1	CHL, PEN, TET, TMP	tet(K)	-	-
Soumbala (BF)	2	<i>S. sciuri</i>	1	CLI, FUS, SMX, TIA, TMP	mecA, mecA1	-	-
Bikalga, Ntoba Mbodi (BF, Co)	4	<i>S. sciuri</i>	6	CLI, FUS, TIA, TMP	mecA1	-	-
Bikalga, Soumbala, Ntoba Mbodi (BF, Co)	5	<i>S. sciuri</i>	13	CLI, FUS, TMP	mecA1	-	-
Bikalga, Ntoba Mbodi (BF, Co)	6	<i>S. sciuri</i>	16	CHL, PEN, SYN, SMX, TIA, TMP	mecA, mecA1	-	-
Soumbala (BF)	7	<i>S. sciuri</i>	1	CLI, ERY, FUS, SYN, TIA, TMP	mecA, mecA1	-	-
Bikalga (BF)	8	<i>S. sciuri</i>	2	CHL, PEN, SYN, TET, TIA, TMP	mecA1	-	-
Ntoba Mbodi (Co)	25	<i>S. sciuri</i>	2	CLI, FUS, SMX, SYN, TIA, TMP	mecA1	-	-
Soumbala (BF)	3	<i>S. simulans</i>	17	CHL, LZD, SYN, SMX, TMP		-	+
Ntoba Mbodi (Co)	12	<i>S. warneri</i>	4	FUS, PEN, SYN, SMX, TMP	blazA, dfr(A)	-	±
Ntoba Mbodi (Co)	13	<i>S. warneri</i>	1	CHL, ERY, PEN, SMX, TMP	blazA, Cat501 dfr(A), msr(A)	-	-
Bikalga (BF)	19	<i>M. caseolyticus</i>	5	CIP, MUP		-	-

867 ^a BF: Burkina Faso, Co: Congo


868 ^b The profile represents the antimicrobials to which the isolates were resistant to; Cefoxitin: FOX, Chloramphenicol: CHL, Ciprofloxacin: CIP,
869 Clindamycin: CLI, Erythromycin: ERY, Fusidate: FUS, Mupirocin: MUP, Quinupristin/Dalfopristin (Synercid): SYN, Linezolid: LZD, Penicillin:
870 PEN, Rifampicin: RIF, Sulfamethoxazole: SMX, Tiamulin: TIA, Trimethoprim: TMP ^c -: negative, ±: partial haemolysis, +: positive haemolysis

871 Table 5: Minimal Inhibitory Concentration (MIC) for donors, recipient and potential *E. faecalis* JH2-2 transconjugants (PET)

MIC (mg/L): Erythromycin		MIC (mg/L): Tetracycline	
Donor <i>S. warneri</i>	>32	Donor <i>S. saprophyticus</i>	64
Recipient <i>E. faecalis</i>	<1	Recipient <i>E. faecalis</i>	1
PET-E1	16	PET-T1	8
PET-E2	<1	PET-T2	8
PET-E3	>32	PET-T3	16
PET-E4	<1	PET-T4	8
PET-E5	<1	PET-T5	8
PET-E6	4	PET-T6	16
PET-E7	<1	PET-T7	16
PET-E8	<1	PET-T8	4
PET-E9	>32	PET-T9	16
PET-E10	<1	PET-T10	4
PET-E11	32		
PET-E12	<1		
PET-E13	>32		
PET-E14	<1		

872

873



874

875 Figure 1