

UWL REPOSITORY
repository.uwl.ac.uk

The role of spinal GABAB receptors in cancer-induced bone pain in rats

Zhou, Ya-Qun, Chen, Shu-Ping, Liu, Dai-Qiang, Manyande, Anne ORCID: <https://orcid.org/0000-0002-8257-0722>, Zhang, Wen, Yang, Shao-Bing, Xiong, Bing-Rui, Fu, Qiao-Chu, Song, Zhenpeng, Rittner, Heike, Ye, Da-Wei and Tian, Yu-Ke (2017) The role of spinal GABAB receptors in cancer-induced bone pain in rats. *Journal of Pain*, 18 (8). pp. 933-946. ISSN 1526-5900

<http://dx.doi.org/10.1016/j.jpain.2017.02.438>

This is the Accepted Version of the final output.

UWL repository link: <https://repository.uwl.ac.uk/id/eprint/3242/>

Alternative formats: If you require this document in an alternative format, please contact: open.research@uwl.ac.uk

Copyright:


Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy: If you believe that this document breaches copyright, please contact us at open.research@uwl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Accepted Manuscript

The Role of Spinal GABAB Receptors in Cancellor-Induced Bone Pain in Rats

Ya-Qun Zhou, Shu-Ping Chen, Dai-Qiang Liu, Anne Manyande, Wen Zhang, Shao-Bing Yang, Bing-Rui Xiong, Qiao-Chu Fu, Zhen-peng Song, Heike Rittner, Da-Wei Ye, Yu-Ke Tian



PII: S1526-5900(17)30514-X

DOI: [10.1016/j.jpain.2017.02.438](https://doi.org/10.1016/j.jpain.2017.02.438)

Reference: YJPAI 3394

To appear in: *Journal of Pain*

Received Date: 22 October 2016

Revised Date: 21 February 2017

Accepted Date: 26 February 2017

Please cite this article as: Zhou Y-Q, Chen S-P, Liu D-Q, Manyande A, Zhang W, Yang S-B, Xiong B-R, Fu Q-C, Song Z-p, Rittner H, Ye D-W, Tian Y-K, The Role of Spinal GABAB Receptors in Cancellor-Induced Bone Pain in Rats, *Journal of Pain* (2017), doi: 10.1016/j.jpain.2017.02.438.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The role of spinal GABAB receptors in cancer-induced bone pain in rats

Ya-Qun Zhou^{1,2}, Shu-Ping Chen^{1,2}, Dai-Qiang Liu^{1,2}, Anne Manyande³, Wen Zhang^{1,2}, Shao-Bing Yang^{1,2}, Bing-Rui Xiong^{1,2}, Qiao-Chu Fu^{1,2}, Zhen-peng Song^{1,2}, Heike Rittner⁴, Da-Wei Ye⁵, Yu-Ke Tian^{1,2}

1. Anesthesiology Institute, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China.

2. Department of Anesthesiology and Pain Medicine, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China.

3. School of Human and Social Sciences, University of West London, London, UK.

4. Department of Anesthesiology, University Hospital of Würzburg, Würzburg, Germany.

5. Cancer Center, Tongji Hospital, Tongji Medical college, Huazhong University of Science and Technology, Wuhan, China.

Corresponding authors:

Da-Wei Ye, MD, PhD.

Cancer Center, Tongji Hospital, Tongji Medical college, Huazhong University of Science and Technology, Wuhan, China.

TEL: 00862783663409, FAX: 00862783662853, E-mail: dy0711@gmail.com.

Yu-Ke Tian, MD, PhD.

Anesthesiology Institute, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China.

TEL: 00862783663173, FAX: 00862783662853, E-mail: yktian@tjh.tjmu.edu.cn.

Running title: The role of GABAB receptors in bone cancer pain

Disclosures: This work was supported by grants from National Natural Science Foundation of P.R. China (Grant No. 81400917, 81371250 and 81571053), and

Specialized Research Fund for the Doctoral Program of Higher Education

No.20130142120102, HUST No. 2014QT021. The authors have no conflict of interest.

Abstract

Cancer-induced bone pain (CIBP) remains a major challenge in advanced cancer patients due to our lack of understanding of its mechanisms. Previous studies have demonstrated the vital role of GABAB receptors (GABABRs) in regulating nociception and various neuropathic pain models have shown diminished activity of GABABRs. However, the role of spinal GABABRs in CIBP remains largely unknown. In this study, we investigated the specific cellular mechanisms of GABABRs in the development and maintenance of CIBP in rats. Our behavioral results show that both acute and chronic intrathecal treatment with baclofen, a GABABR agonist, significantly attenuated CIBP-induced mechanical allodynia and ambulatory pain. The expression levels of GABABRs were significantly decreased in a time-dependent manner and colocalized mostly with neuron and a minority with astrocyte and microglia. Chronic treatment with baclofen restored the expression of GABABRs and markedly inhibited the activation of cAMP-dependent protein kinase (PKA) and the cAMP-response element-binding protein (CREB) signaling pathway.

Perspective: Our findings provide the first evidence that downregulation of GABABRs contribute to the development and maintenance of CIBP and restore diminished GABABRs attenuates CIBP-induced pain behaviors at least partially by inhibiting the PKA/CREB signaling pathway. Therefore, spinal GABABR may become a potential therapeutic target for the management of CIBP.

Keywords: Cancer-induced bone pain; GABAB receptor; PKA; CREB; Baclofen

1. Introduction

More than one-third patients with metastatic bone disease experience moderate to severe pain, which significantly debilitates their quality of life^{39, 41, 56}. Cancer-induced bone pain (CIBP) is a complex pain state involving background, spontaneous and evoked pain^{21, 35, 54}. Currently, the management of CIBP is limited to the use of opioids^{28, 55}. However, high dose of opioids often lead to unwanted side effects including addiction, sedation, constipation, pruritus, nausea and vomiting^{6, 9}.

Although marked advances have been made in recent years, the mechanisms of CIBP are far from clear. Therefore, further elucidation of the mechanisms of CIBP is warranted to reveal novel therapeutic targets.

γ -aminobutyric acid (GABA), the main inhibitory neurotransmitter in the central nervous system (CNS), exerts its action via ionotropic GABAA receptors (GABAARs) and metabotropic GABAB receptors (GABABRs)⁴. GABABRs, composed of GABABR1 and GABABR2, are G-protein-coupled receptors (GPCRs) that elicit primarily inhibitory effects via the inhibition of presynaptic voltage-gated Ca^{2+} channels, activation of postsynaptic K^{+} channels, and inhibition of adenylyl cyclase^{5, 30}. In the spinal cord, GABABRs are mainly located on laminae I-III where nociceptive primary afferent fibers principally terminate¹³. Evidence is accumulating that GABABRs play a vital role in regulating nociception^{20, 34, 40}. Moreover, studies suggest that impairment of GABABRs contribute to neuropathic pain in various models including the spinal nerve ligation (SNL)²³, spinal cord injury¹⁶ and diabetic neuropathy^{3, 26, 44}. However, the role of spinal GABABRs in CIBP remains unknown. It has been shown that activation of GABABRs could inhibit adenylyl cyclase, and thus reduce the activity of cAMP-dependent protein kinase (PKA) signaling¹¹. The

PKA complex consists of two catalytic and two regulatory subunits, which bind cAMP and phosphorylated cAMP-response element-binding protein (p-CREB), respectively⁵³. Several lines of evidence have demonstrated that upregulation of PKA and p-CREB contribute to the development of chronic pain including CIBP^{17, 19, 29, 57}. If activation of GABABRs could alleviate CIBP, maybe downregulation of PKA and p-CREB is one of its intracellular mechanisms, which needs to be investigated. In the present study, using a CIBP model induced by intramedullary injection of Walker 256 cells into the tibia of rats, we tested the hypothesis that activation of GABABRs could alleviate CIBP via inhibiting PKA/CREB signaling pathway.

2. Material and Methods

2.1. Animals and ethical statement

Since female rats are more susceptible to Walker 256 mammary gland carcinoma cells, we chose virgin female Sprague-Dawley rats (180-200 g, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, PR China) in this study. Rats were kept under controlled conditions ($24 \pm 0.5^{\circ}\text{C}$, 12 h alternating light-dark cycle, with *ad libitum* access to water and food). All experimental protocols were approved by the Animal Care and Use Committee of Huazhong University of Science & Technology and were in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals and the guidelines of the International Association for the Study of Pain⁵⁸. All efforts were made to minimize the number of animals used and their suffering.

2.2. Intrathecal catheters and drug administration

As described previously, the intrathecal catheters were administered 5 days prior to the establishment of CIBP models^{22, 37, 49}. In brief, the rats were anesthetized by intraperitoneal injection (i.p.) of pentobarbital sodium (50 mg/kg). A PE-10

polyethylene catheter was then inserted into the subarachnoid space of the spinal cord between L5 and L6 spinous processes. The correct position of the catheter was verified by a tail flick response immediately after inserting the catheter and further confirmed by an intrathecal injection (i.t.) of 2% lidocaine (10 μ L). Animals exhibiting motor dysfunction were excluded from the experiments.

R(+)-Baclofen hydrochloride, a GABAB receptor agonist, was purchased from Sigma-Aldrich (USA) and dissolved in saline. 10 μ L baclofen (0.1 μ g, 0.5 μ g or 1 μ g) or vehicle (saline) was intrathecally injected via the implanted catheter, followed by 10 μ L saline for flushing. The dose of baclofen was based on our preliminary results and previous reports^{3, 26}. The drug administration protocol was as follows: for acute treatment, baclofen (i.t., 0.1 μ g, 0.5 μ g or 1 μ g /10 μ L) or vehicle (10 μ L) was administered on day 21 following CIBP model establishment; for chronic treatment, baclofen (i.t., 0.5 μ g/10 μ L, once a day) or vehicle (10 μ L) was administered from day 3 to day 14.

2.3. Preparation of carcinoma cells

Walker 256 rat mammary gland carcinoma cells (4×10^7 cells/mL, 1 mL) were inoculated into the abdominal cavity of a female Sprague-Dawley rat. After 7 days, cells were harvested from the ascitic fluid of the above rat. Cells were diluted to achieve a final concentration (4×10^7 cells/mL) for injection using a hemocytometer. The single-cell suspensions were maintained on ice prior to surgery.

2.4. Bone cancer pain model

The model of CIBP was established as described previously^{10, 15, 27}. Briefly, rats were anesthetized with pentobarbital sodium (50 mg/kg, i.p.). After shaving and disinfecting the left leg, a minimal incision was made to expose the proximal tibia. Then, the prepared Walker 256 cells (4×10^7 cells/mL, 10 μ L) were slowly injected

into the bone cavity using a 10 μ L Hamilton syringe. For the sham group, 10 μ L D-Hank's solution was injected instead. The injection site was closed using bone wax as soon as the syringe was removed. Finally, the wound was sutured with 3-0 silk thread.

2.5. Behavioral tests

Behavioral tests were performed on day 0, 3, 7, 14 and 21 after surgery between 8:00 and 12:00 to avoid diurnal variation. Mechanical allodynia was assessed by ipsilateral hind paw withdrawal threshold (PWT) to von Frey filament stimulation as described previously³⁸. Briefly, animals were placed in individual plastic boxes on a metal mesh floor and allowed to acclimatize for 30 min. Von Frey filaments, with ascending order of forces (1, 2, 4, 6, 8, 10, and 15 g), were applied for up to 6 s per filament to the mid-plantar of the right hind paw. Abrupt paw withdrawal, lickings, and shaking were considered as positive responses. Once a positive response was established, the paw was re-tested after a 5 min-rest, starting with the next descending von Frey filament until no response occurred. The lowest amount of force required to elicit a positive response was recorded as the PWT (in grams). Ambulatory pain was assessed by ipsilateral limb use during normal ambulation in a plastic observation box as described previously¹⁴. Limb use during spontaneous ambulation was scored on a scale of 0 to 4: 0 = complete lack of limb use, 1 = marked limping, 2 = extent between 1 and 3, 3 = slight limping, 4 = normal use. All the behavioral tests were performed by an investigator who was blinded to the experimental design.

2.6. Immunohistochemistry

Under deep anesthesia with pentobarbital sodium (60 mg/kg, i.p.), the rats were perfused intracardially with saline followed by 4% ice-cold paraformaldehyde in 0.1 M phosphate buffer saline (PBS). The L4-L5 spinal segments were removed and post-fixed in 4% paraformaldehyde for 4 h, and subsequently dehydrated in 30 % sucrose

solution overnight at 4°C. The samples were sectioned 30 µm thick in a cryostat (CM1900, Leica, Germany) and stored in PBS.

2.6.1. Single immunostainings

After washing three times in PBS, the sections were blocked with 5% donkey serum and 0.3% Triton X-100 for 1 h at 37°C, and then incubated overnight at 4°C with rabbit anti-GABABR1 antibody (1:50; DF4934; Affinity) or rabbit anti-GABABR2 antibody (1:50; A6594; ABclonal) or rabbit anti-PKAα antibody (1:50; ab26322; Abcam) or rabbit anti-phospho-CREB (pSer133) antibody (1:50; MA5-11192; Thermo). After washing three times in PBS, the sections were incubated with Alexa Fluor 488-labeled donkey anti-rabbit secondary antibody (1:500; A-21206; Invitrogen) for 2 h at 37°C and were washed in PBS. Nonspecific staining was determined by replacing primary or secondary antibodies with an antibody dilution buffer. Sections were rinsed, mounted and cover-slipped with 50% glycerol. Images were captured using a fluorescence microscope (DM2500, Leica, Germany). The GABABR1/GABABR2/PKA/p-CREB-immunolabeled surface areas were measured in laminae I-IV of the spinal cord dorsal horn using Image Pro Plus software. Quantification of the immunoreactivity was accomplished by calculating the percentages of immunostaining ($[\text{positive immunofluorescent surface area}]/[\text{total measured picture area}] \times 100$). Six rats of each group were used for statistical analysis. All the image analyses were performed by an investigator who was blinded to the experimental design.

2.6.2. Double-labeling procedures

For double immunofluorescence, the sections were incubated with a mixture of two primary antibodies followed by a mixture of Alexa 488-conjugated and Alexa 594-conjugated secondary antibodies (1:500; A-21207; Invitrogen). Specifically, to

identify the cell types that expressed GABABR1, GABABR2, PKA, and p-CREB, each of the antibodies for these molecules was mixed with mouse anti-neuronal nuclei antibody (NeuN; neuronal marker; 1:200; MAB377; Millipore), mouse anti-glial fibrillary acidic protein antibody (GFAP, astrocytic marker; 1:300; 3670; Cell Signaling Technology), or goat anti-Iba1 antibody (microglial marker; 1:300; ab5076; Abcam).

2.7. Western blotting

Under deep anesthesia with pentobarbital sodium (60 mg/kg, i.p.), the L4-L5 spinal segments were removed and homogenized in RIPA lysis buffer containing 50mM Tris (pH 7.4), 150mM NaCl, 1% Triton X-100, 1% sodium deoxycholate, 0.1% sodium dodecyl sulfate (SDS), 1mM NaF and 2mM EDTA. Then the homogenates were centrifuged at 12000 g for 30 minutes at 4°C. The supernatants were collected and the protein concentration was measured using the Bradford method. The samples were then heated at 95°C for 10 min in a loading buffer (pH 6.8): 250 mM Tris-HCl, 200 mM Sucrose, 300 mM DTT, 0.01% Coomassie brilliant blue-G, 8% sodium dodecyl sulfate (SDS). Equivalent amounts of proteins (50 µg) were separated using 10% SDS-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred onto polyvinylidene fluoride membranes (PVDF; IPVH00010; Millipore). After blocking with 5% bovine serum albumin in Tris-buffered saline and Tween 20 (TBST, 0.1%) for 2 h at 37°C, the membranes were incubated overnight at 4 °C with rabbit anti-GABABR1 antibody (1:1000; DF4934; Affinity), rabbit anti-GABABR2 antibody (1:500; A6594; ABclonal), rabbit anti-PKAc antibody (1:1000; ab26322; Abcam), rabbit anti-phospho-CREB (pSer133) antibody (1:500; MA5-11192; Thermo) and mouse anti-glyceraldehyde-3-phosphate dehydrogenase antibody (GAPDH; 1:5000; AS1039; Aspen), respectively. The membranes were then washed in TBST and

incubated with horseradish peroxidase-conjugated goat-anti-rabbit (1:5000; A21020; Abbkine) or goat-anti-mouse secondary antibody (1:5000; A21010; Abbkine) for 2 h at 37°C. The bands were finally visualized with SuperLumia ECL Plus HRP Substrate Kit (K22030; Abbkine) and then detected by a computerized image analysis system (BIO-RAD, ChemiDoc XRS+, USA). The intensity of protein blots was quantified using System with image Lab software (Bio-rad Laboratories), normalized to loading control GAPDH and expressed as the fold of control. The blot density of control groups was set as 1.

2.8. Statistical analysis

All data are presented as mean \pm SEM and analyzed using the GraphPad Prism version 5.01 for Windows (Graph Pad Software, San Diego, CA, USA). One-way analysis of variance (ANOVA) followed by Bonferroni *post hoc* test was used for Western blot and immunochemistry data. Two-way ANOVA with repeated measures, followed by Bonferroni *post hoc* test was used for PWT and limb use scores. $P < 0.05$ was considered statistically significant.

3. Results

3.1. Role of spinal GABABRs in CIBP-related behaviors

No significant differences in pain-related behaviors were observed among all groups at baseline. The ipsilateral PWT of CIBP rats were significantly decreased from day 7 after tumor cell implantation (TCI) to the last observation on day 21 (Figure 1A). Similarly, the ipsilateral limb use scores of CIBP rats were clearly decreased over the same time frame (Figure 1B). These results indicate that CIBP rats developed mechanical allodynia and ambulatory pain.

To investigate the role of spinal GABABRs in CIBP rats, we intrathecally injected baclofen, a GABABR agonist, into sham and CIBP rats. We first tested the analgesic

effect of a single dose of baclofen on day 21 following TCI (i.t., 0.1 μg , 0.5 μg or 1 μg /10 μL). The behavioral tests were conducted at 0, 0.25, 0.5, 1, 1.5, 2, 2.5, 3 h after baclofen injection. Compared with that of the vehicle group, intrathecal injection of 0.1 μg baclofen had no significant influence on PWT and limb use scores. However, baclofen (i.t., 0.5 μg or 1 μg /10 μL) treatment notably reversed the CIBP-related behaviors, beginning at 0.25 h, peaking at 0.5 h and lasting for at least 2 h (Figure 1C and D). Moreover, in terms of PWT and limb use scores, significant differences were found between these two groups (0.5 μg vs. 1 μg). These results indicate that the antinociceptive effects of baclofen are dose-dependent. For chronic treatment, baclofen (i.t., 0.5 μg /10 μL , once a day) or vehicle (10 μL) was administered for 12 days (from day 3 following TCI to day 14), and the behavioral tests were conducted before baclofen injection at day 0, 3, 7, 14 and 21. The results showed that both mechanical allodynia and ambulatory pain were alleviated considerably in CIBP rats treated with baclofen rather than vehicle (Figure 1E and F). To examine whether baclofen affect the pain threshold of naïve rats, we intrathecal injection of baclofen (0.1 μg , 0.5 μg or 1 μg /10 μL) into naïve rats. The behavioral tests were conducted at 0, 0.5, 1, 2 h after baclofen injection. The results showed that intrathecal injection of baclofen had no significant influence on PWT and limb use scores compared with vehicle group (Figure 1G and H).

Taken together, these results demonstrate that activation of GABABRs attenuates the development and maintenance of mechanical allodynia and ambulatory pain following TCI, which indicates an important role of GABABRs in CIBP.

3.2. Expression and cellular localization of spinal GABABRs in CIBP rats

To investigate the expression and cellular localization of GABABs in the spinal cord of CIBP rats, we performed western blot analysis and immunohistochemistry. As

shown in Figures 2A-B and 3A-B, the expression of GABABR1 and GABABR2 protein were significantly and time dependently decreased in CIBP rats, beginning at day 7 after TCI and remained at low levels until day 21, the last test day. In contrast, GABABR1 and GABABR2 proteins were at a higher level in the spinal cord of sham rats. Moreover, our immunohistochemistry data also showed that the expression of GABABR1 and GABABR2 were evidently reduced in the ipsilateral spinal cord of CIBP rats compared with sham rats (Figures 2C-E and 3C-E). We further conducted double immunofluorescence staining of GABABR1 or GABABR2 with NeuN (a neuronal marker), GFAP (an astrocytic marker), or Iba-1 (a microglial marker). As illustrated in Figures 2F-N and 3F-N, GABABR1 and GABABR2 were predominantly colocalized with NeuN, and a minority with GFAP and Iba-1 in the superficial spinal cord of CIBP rats. These results demonstrate that spinal GABABRs are downregulated after TCI, and expressed mostly in neurons and a minority in astrocyte and microglia.

3.3. Chronic treatment with baclofen restored the expression of GABABRs.

To investigate whether baclofen exerts its analgesic effect by restoring the diminished expression of GABABRs, we chronically treated CIBP rats with baclofen (i.t., 0.5 $\mu\text{g}/10 \mu\text{L}$) once a day from day 3 to day 14 following TCI. Three hours after the final administration, spinal samples were collected to detect the expressions of GABABR1 and GABABR2 using western blot and immunohistochemistry. As shown in Figure 4A-B, the protein expression of GABABRs were significantly upregulated after baclofen treatment. Consistent with the western blot data, our immunohistochemistry also showed increased immunoreactivity of GABABRs in the superficial dorsal horn of spinal cord (Figure 4C-L). These results suggested that chronic treatment with baclofen could restore the expression of GABABRs.

3.4. Involvement of spinal PKA/CREB signaling pathway in the analgesic effect of baclofen

The PKA/CREB signaling pathway has been shown to participate in the development of chronic pain including CIBP. We consistently found that the expression of spinal PKA and p-CREB were significantly increased in a time-dependent manner, beginning at day 3 after TCI and remained at high levels until day 21, whereas PKA and p-CREB were at lower levels in the spinal cord of sham rats (Figures 5A-B and 6A-B). Moreover, our immunohistochemistry data also showed that the expression of PKA and p-CREB was significantly increased in the ipsilateral spinal cord of CIBP rats compared with sham rats (Figures 5C-E and 6C-E). However, the cellular localization of PKA and p-CREB in the spinal cord under CIBP condition remains unknown. Our double immunofluorescence staining showed that PKA was colocalized with NeuN, GFAP and Iba-1 and (Figures 5F-N), whereas p-CREB was strongly colocalized with NeuN (Figures 6F-N). These results confirm the activation of PKA/CREB signaling pathway in the CIBP condition.

To further examine the involvement of spinal PKA/CREB signaling pathway in the analgesic effect of baclofen, we intrathecally injected baclofen (0.5 μ g/10 μ L) or vehicle (10 μ L) into CIBP rats or sham rats daily for 12 consecutive days (from day 3 following TCI to day 14). Three hours after the final administration, spinal samples were collected to detect the expressions of PKA and p-CREB using western blot and immunohistochemistry. Our results showed that the upregulated protein levels of PKA and p-CREB were considerably attenuated by repeated intrathecal injection of baclofen (Figures 7A-B). Additionally, our immunofluorescence results showed that CIBP-induced activation of PKA and p-CREB was markedly suppressed by baclofen

treatment (Figures 7C-L). These results indicate that restored expression of GABABRs could suppress the activation of PKA/CREB signaling pathway. Collectively, these data suggest that activation of GABABRs alleviates CIBP at least partly by inhibiting spinal PKA/CREB signaling pathway.

4. Discussion

The current study demonstrated that (1) activation of GABABRs via intrathecal administration of baclofen, a GABABR agonist, significantly attenuated CIBP-induced mechanical allodynia and ambulatory pain in a dose-dependent manner, (2) inoculation of Walker 256 mammary gland carcinoma cells into a rat tibia induced downregulation of GABABRs in the spinal cord in a time-dependent manner, (3) spinal GABABRs were mostly expressed in neurons and a minority in astrocytes and microglia, (4) chronic treatment with baclofen restored the expression of GABABRs and significantly inhibited the activation of PKA/CREB signaling pathway in CIBP rats. Taken together, these results provide the first evidence that activation of spinal GABABRs mitigates CIBP at least partially by inhibiting the PKA/CREB signaling pathway.

Despite advanced progress has been made in the last decades for the management of chronic pain^{1, 24, 43, 47}, its mechanisms remains far from clear. It has long been suggested that diminished inhibitory neurotransmission plays a pivotal role in chronic pain states^{46, 50}. As the main inhibitory neurotransmitter in the CNS, GABA exerts its effect via ionotropic GABAARs and metabotropic GABABRs⁷. Spinal GABABRs are abundantly expressed in primary afferent fibers and interneurons on laminae I-III⁴⁸. Genetic deletion of GABABR1 or GABABR2 produces prominent hyperalgesia^{12, 36}. Previously, impairment of GABABRs has been demonstrated to participate in the development of neuropathic pain induced by SNL²³, spinal cord injury¹⁶ and diabetic

neuropathy^{3, 26, 44}. Moreover, activation of GABABRs by intrathecal injection of baclofen produces analgesic effects^{2, 42, 51}. In our study, we provided consistent results showing that GABABRs were significantly decreased in the ipsilateral spinal cord dorsal horn after TCI (Figures 2A-E and 3A-E). Furthermore, the decreased expression of GABABRs matches the time frame that coincides with the development of mechanical allodynia and ambulatory pain. Notably, both acute and chronic treatment with baclofen markedly alleviated the CIBP-induced mechanical allodynia and ambulatory pain (Figures 1C-F). These findings indicate that downregulation of GABABRs may be important contributors to diminished GABAergic inhibition under CIBP condition. More importantly, repeated intrathecal administration of baclofen upregulated the protein expression (Figure 3A-B) and immunoreactivity of GABABRs (Figure 3C-L), indicating restored expression of GABABRs after baclofen treatment. However, it is worth mentioning that several studies reported that loss of GABAergic neurons is not necessary for the development of neuropathic pain³¹⁻³³. These inconsistent results might be due to the difference in animal model, time of behavioral tests and detection methods.

PKA has been demonstrated to play a vital role in the processing of nociception at both peripheral and central levels^{8, 53}. Moreover, electrophysiological results also illustrate that PKA is involved in modulating neuronal excitability in response to noxious stimuli^{19, 52}. When PKA is activated, its catalytic subunit can phosphorylate CREB in the spinal cord, which contributes to the prolonged synaptic plasticity strengthened during central sensitization^{25, 45}. Previous studies have established that activation of PKA/CREB signaling pathway is involved in various types of chronic pain including CIBP^{17, 29}, and intrathecal treatment with H-89, a PKA inhibitor, significantly attenuates CIBP-related behaviors¹⁸. It is also shown that GABABR-

mediated inhibition of adenylyl cyclase reduces PKA activity, thereby influencing the regulation of gene expression⁷. Therefore, we determined whether inhibiting PKA/CREB signaling pathway is one of the mechanisms underlying the analgesic effect of the GABABR agonist. Consistent with previous researches, we found upregulated expression of PKA and p-CREB in the ipsilateral spinal cord of CIBP rats, whereas their expression in sham rats was at lower levels (Figures 5A-E and 6A-E). Furthermore, we performed double immunofluorescence to examine their cellular localization. We provided the first evidence that PKA was colocalized with neurons, astrocytes and microglia, whereas p-CREB was strongly colocalized with neurons under the CIBP condition (Figures 4F-N and 5F-N). In addition, CIBP rats that received chronic treatment with baclofen showed decreased expression of PKA and p-CREB in the ipsilateral spinal cord dorsal horn (Figure 7A-L). These results indicate that inhibition of PKA/CREB signaling pathway may contribute to the analgesic effect of the GABABR agonist.

In summary, the present study suggests that downregulation of GABABRs in the spinal cord contributes to the initiation and development of CIBP, and restores diminished GABABRs attenuates CIBP-induced mechanical allodynia and ambulatory pain at least partially by inhibiting the PKA/CREB signaling pathway. Therefore, spinal GABABR may become a potential therapeutic target for the management of CIBP.

Conflict of interest

The authors declare no conflict of interests

Acknowledgements

This work was supported by grants from National Natural Science Foundation of P.R. China (Grant No. 81400917, 81371250 and 81571053) and Specialized Research

Fund for the Doctoral Program of Higher Education No.20130142120102, HUST No. 2014QT021.

References

1. An K, Xu Y, Yang H, Shu HH, Xiang HB, Tian YK. Subarachnoid transplantation of immortalized galanin-overexpressing astrocytes attenuates chronic neuropathic pain. *European journal of pain*. 14:595-601, 2010
2. Ataka T, Kumamoto E, Shimoji K, Yoshimura M. Baclofen inhibits more effectively C-afferent than Adelta-afferent glutamatergic transmission in substantia gelatinosa neurons of adult rat spinal cord slices. *Pain*. 86:273-282, 2000
3. Bai HP, Liu P, Wu YM, Guo WY, Guo YX, Wang XL. Activation of spinal GABAB receptors normalizes N-methyl-D-aspartate receptor in diabetic neuropathy. *Journal of the neurological sciences*. 341:68-72, 2014
4. Benarroch EE. GABAB receptors: structure, functions, and clinical implications. *Neurology*. 78:578-584, 2012
5. Benke D, Balakrishnan K, Zemoura K. Regulation of cell surface GABA(B) receptors: contribution to synaptic plasticity in neurological diseases. *Advances in pharmacology*. 73:41-70, 2015
6. Boland JW, Ziegler L, Boland EG, McDermid K, Bennett MI. Is regular systemic opioid analgesia associated with shorter survival in adult patients with cancer? A systematic literature review. *Pain*. 156:2152-2163, 2015
7. Chalifoux JR, Carter AG. GABAB receptor modulation of synaptic function. *Current opinion in neurobiology*. 21:339-344, 2011
8. Chen HS, Lei J, He X, Qu F, Wang Y, Wen WW, You HJ, Arendt-Nielsen L. Peripheral involvement of PKA and PKC in subcutaneous bee venom-induced

- persistent nociception, mechanical hyperalgesia, and inflammation in rats. *Pain*. 135:31-36, 2008
9. Christo PJ, Mazloomdoost D. Interventional pain treatments for cancer pain. *Annals of the New York Academy of Sciences*. 1138:299-328, 2008
10. Fu Q, Shi D, Zhou Y, Zheng H, Xiang H, Tian X, Gao F, Manyande A, Cao F, Tian Y, Ye D. MHC-I promotes apoptosis of GABAergic interneurons in the spinal dorsal horn and contributes to cancer induced bone pain. *Experimental neurology*. 286:12-20, 2016
11. Gassmann M, Bettler B. Regulation of neuronal GABA(B) receptor functions by subunit composition. *Nature reviews. Neuroscience*. 13:380-394, 2012
12. Gassmann M, Shaban H, Vigot R, Sansig G, Haller C, Barbieri S, Humeau Y, Schuler V, Müller M, Kinzel B, Klebs K, Schmutz M, Froestl W, Heid J, Kelly PH, Gentry C, Jatton AL, Van der Putten H, Mombereau C, Lecourtier L, Mosbacher J, Cryan JF, Fritschy JM, Lüthi A, Kaupmann K, Bettler B. Redistribution of GABAB(1) protein and atypical GABAB responses in GABAB(2)-deficient mice. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 24:6086-6097, 2004
13. Goudet C, Magnaghi V, Landry M, Nagy F, Gereau RWt, Pin JP. Metabotropic receptors for glutamate and GABA in pain. *Brain research reviews*. 60:43-56, 2009
14. Guan X, Fu Q, Xiong B, Song Z, Shu B, Bu H, Xu B, Manyande A, Cao F, Tian Y. Activation of PI3Kgamma/Akt pathway mediates bone cancer pain in rats. *Journal of neurochemistry*. 134:590-600, 2015
15. Guan XH, Fu QC, Shi D, Bu HL, Song ZP, Xiong BR, Shu B, Xiang HB, Xu B, Manyande A, Cao F, Tian YK. Activation of spinal chemokine receptor CXCR3 mediates bone cancer pain through an Akt-ERK crosstalk pathway in rats. *Experimental neurology*. 263:39-49, 2015

16. Gwak YS, Tan HY, Nam TS, Paik KS, Hulsebosch CE, Leem JW. Activation of spinal GABA receptors attenuates chronic central neuropathic pain after spinal cord injury. *Journal of neurotrauma*. 23:1111-1124, 2006
17. Hang LH, Yang JP, Shao DH, Chen Z, Wang H. Involvement of spinal PKA/CREB signaling pathway in the development of bone cancer pain. *Pharmacological reports : PR*. 65:710-716, 2013
18. Hang LH, Yang JP, Shao DH, Chen Z, Wang H. Activation of spinal TDAG8 and its downstream PKA signaling pathway contribute to bone cancer pain in rats. *The European journal of neuroscience*. 36:2107-2117, 2012
19. Huang ZJ, Li HC, Cowan AA, Liu S, Zhang YK, Song XJ. Chronic compression or acute dissociation of dorsal root ganglion induces cAMP-dependent neuronal hyperexcitability through activation of PAR2. *Pain*. 153:1426-1437, 2012
20. Ishii H, Izumi H. GABAB receptors in the NTS mediate the inhibitory effect of trigeminal nociceptive inputs on parasympathetic reflex vasodilation in the rat masseter muscle. *American journal of physiology. Regulatory, integrative and comparative physiology*. 302:R776-784, 2012
21. Ke C, Gao F, Tian X, Li C, Shi D, He W, Tian Y. Slit2/Robo1 Mediation of Synaptic Plasticity Contributes to Bone Cancer Pain. *Molecular neurobiology*. 54:295-307, 2017
22. Ke CB, He WS, Li CJ, Shi D, Gao F, Tian YK. Enhanced SCN7A/Nax expression contributes to bone cancer pain by increasing excitability of neurons in dorsal root ganglion. *Neuroscience*. 227:80-89, 2012
23. Laffray S, Bouali-Benazzouz R, Papon MA, Favereaux A, Jiang Y, Holm T, Spriet C, Desbarats P, Fossat P, Le Feuvre Y, Decossas M, Héliot L, Langel U, Nagy

- F, Landry M. Impairment of GABAB receptor dimer by endogenous 14-3-3zeta in chronic pain conditions. *The EMBO journal*. 31:3239-3251, 2012
24. Li M, Mei W, Wang P, Yu Y, Qian W, Zhang ZG, Tian YK. Propofol reduces early post-operative pain after gynecological laparoscopy. *Acta anaesthesiologica Scandinavica*. 56:368-375, 2012
25. Liou JT, Liu FC, Hsin ST, Yang CY, Lui PW. Inhibition of the cyclic adenosine monophosphate pathway attenuates neuropathic pain and reduces phosphorylation of cyclic adenosine monophosphate response element-binding in the spinal cord after partial sciatic nerve ligation in rats. *Anesthesia and analgesia*. 105:1830-1837, 2007
26. Liu P, Guo WY, Zhao XN, Bai HP, Wang Q, Wang XL, Zhang YZ. Intrathecal baclofen, a GABAB receptor agonist, inhibits the expression of p-CREB and NR2B in the spinal dorsal horn in rats with diabetic neuropathic pain. *Canadian journal of physiology and pharmacology*. 92:655-660, 2014
27. Liu X, Bu H, Liu C, Gao F, Yang H, Tian X, Xu A, Chen Z, Cao F, Tian Y. Inhibition of glial activation in rostral ventromedial medulla attenuates mechanical allodynia in a rat model of cancer-induced bone pain. *Journal of Huazhong University of Science and Technology. Medical sciences*. 32:291-298, 2012
28. Mantyh P. Bone cancer pain: causes, consequences, and therapeutic opportunities. *Pain*. 154 Suppl 1:S54-62, 2013
29. Miletic G, Pankratz MT, Miletic V. Increases in the phosphorylation of cyclic AMP response element binding protein (CREB) and decreases in the content of calcineurin accompany thermal hyperalgesia following chronic constriction injury in rats. *Pain*. 99:493-500, 2002

30. Pinard A, Seddik R, Bettler B. GABAB receptors: physiological functions and mechanisms of diversity. *Advances in pharmacology*. 58:231-255, 2010
31. Polgar E, Gray S, Riddell JS, Todd AJ. Lack of evidence for significant neuronal loss in laminae I-III of the spinal dorsal horn of the rat in the chronic constriction injury model. *Pain*. 111:144-150, 2004
32. Polgar E, Hughes DI, Arham AZ, Todd AJ. Loss of neurons from laminae I-III of the spinal dorsal horn is not required for development of tactile allodynia in the spared nerve injury model of neuropathic pain. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 25:6658-6666, 2005
33. Polgar E, Hughes DI, Riddell JS, Maxwell DJ, Puskar Z, Todd AJ. Selective loss of spinal GABAergic or glycinergic neurons is not necessary for development of thermal hyperalgesia in the chronic constriction injury model of neuropathic pain. *Pain*. 104:229-239, 2003
34. Reis GM, Duarte ID. Baclofen, an agonist at peripheral GABAB receptors, induces antinociception via activation of TEA-sensitive potassium channels. *British journal of pharmacology*. 149:733-739, 2006
35. Schneider G, Voltz R, Gaertner J. Cancer Pain Management and Bone Metastases: An Update for the Clinician. *Breast care*. 7:113-120, 2012
36. Schuler V, Luscher C, Blanchet C, Klix N, Sansig G, Klebs K, Schmutz M, Heid J, Gentry C, Urban L, Fox A, Spooren W, Jatton AL, Vigouret J, Pozza M, Kelly PH, Mosbacher J, Froestl W, Käslin E, Korn R, Bischoff S, Kaupmann K, van der Putten H, Bettler B. Epilepsy, hyperalgesia, impaired memory, and loss of pre- and postsynaptic GABA(B) responses in mice lacking GABA(B1). *Neuron*. 31:47-58, 2001

37. Song Z, Xiong B, Zheng H, Manyande A, Guan X, Cao F, Ren L, Zhou Y, Ye D, Tian Y. STAT1 as a downstream mediator of ERK signaling contributes to bone cancer pain by regulating MHC II expression in spinal microglia. *Brain, behavior, and immunity*. 60:161-173, 2017
38. Song ZP, Xiong BR, Guan XH, Cao F, Manyande A, Zhou YQ, Zheng H, Tian YK. Minocycline attenuates bone cancer pain in rats by inhibiting NF-kappaB in spinal astrocytes. *Acta pharmacologica Sinica*. 37:753-762, 2016
39. Sun Y, Ye DW, Zhang P, Wu YX, Wang BY, Peng G, Yu SY. Anti-rheumatic drug iguratimod (T-614) alleviates cancer-induced bone destruction via down-regulating interleukin-6 production in a nuclear factor-kappaB-dependent manner. *Journal of Huazhong University of Science and Technology. Medical sciences*. 36:691-699, 2016
40. Takeda M, Ikeda M, Takahashi M, Kanazawa T, Nasu M, Matsumoto S. Suppression of ATP-induced excitability in rat small-diameter trigeminal ganglion neurons by activation of GABAB receptor. *Brain research bulletin*. 98:155-162, 2013
41. van den Beuken-van Everdingen MH, de Rijke JM, Kessels AG, Schouten HC, van Kleef M, Patijn J. Prevalence of pain in patients with cancer: a systematic review of the past 40 years. *Annals of oncology*. 18:1437-1449, 2007
42. van der Plas AA, van Rijn MA, Marinus J, Putter H, van Hilten JJ. Efficacy of intrathecal baclofen on different pain qualities in complex regional pain syndrome. *Anesthesia and analgesia*. 116:211-215, 2013
43. Wang GM, Tian XB, Chen JP, Yang SB, Gao F, Yang H, An K, Tian YK. Prevention of neuropathic pain in an animal model of spare nerve injury following oral immunization with recombinant adenovirus serotype 5-mediated NR2B gene transfer. *Gene therapy*. 14:1681-1687, 2007

44. Wang XL, Zhang Q, Zhang YZ, Liu YT, Dong R, Wang QJ, Guo YX. Downregulation of GABAB receptors in the spinal cord dorsal horn in diabetic neuropathy. *Neuroscience letters*. 490:112-115, 2011
45. Wu J, Su G, Ma L, Zhang X, Lei Y, Li J, Lin Q, Fang L. Protein kinases mediate increment of the phosphorylation of cyclic AMP-responsive element binding protein in spinal cord of rats following capsaicin injection. *Molecular pain*. 1:26, 2005
46. Xu B, Guan XH, Yu JX, Lv J, Zhang HX, Fu QC, Xiang HB, Bu HL, Shi D, Shu B, Qin LS, Manyande A, Tian YK. Activation of spinal phosphatidylinositol 3-kinase/protein kinase B mediates pain behavior induced by plantar incision in mice. *Experimental neurology*. 255:71-82, 2014
47. Xu Y, Tian XB, An K, Yang H, Tian YK. Lumbar transplantation of immortalized enkephalin-expressing astrocytes attenuates chronic neuropathic pain. *European journal of pain*. 12:525-533, 2008
48. Yang K, Ma WL, Feng YP, Dong YX, Li YQ. Origins of GABA(B) receptor-like immunoreactive terminals in the rat spinal dorsal horn. *Brain research bulletin*. 58:499-507, 2002
49. Ye D, Bu H, Guo G, Shu B, Wang W, Guan X, Yang H, Tian X, Xiang H, Gao F. Activation of CXCL10/CXCR3 signaling attenuates morphine analgesia: involvement of Gi protein. *Journal of molecular neuroscience*. 53:571-579, 2014
50. Zeilhofer HU, Benke D, Yevenes GE. Chronic pain states: pharmacological strategies to restore diminished inhibitory spinal pain control. *Annual review of pharmacology and toxicology*. 52:111-133, 2012
51. Zemoura K, Ralvenius WT, Malherbe P, Benke D. The positive allosteric GABAB receptor modulator rac-BHFF enhances baclofen-mediated analgesia in neuropathic mice. *Neuropharmacology*. 108:172-178, 2016

52. Zhang JM, Li H, Liu B, Brull SJ. Acute topical application of tumor necrosis factor alpha evokes protein kinase A-dependent responses in rat sensory neurons. *Journal of neurophysiology*. 88:1387-1392, 2002
53. Zhou XL, Wang Y, Zhang CJ, Yu LN, Cao JL, Yan M. PKA is required for the modulation of spinal nociceptive information related to ephrinB-EphB signaling in mice. *Neuroscience*. 284:546-554, 2015
54. Zhou YQ, Gao HY, Guan XH, Yuan X, Fang GG, Chen Y, Ye DW. Chemokines and Their Receptors: Potential Therapeutic Targets for Bone Cancer Pain. *Current pharmaceutical design*. 21:5029-5033, 2015
55. Zhou YQ, Liu Z, Liu HQ, Liu DQ, Chen SP, Ye DW, Tian YK. Targeting glia for bone cancer pain. *Expert opinion on therapeutic targets*. 20:1365-1374, 2016
56. Zhou YQ, Liu Z, Liu ZH, Chen SP, Li M, Shahveranov A, Ye DW, Tian YK. Interleukin-6: an emerging regulator of pathological pain. *Journal of neuroinflammation*. 13:141, 2016
57. Zhu GQ, Liu S, He DD, Liu YP, Song XJ. Activation of the cAMP-PKA signaling pathway in rat dorsal root ganglion and spinal cord contributes toward induction and maintenance of bone cancer pain. *Behavioural pharmacology*. 25:267-276, 2014
58. Zimmermann M. Ethical guidelines for investigations of experimental pain in conscious animals. *Pain*. 16:109-110, 1983

Figure legends

Fig. 1. Activation of GABAB receptors (GABABRs) alleviated cancer-induced bone pain (CIBP)-related behaviors. (A, B) The ipsilateral paw withdrawal thresholds (PWT) and limb use scores were significantly decreased from day 7 after tumor cell implantation (TCI) to the last observation on day 21 ($***P < 0.001$ compared with the naive group, $n = 6$ in each group). (C, D) A single dose of baclofen (i.t., 0.5 μg and 1 μg , but not 0.1 μg , on day 21) significantly reversed the CIBP-induced mechanical allodynia and ambulatory pain ($*P < 0.05$, $**P < 0.01$, $***P < 0.001$ compared with the vehicle group, $\#P < 0.05$, $\##P < 0.01$ compared with the group treated with 0.5 μg baclofen, $n = 6$ in each group). The behavioral tests were conducted at 0, 0.25, 0.5, 1, 1.5, 2, 2.5, 3 h after baclofen injection. (E, F) For chronic treatment, baclofen (i.t., 0.5 $\mu\text{g}/10 \mu\text{L}$, once a day) or vehicle (10 μL) was administered for 12 days (from day 3 following TCI to day 14). The behavioral tests were conducted before baclofen injection at day 0, 3, 7, 14 and 21 after surgery. The development of ipsilateral mechanical allodynia and ambulatory pain of CIBP rats were significantly delayed and attenuated by baclofen ($**P < 0.01$, $***P < 0.001$ compared with the sham+vehicle group, $\#P < 0.05$, $\##P < 0.01$, $\###P < 0.001$ compared with the CIBP+vehicle group, $n = 6$ in each group). (G, H) Intrathecal injection of baclofen (0.1 μg , 0.5 μg or 1 $\mu\text{g}/10 \mu\text{L}$) into naïve rats had no significant influence on PWT and limb use scores compared with vehicle group ($n = 3$ in each group). The behavioral tests were conducted at 0, 0.5, 1, 2 h after baclofen injection.

Fig. 2. Expression and cellular localization of GABAB receptor 1 (GABABR1) in spinal cord dorsal horn of cancer-induced bone pain (CIBP) rats. (A, B) Western blot analysis showed the time course of GABABR1 expression in sham and CIBP rats ($***P < 0.001$ compared with the sham group, $n = 6$ in each group). The fold change for the density of GABABR1 was normalized to GAPDH for each sample

respectively. The fold change of GABABR1 in the sham group was set at 1 for quantification. (C-E) Immunohistochemistry data showed that GABABR1 was downregulated in the ipsilateral spinal cord dorsal horn at day 21 after tumor cell implantation (TCI) (n = 4 in each group). (F-N) Representative photomicrographs of GABABR1 (green) double fluorescence labeling with NeuN (red) for neurons, GFAP (red) for astrocytes and Iba1 (red) for microglia in the ipsilateral spinal cord at day 21 after TCI. Photomicrographs were taken from ipsilateral spinal cord dorsal horns (laminae I-II; as indicated in C) of CIBP rats (n = 4 in each group). The results showed that GABABR1 was co-expressed mostly with neurons (yellow) and a minority with astrocytes (yellow) and microglia (yellow).

Fig. 3. Expression and cellular localization of GABAB receptor 2 (GABABR2) in spinal cord dorsal horn of cancer-induced bone pain (CIBP) rats. (A, B) Western blot analysis showed the time course of GABABR2 expression in sham and CIBP rats (** $P < 0.05$, *** $P < 0.001$ compared with the sham group, n = 6 in each group). The fold change for the density of GABABR2 was normalized to GAPDH for each sample respectively. The fold change of GABABR2 in the sham group was set at 1 for quantification. (C-E) Immunohistochemistry data showed that GABABR2 was downregulated in the ipsilateral spinal cord dorsal horn at day 21 after tumor cell implantation (TCI) (n = 4 in each group). (F-N) Representative photomicrographs of GABABR2 (green) double fluorescence labeling with NeuN (red) for neurons, GFAP (red) for astrocytes and Iba1 (red) for microglia in the ipsilateral spinal cord at day 21 after TCI. Photomicrographs were taken from ipsilateral spinal cord dorsal horns (laminae I-II; as indicated in C) of CIBP rats (n = 4 in each group). The results showed that GABABR2 was co-expressed mostly with neurons (yellow) and a minority with astrocytes (yellow) and microglia (yellow).

Fig. 4. Repeated intrathecal administration of baclofen reversed tumor cell implantation (TCI)-induced downregulation of GABABRs in the spinal cord. (A, B) Western blot and data summary showed that baclofen restored the protein expression of GABABR1 and GABABR2 (** $P < 0.01$, *** $P < 0.001$ compared with the sham+vehicle group. # $P < 0.05$ compared with the CIBP+vehicle group. $n = 6$ in each group). Baclofen (i.t., 0.5 $\mu\text{g}/10 \mu\text{L}$) or vehicle (10 μL) was administered once a day from day 3 following tumor cell implantation (TCI) to day 14. Tissues were collected three hours after the last spinal injection. (C-J) Representative photomicrographs showed that baclofen reversed TCI-induced downregulation of GABABRs in the spinal cord. Baclofen (i.t., 0.5 $\mu\text{g}/10 \mu\text{L}$) or vehicle (10 μL) was administered once a day from day 3 following tumor cell implantation (TCI) to day 14. Tissues were collected three hours after the last spinal injection. (K, L) GABABRs-immunolabeled surface area were quantified from the spinal dorsal horn (laminae I-IV; as indicated in C) using Image Pro Plus software. Quantification of GABABRs immunoreactivity was accomplished by calculating the percentages of immunostaining ($[\text{positive immunofluorescent surface area}]/[\text{total measured picture area}] \times 100$). Six rats of each group were used for statistical analysis (** $P < 0.01$, *** $P < 0.001$ compared with the sham+vehicle group. # $P < 0.05$ compared with the CIBP+vehicle group. $n = 6$ in each group).

Fig. 5. Expression and cellular localization of cAMP-dependent protein kinase (PKA) in spinal cord dorsal horn of cancer-induced bone pain (CIBP) rats. (A, B) Western blot analysis showed the time course of PKA expression in CIBP rats (** $P < 0.001$ compared with the sham group, $n = 6$ in each group). The fold change for the density of PKA was normalized to GAPDH for each sample respectively. The fold change of PKA in the sham group was set at 1 for quantification. (C-E)

Immunohistochemistry data showed that PKA was upregulated in the ipsilateral spinal cord dorsal horn at day 21 after tumor cell implantation (TCI) (n = 4 in each group).


(F-N) Representative photomicrographs of PKA (green) double fluorescence labeling with NeuN (red) for neurons, GFAP (red) for astrocytes and Iba1 (red) for microglia in the ipsilateral spinal cord at day 21 after TCI. Photomicrographs were taken from ipsilateral spinal cord dorsal horns (laminae I-II; as indicated in E) of CIBP rats (n = 4 in each group). The results showed that PKA was colocalized mostly with astrocytes (yellow) and microglia (yellow) and a minority with neurons (yellow).


Fig. 6. Expression and cellular localization of phosphorylated cAMP-response element-binding protein (p-CREB) in spinal cord dorsal horn of cancer-induced bone pain (CIBP) rats.


(A, B) Western blot analysis showed the time course of p-CREB expression in CIBP rats (***) $P < 0.001$ compared with the sham group, n = 6 in each group). The fold change for the density of p-CREB was normalized to GAPDH for each sample respectively. The fold change of p-CREB in the sham group was set at 1 for quantification. (C-E) Immunohistochemistry data showed that p-CREB as upregulated in the ipsilateral spinal cord dorsal horn at day 21 after tumor cell implantation (TCI) (n = 4 in each group). (F-N) Representative photomicrographs of p-CREB (green) double fluorescence labeling with NeuN (red) for neurons, GFAP (red) for astrocytes and Iba1 (red) for microglia in the ipsilateral spinal cord at day 21 after TCI. Photomicrographs were taken from ipsilateral spinal cord dorsal horns (laminae I-II; as indicated in E) of CIBP rats (n = 4 in each group). The results showed that p-CREB was colocalized mostly with neurons (yellow) and a minority with astrocytes (yellow) and microglia (yellow).


Fig. 7. Repeated intrathecal administration of baclofen reversed tumor cell implantation (TCI)-induced upregulation of PKA and p-CREB in the spinal cord.















(A, B) Western blot and data summary showed inhibitory effect of baclofen on CIBP-induced increased expression of PKA and p-CREB protein (*P < 0.05, ***P < 0.001 compared with the sham+vehicle group. ####P < 0.001 compared with the CIBP+vehicle group. n = 6 in each group). Baclofen (i.t., 0.5 µg/10 µL) or vehicle (10 µL) was administered once a day from day 3 following tumor cell implantation (TCI) to day 14. Tissues were collected three hours after the last spinal injection. (C-J) Representative photomicrographs showed that baclofen inhibited TCI-induced upregulation of PKA and p-CREB in the spinal cord. Baclofen (i.t., 0.5 µg/10 µL) or vehicle (10 µL) was administered once a day from day 3 following tumor cell implantation (TCI) to day 14. Tissues were collected three hours after the last spinal injection. (K, L) PKA/p-CREB-immunolabeled surface area were quantified from the spinal dorsal horn (laminae I-IV; as indicated in C) using Image Pro Plus software. Quantification of PKA/p-CREB immunoreactivity was accomplished by calculating the percentages of immunostaining ([positive immunofluorescent surface area]/[total measured picture area] × 100). Six rats of each group were used for statistical analysis (***P < 0.001 compared with the sham+vehicle group. ##P < 0.01 compared with the CIBP+vehicle group. n = 6 in each group).











A**B****C****D****E****F****G****H****I****J****K****L****M****N**





Highlights

1. The expression levels of GABABRs are significantly decreased in CIBP rats.
2. Restore diminished GABABRs attenuates CIBP-induced pain behaviors.
3. Inhibition the PKA/CREB signaling contributes to the analgesic effect of baclofen.