

UWL REPOSITORY

repository.uwl.ac.uk

Assessing the reliability of uptake and elimination kinetics modelling approaches for estimating bioconcentration factors in the freshwater invertebrate, *Gammarus pulex*

Miller, Thomas H., McEneff, Gillian ORCID logo [ORCID: https://orcid.org/0000-0003-2590-413X](https://orcid.org/0000-0003-2590-413X), Stott, Lucy C., Owen, Stewart F., Bury, Nicolas R. and Barron, Leon (2016) Assessing the reliability of uptake and elimination kinetics modelling approaches for estimating bioconcentration factors in the freshwater invertebrate, *Gammarus pulex*. *Science of The Total Environment*, 547. pp. 396-404. ISSN 0048-9697

<http://dx.doi.org/10.1016/j.scitotenv.2015.12.145>

This is the Published Version of the final output.

UWL repository link: <https://repository.uwl.ac.uk/id/eprint/3694/>

Alternative formats: If you require this document in an alternative format, please contact: open.research@uwl.ac.uk

Copyright: Creative Commons: Attribution 4.0

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy: If you believe that this document breaches copyright, please contact us at open.research@uwl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Assessing the reliability of uptake and elimination kinetics modelling approaches for estimating bioconcentration factors in the freshwater invertebrate, *Gammarus pulex*

Thomas H. Miller^{a,b}, Gillian L. McEneff^a, Lucy C. Stott^c, Stewart F. Owen^b, Nicolas R. Bury^{c,1}, Leon P. Barron^{a,*,1}

^a Analytical & Environmental Sciences Division, Faculty of Life Sciences and Medicine, King's College London, 150 Stamford Street, London SE1 9NH, United Kingdom


^b AstraZeneca, Global Environment, Alderley Park, Macclesfield, Cheshire SK10 4TF, United Kingdom

^c Division of Diabetes and Nutritional Sciences, Faculty of Life Sciences and Medicine, King's College London, Franklin Wilkins Building, 150 Stamford Street, London SE1 9NH, United Kingdom

HIGHLIGHTS

- Toxicokinetics for eight pharmaceuticals are presented in *Gammarus pulex*.
- Bioconcentration factors ranged from 12 to 4533 and depended on the method used.
- Decreasing trends in the uptake (k_1) rate constants were observed.
- Recognised models and their assumptions may lead to BCF estimation inaccuracies.
- Meta-analysis of previous toxicokinetic work revealed similar trends in k_1 .

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 13 October 2015

Received in revised form 18 December 2015

Accepted 29 December 2015

Available online 12 January 2016

Editor: Kevin V. Thomas

Keywords:

Pharmaceuticals

Pesticides

Toxicokinetics

Bioconcentration

Invertebrates

ABSTRACT

This study considers whether the current standard toxicokinetic methods are an accurate and applicable assessment of xenobiotic exposure in an aquatic freshwater invertebrate. An *in vivo* exposure examined the uptake and elimination kinetics for eight pharmaceutical compounds in the amphipod crustacean, *Gammarus pulex* by measuring their concentrations in both biological material and in the exposure medium over a 96 h period. Selected pharmaceuticals included two anti-inflammatories (diclofenac and ibuprofen), two beta-blockers (propranolol and metoprolol), an anti-depressant (imipramine), an anti-histamine (ranitidine) and two beta-agonists (formoterol and terbutaline). Kinetic bioconcentration factors (BCFs) for the selected pharmaceuticals were derived from a first-order one-compartment model using either the simultaneous or sequential modelling methods. Using the simultaneous method for parameter estimation, BCF values ranged from 12 to 212. In contrast, the sequential method for parameter estimation resulted in bioconcentration factors ranging from 19 to 4533. Observed toxicokinetic plots showed statistically significant lack-of-fits and further interrogation of the models revealed a decreasing trend in the uptake rate constant over time for ranitidine, diclofenac, imipramine, metoprolol, formoterol and terbutaline. Previous published toxicokinetic data for 14 organic micro-pollutants

* Corresponding author.

E-mail address: leon.barron@kcl.ac.uk (L.P. Barron).

¹ Both of these authors co-led this work.

were also assessed and similar trends were identified to those observed in this study. The decreasing trend of the uptake rate constant over time highlights the need to interpret modelled data more comprehensively to ensure uncertainties associated with uptake and elimination parameters for determining bioconcentration factors are minimised.

© 2016 Elsevier B.V. All rights reserved.

1. Introduction

The pseudo-persistent nature of pharmaceuticals and personal care products (PPCPs) has been highlighted in recent years as an environmental concern and has led to the introduction of a watch list under the EU Water Framework Directive which includes an anti-inflammatory, diclofenac, and two hormones, the synthetic ethinyl-estradiol (EE2) and natural estradiol (2013/39/EU, 2013). Several thousand PPCPs are currently available worldwide and whilst measured environmental concentrations typically range from low ng L^{-1} to high $\mu\text{g L}^{-1}$, their potential to effect an ecotoxicological response and/or bioaccumulate in a range of biota still remains understudied (De Lange et al., 2006; Contardo-Jara et al., 2011).

Ecotoxicological studies have shown that measured PPCP concentrations in surface waters would be highly unlikely to cause acute effects on exposed organisms (Crane et al., 2006). However, chronic exposure has been linked to behavioural activity changes, increased oxidative stress and alterations to the function of several vital organs in fish and invertebrates (Heckmann et al., 2007; Fernández et al., 2013). Aquatic invertebrates such as molluscs and smaller crustacean species have been previously utilised for monitoring PPCPs in the natural aquatic environment. Most recently, the freshwater amphipod, *Gammarus pulex*, was found to contain residues of carbamazepine, diazepam, nimesulide, trimethoprim and warfarin measuring at low ng g^{-1} concentrations in UK streams (Miller et al., 2015). PPCP uptake has also been previously observed at low ng g^{-1} concentrations in wild and caged mussel species collected from the coast of Ireland, the Bohai Sea in China, the Mediterranean Sea and San Francisco Bay, highlighting the extent of PPCP contamination worldwide (McEneff et al., 2014; Li et al., 2012; Bueno et al., 2013; Klosterhaus et al., 2013). EU Directive 93/39/EEC requires an environmental risk assessment to be carried out prior to drug licencing in order to determine any significant toxicological risks associated with a xenobiotic (Straub, 2002). Under the regulatory guidelines, environmental toxicity testing of pharmaceuticals requires standard acute toxicity tests, such as LC_{50} testing, to be carried out unless the predicted environmental concentration (PEC)/predicted no effect concentration (PNEC) ratio is <1 , whereby no further toxicity testing is required. Standardised toxicity tests on aquatic organisms are generally limited to algae (*Desmodesmus subspicatus* or *Pseudokirchneriella subcapitata*), *Daphnia magna* and/or fish (e.g. *Danio rerio*) considered as good model species from freshwater environments. Furthermore, a lack of published research generally exists on the uptake and depuration kinetics of PPCPs both in target and non-target aquatic species to help elucidate potential acute versus chronic effects.

Toxicokinetic studies identify whether a compound will accumulate to potentially toxic levels in/on the organism itself over time or potentially act as a source of toxicity in higher trophic organisms (Ashauer & Escher, 2010). In aquatic species, this can involve the study of either accumulation of compounds via water exposure only (i.e. bioconcentration), or via exposure through both water and diet (i.e. bioaccumulation or biomagnification) (Oliver & Niimi, 1983; Meador et al., 1995). Fish exposure studies often allow a time period for the compound of interest to reach steady-state within the organism, where the rate of uptake is equal to the rate of depuration. However, this time can vary considerably and has led to the application of kinetic modelling where uptake and elimination rates are estimated and used to derive a bioconcentration factor (BCF) (Veith et al., 1979). This factor can be determined in two ways: (a) as a ratio of either the compound concentrations in the organism and the water phase at steady-state, or

(b) as the ratio of the uptake (k_1) and elimination (k_2) rate constants (Kenaga, 1972). This approach has been widely evaluated in the literature. Earlier models, such as those for methylmercury in fish (Norstrom et al., 1976) considered several variables including volume of water passing the gills, assimilation across the gills and body weight of the organism. More recent models have been developed to also account for water-phase and lipid-phase resistance, fish lipid content and compound $\log K_{ow}$ (Veith et al., 1979; Gobas & MacKay, 1987; Hendriks & Heikens, 2001). A widely known and accepted model used to calculate the bioaccumulation of a compound in fish via aqueous and dietary exposure is outlined in the Organisation for Economic Co-Operation and Development (OECD) 305 guidelines (OECD, Test No. 305). These guidelines present two methods for estimation of k_1 and k_2 . The sequential method can be performed in one of two ways, a k_2 value can be estimated by linear regression and then curve fitting methods are applied to find k_1 . Alternatively, curve fitting methods can be used to estimate k_2 first which is then used to estimate k_1 . The simultaneous model calculates both k_1 and k_2 together and is considered a potentially more reliable and realistic model for concurrent uptake and elimination processes occurring in biological systems. Considering the number of PPCPs available on the market that may require testing under EU REACH legislation (European Commission, 2006), the time scales (2 week acclimatisation followed by 28 days for the uptake phase alone unless steady-state is achieved sooner) and number of organisms required for each test ($n = 4$ per time-point for each exposure) the testing regime to apply to all chemicals under REACH would appear unfeasible. Furthermore, current policy aims to reduce the number of fish used for scientific research, thus current methods proposed such as the OECD guidelines should account for this more ethical approach (Carter et al., 2014; Browne, 2013). Several recent studies assessing BCF have utilised shorter exposure times with experiments lasting only 4–7 days using aquatic invertebrates as a means to assess the potential for substance to bioaccumulate in aquatic organisms (Ashauer et al., 2006; Meredith-Williams et al., 2012). These ecotoxicological studies are important to direct future risk assessment and essential when considering contaminant monitoring in water, sediment and biota.

In the present study, an *in vivo* experiment was carried out to determine the uptake and depuration kinetics of environmentally relevant (low $\mu\text{g} \cdot \text{L}^{-1}$) concentrations of several selected PPCPs in the common freshwater invertebrate, *G. pulex*, using radioactive labels and liquid scintillation analysis. Lastly, the OECD 305 guidelines currently used for modelling of uptake and elimination kinetics in aquatic species are critically evaluated for the first time based on the results obtained both in this study and other published works on micropollutants.

2. Materials and methods

2.1. Reagents, chemicals and consumables

Radio-labelled pharmaceuticals including ^3H -propranolol hydrochloride ($29.0 \text{ Ci mmol}^{-1}$) were acquired from Amersham Biosciences. ^3H -metoprolol ($29.7 \text{ Ci mmol}^{-1}$), ^3H -formoterol ($18.5 \text{ Ci mmol}^{-1}$) and ^3H -terbutaline ($29.0 \text{ Ci mmol}^{-1}$) were obtained from Vitrex. ^{14}C -ibuprofen ($2.03 \text{ Ci mmol}^{-1}$) was obtained from American Radiolabelled Chemicals Inc. (St Louis, US). ^3H -ranitidine (2.5 Ci mmol^{-1}) was obtained from Moravek Biochemicals, ^{14}C -diclofenac ($0.063 \text{ Ci mmol}^{-1}$) and ^3H -imipramine hydrochloride ($48.5 \text{ Ci mmol}^{-1}$) from Perkin-Elmer. All stock solutions were stored in ethanol. Hydrogen peroxide solution

(30% w/w) and analytical grade salts (>99%) including sodium hydrogen carbonate, magnesium sulphate, calcium sulphate, potassium chloride were purchased from Sigma (Dorset, UK). Tissue solubiliser (Solvable™) and liquid scintillation cocktail (Hionic Fluor™) were purchased from Fischer Scientific Ltd. (Loughborough, UK). Ultra-pure water was obtained from a Millipore Milli-Q water purification system with a specific resistance of 18.2 MΩ·cm or greater (Millipore, Bedford, MA, USA). 6-Well culture plates were obtained from VWR (Leicestershire, UK).

2.2. Sample collection and culture maintenance

G. pulex were collected by kick-sampling from the River Cray, South-East London, UK, 51°23'09.5"N 0°06'32.4"E. This site was previously shown to have low pharmaceutical contamination in both collected surface water and animal samples (Miller et al., 2015). The populations were transported to the laboratory in 500 mL Nalgene™ flasks filled with surface water from the sample collection site. Populations were rinsed with artificial freshwater (AFW) and then acclimatised to laboratory conditions (as specified below) for a minimum of 7 days before any exposure experiments were performed. AFW was prepared from 1.15 mM of NaHCO₃, 0.50 mM MgSO₄, 0.44 mM CaSO₄ and 0.05 mM of KCl dissolved in 20 L of ultra-pure water. This water was subsequently aerated for several hours to remove dissolved carbonic acid and maximise the dissolved oxygen concentrations. Each culture tank (n = 8) was filled with 2.5 L of AFW and animals were fed with alder leaves that were previously collected from the sampling site and conditioned by submersion in surface water for two days prior to use.

2.3. Toxicokinetic exposure and conditions

Toxicokinetic experiments were performed separately for each pharmaceutical for a total of 96 h which included a 48 h uptake phase followed by a 48 h depuration period. Individual adult organisms, both male and female and each >5 mg wet weight, were placed in each well of 6-well culture plates. *G. pulex* were carefully transferred to well plates using blunt forceps to avoid any harm to the organisms before exposure. A single well contained one organism in 10 mL of exposure media (AFW and test compound) and only non-parasitised individuals were used (absence of *Pomphorhynchus laevis* indicated by the lack of an orange dot on the dorsal side of the animal). *G. pulex* were exposed to individual PPCPs at a concentration of 1 µg·L⁻¹, except for diclofenac and ibuprofen which were present at 10 µg·L⁻¹. The higher exposure of these two compounds was due to the low activity of the radiolabel. All exposure media contained <0.05% of solvent (ethanol). A total of 33 organisms were used per exposure and were sampled (n = 3/time-point) at 2, 5, 18, 24 and 48 h in the uptake phase followed by the same time-points in the depuration phase. Along with *G. pulex*, 50 µL water was also sampled from each well for analysis of radioactivity. Each sampled organism was washed in 10 mL of ultra-pure water for 10 s (n = 6) and gently blotted dry to remove any excess exposure media and unbound compound to the cuticle of the animal. Organisms were weighed after sampling to determine body mass and then transferred to scintillation tubes for tissue solubilisation. Three individual organisms were also exposed to unspiked AFW in culture plates and sampled after 96 h in a control experiment to account for any background radiation. Additionally, for each experiment, three wells without *G. pulex* were filled with exposure media to account for losses of the compound by sorption to the walls of culture plates. Culture plates were stored in sealed plastic containers with wet tissue to prevent evaporative losses during the static exposure. The light cycle followed 12:12 h light:dark without a dusk/dawn transition period. All experiments were performed in a temperature controlled room at 15 °C (± 2 °C) and water pH was also measured across each experiment at 8.2 ± 0.1.

2.4. Sample preparation and liquid scintillation counting

Water samples (50 µL) collected from each exposure well were added to 2 mL of Hionic Fluor liquid scintillation cocktail and counted for radioactivity on a Beckman LS6500 instrument (Beckman Coulter, Inc.). Sampled *G. pulex* individuals were placed in a scintillation tube with 2 mL of tissue solubiliser and maintained at room temperature (approx. 20 °C) for 96 h. Samples were shaken vigorously and then a 50 µL aliquot of the solubilised biotic extract was added to 2 mL of Hionic Fluor to be counted. To account for any difference in counts caused by colour quenching, hydrogen peroxide (200 µL) was added to a previously counted biotic extract and re-analysed. No difference in counts was observed with or without the presence of hydrogen peroxide, therefore, all other biotic samples were counted without the addition of hydrogen peroxide. In addition, chemiluminescence accounted for <0.01% of the overall counts, and was therefore ignored.

2.5. Modelling bioconcentration factors

Parameter estimation of uptake rate constant (k_1) and depuration rate constant (k_2) was performed using a curve fitting algorithm via Minitab statistical software (Minitab Ltd., Coventry, UK) and as outlined in the OECD 305 Fish Bioconcentration Guidelines (OECD, Test No. 305). The concentration of compound in the organism is assumed to follow first order kinetics and is expressed in Eq. (1),

$$\frac{dC_{\text{organism}}}{dt} = k_1 \times [C_{\text{water}}] - k_2 \times [C_{\text{organism}}] \quad (1)$$

where, dC_{organism}/dt is the rate of change in the concentration of a compound within/on *G. pulex* (mg kg⁻¹ day⁻¹), k_1 is the uptake rate constant (L kg⁻¹ day⁻¹), k_2 is the elimination rate constant (day⁻¹), C_{water} is the concentration in the water (mg L⁻¹) and C_{organism} is the concentration in the organism (mg kg⁻¹). Eq. (1) was integrated into Eqs. (2) and (3) for fitting of curves to the uptake and depuration data. This method, known as the Levenberg–Marquardt algorithm, uses an iterative formula to minimise the residual errors between the observed and predicted data points and simultaneously estimates k_1 and k_2 values from the fitted curve i.e.

$$[C_{\text{organism}}] = [C_{\text{water}}] \times \frac{k_1}{k_2} \times (1 - e^{-k_2 t}), \quad \text{when } 0 < t < t_e \quad (2)$$

$$[C_{\text{organism}}] = [C_{\text{water}}] \times \frac{k_1}{k_2} \times (1 - e^{-k_2(t-t_e)} - e^{-k_2 t}), \quad \text{when } t > t_e \quad (3)$$

where, t is the time (days) and t_e is the end time of the uptake phase (days). At steady-state, the rate of uptake should be equal to the rate of depuration and there should be no overall change in analyte concentration within *G. pulex*, as expressed by Eq. (4),

$$k_1 \times [C_{\text{water}}] = k_2 \times [C_{\text{organism}}] \leftrightarrow \frac{k_1}{k_2} = \frac{[C_{\text{fish}}]}{[C_{\text{water}}]} = \text{BCF} \quad (4)$$

where, BCF is the bioconcentration factor (L kg⁻¹). BCF can also be estimated using a sequential method where a simple linear regression model is developed based on the depuration data only. With the assumption of first order kinetics, the model should fit a straight line and its slope represents the elimination rate constant as shown in Eq. (5), i.e.

$$\ln [C_{\text{organism}}] = -k_2 \times t + c \quad (5)$$

where, $\ln[C_{\text{organism}}]$ is the natural log of the analyte concentration within *G. pulex* and c is the intercept, which here equals the natural log of the analyte concentration in the *G. pulex* at the start of the depuration phase. The k_2 from Eq. (5) can then be used as a parameter in the

curve fitting algorithm to estimate k_1 . The rearrangement of Eq. (2) allows the value for k_1 to be calculated over the time interval specified, as shown in Eq. (6) (Crookes & Brooke, 2011). The assumptions of the equation are that analyte concentration in the water and k_2 remain constant. The k_2 used in Eq. (6) was directly estimated by using linear regression of the depuration data to obtain the slope (k_2). The value of k_1 should remain constant over the entire experiment.

$$k_1 = \frac{[C_{\text{organism}}] \times k_2}{[C_{\text{water}}] \times (1 - e^{-k_2 \times t})} \quad (6)$$

For this study, initial parameters for k_1 and k_2 were arbitrarily set at 0.1 in the software with C_{water} set in $\mu\text{g L}^{-1}$, t set at 48 h and the maximum number of iterations was set at 200 upon which optimised k_1 and k_2 values were subsequently derived. Confidence intervals (95%) were plotted for curves and the overall model fits were assessed. The lack-of-fit test was calculated in the Minitab software and was used to assess the fit of the line by comparing the variation in response of the replicate data. Lack-of-fit was assessed at a significance level of 0.05. Correlation coefficients (r^2) were evaluated when the sequential method was used to estimate k_2 . The distribution coefficient ($\log D$) was generated using ACD Labs Percepta software for the interpretation of estimated BCF values. All compound information is displayed in Table S2 of the SI.

3. Results and discussion

3.1. Uptake and elimination kinetics for selected PPCPs within *G. pulex*

The exposure concentration of each PPCP was selected to approximate the higher ranges of trace pharmaceutical occurrence in the aquatic environment to maintain practically quantifiable limits for reliable analysis (Miller et al., 2015; Hilton & Thomas, 2003; Thomas & Hilton, 2004). Considering that natural uptake and depuration are not separate processes, the BCF values for the selected compounds were determined using the simultaneous model described above (Table 1). Uptake of each pharmaceutical was observed in *G. pulex* as early as 2 h from the point of exposure. The highest residue concentrations measured in *G. pulex* at the 48 h timepoint were ibuprofen and diclofenac, potentially corresponding to the elevated exposure concentrations of $10 \mu\text{g L}^{-1}$. All other compounds exposed at $1 \mu\text{g L}^{-1}$ measured $<80 \text{ ng g}^{-1}$ ww after 48 h uptake (Fig. 1). The rate of PPCP uptake measured in the exposed *G. pulex* corresponds to the decreases in PPCP concentration measured in the spiked AFW. The largest decrease in PPCP concentration was observed for imipramine, where analyte concentrations in the water decreased to an average of $0.478 \mu\text{g L}^{-1}$ corresponding to a 52.2% loss. After 48 h, formoterol concentration also decreased in water by 15% to an average of $0.85 \mu\text{g L}^{-1}$. The exposure concentrations of the remaining compounds did not decrease by $\geq 10\%$ (Fig. 2 and Table S1). Additional sources of potential PPCP loss in the aqueous phase should be mentioned and include photolysis, volatilisation, metabolism by microorganisms and sorption to the walls of the exposure well. Of these processes sorption was accounted for by control wells with exposure media only and was shown to account for negligible losses in water concentration except in the case of imipramine (Table S1). Within 2 h, there was a 27% loss of imipramine and within 48 h the loss increased to 39%. As quantification was performed by LSC, any degradation products resulting from transformation or photolysis would contribute towards the total radioactivity and counted as the precursor compound. However, it should be considered that these formed products may potentially have different accumulation potentials and hence latent uptake and elimination kinetics.

Following removal from the contaminated source, relatively high elimination rates were measured for most of the selected compounds. However, imipramine showed increased uptake ($k_1 = 1.408 \text{ L kg}^{-1} \text{ day}^{-1}$), but lower elimination ($k_2 = 0.007 \text{ day}^{-1}$), resulting in the highest BCF value measured at 212. Diclofenac has the

Table 1
Toxicokinetic parameters and bioconcentration factors for eight PPCPs.

Compound	Simultaneous BCF					Sequential BCF ^a					Sequential BCF ^b				
	k_1 ($\text{L kg}^{-1} \text{ day}^{-1}$)	SE	k_2 (day^{-1})	SE	p-Value	BCF	k_1 ($\text{L kg}^{-1} \text{ day}^{-1}$)	SE	p-Value	BCF	k_1 ($\text{L kg}^{-1} \text{ day}^{-1}$)	SE	p-Value	k_2 (day^{-1})	r^2
Propranolol	0.538	0.068	0.017	0.004	0.266	32	0.618	0.047	0.831	39	0.604	0.045	0.860	0.015	0.490
Formoterol	0.408	0.093	0.029	0.009	0.335	14	0.451	0.051	0.914	18	0.357	0.040	0.942	0.011	0.121
Imipramine	1.408	0.205	0.007	0.004	0.008	212	1.361	0.177	0.017	3811	1.360	0.177	0.017	0.000	0.001
Metoprolol	0.076	0.022	0.005	0.008	0.073	16	N/A								
Terbutaline	0.136	0.020	0.011	0.004	0.026	12	0.135	0.016	0.027	22	0.117	0.016	0.027	0.006	0.200
Ranitidine	0.479	0.126	0.028	0.011	0.005	17	0.310	0.071	0.007	81	0.301	0.070	0.007	0.003	0.015
Diclofenac	0.273	0.037	0.020	0.005	0.002	14	0.253	0.025	0.017	27	0.269	0.026	0.023	0.013	0.243
Ibuprofen	0.338	0.094	0.012	0.008	0.000	27	0.582	0.097	0.013	27	0.488	0.073	0.029	0.010	0.093

p-Values were assessed via standard error (SE) and lack-of-fit tests.

^a Sequential using curve fitting method.

^b Sequential using linear regression.




Fig. 1. Uptake and elimination data for PPCPs in *G. pulex*. Dashed lines indicate 95% confidence limits.




Fig. 2. Relationship of uptake rate constants (k_1) over time for eight PPCPs (black circles) and the respective water concentrations (C_w) over time (crosses).

same $\log P$ value as imipramine at 4.4 ($\log D_{8.2} = -1.1$) but attained a significantly lower BCF value of 14 due to its high rate of elimination. Ibuprofen, another acidic drug with a $\log P$ of 3.5 (calculated $\log D_{8.2} = -0.1$), also had a low BCF value determined at 27. The BCF values for the four compounds with $\log P < 2$. (i.e. metoprolol, ranitidine,

terbutaline and formoterol) were determined between 12 and 17. Hydrophobicity is generally considered a major factor when determining the bioaccumulation potential of a compound. However, uptake studies related to pharmaceuticals in several species of plants, for example, showed poor correlations between $\log D_{ow}$ and $\log BCF$ and especially

so for ionised molecules (Wu et al., 2013). Low bioconcentration of the selected PPCPs was in agreement with a study by Meredith-Williams et al., in which toxicokinetic data for six pharmaceuticals within *G. pulex* was shown with the exception of fluoxetine, a selective serotonin reuptake inhibitor ($BCF = 185,900$) (Meredith-Williams et al., 2012). In the cases of diclofenac, ibuprofen, imipramine and propranolol, $\log P$ is similar (3.3–4.7). Therefore, using an uptake model based on hydrophobicity, it would be logical to assume similar uptake rates. A potential reason for their difference could be physicochemical in nature, e.g. due to their anionic or cationic nature as well as the degree of ionisation and $\log D_{ow}$ value (Erickson et al., 2006). It could also be due to biological factors such as gill surface charge or the boundary layer between the bulk water and the gill surface (Tao et al., 2001). Uptake across the gill may also occur by more than simple passive diffusion for these ionic compounds and thus carrier mediated transport may also have influence on the different ionic species (Sugano et al., 2010; Kell & Oliver, 2014). The increased uptake constants of imipramine, propranolol and formoterol are in agreement with reported gill cell permeabilities to these compounds in the same order of imipramine > propranolol > formoterol (Stott et al., 2015). The low concentrations of PPCP residues measured in the *G. pulex* and unspiked AFW post-exposure highlights the ability for *G. pulex* to readily metabolise and eliminate xenobiotics, as previously shown by Nyman et al. (2014) and Ashauer et al. (2012). This evidence suggests there is conservation of cytochrome P450 enzymes, similarly observed in other aquatic invertebrate species (Solé & Livingstone, 2005).

3.2. Comparison of simultaneous versus sequential uptake and depuration process models

Methods used for the calculation of BCF values in *G. pulex* are summarised in Table 1 and also include uptake and elimination constants (\pm standard error). Many of the toxicokinetic plots in Fig. 1 are shown to have some lack-of-fit. The p -value generated from a lack-of-fit test shows that in the simultaneous method there are 5 models that have a statistically significant lack-of-fit indicating potentially inaccurate and unreliable BCF values. It is possible that several large outliers could influence the lack-of-fit test, thus resulting in a statistical significance when potentially none exists. When using the simultaneous method, if a poor fit exists, then the sequential method should be investigated as a potential alternative. The linear regression of the depuration phase data points gives a direct estimate of k_2 . The goodness-of-fit is interpreted by visual inspection of the linearity and the r^2 (Fig. S1). Consideration of the sequential method showed an over-estimation of BCF values when compared to the simultaneous model. Deviations from linearity can indicate higher order kinetics. Simple plots of $1/[C_{organism}]$

here did not indicate second order kinetics and therefore k_2 values from plots of $\ln[C_{organism}]$ were accepted. Low r^2 values for some compounds were likely due to the scatter in measured internal concentrations. Comparison of derived k_2 values showed that imipramine, formoterol, ranitidine, diclofenac and terbutaline had significantly lower elimination constants in comparison to the simultaneous model approach (Table 1). Markedly reduced estimations in k_2 for imipramine and ranitidine corresponded to a large increase in BCF for ranitidine, increasing 4-fold, and imipramine, increasing >10-fold to ~4200 on average between curve fitting and linear regression approaches. Given the inherent non-standard method we have applied, further work would be necessary to better understand this apparently high BCF and we would caution reliance on this value from such a limited study. When using a curve fitting method to calculate an elimination constant in the sequential method there was good agreement between the linear regression estimates of k_2 , indicating the estimate of k_2 was correct. The p -values for the curve fits indicated that there was only one statistical lack-of-fit for the k_2 value generated for ibuprofen. Uptake curves displayed a poor fit (as shown for imipramine concentration in *G. pulex*, which was consistently under-estimated). In addition, uptake constants in Table 1 specifically showed significant lack-of-fit for all compounds except propranolol and formoterol (p -value > 0.05). In fitting the depuration data using the sequential approach, a zero to mildly increasing slope was observed overall for metoprolol due to a wider scatter of data. A k_2 value could not therefore be calculated for metoprolol. The potential for model uncertainty highlighted in this study is significant from a regulatory perspective, especially for compounds such as imipramine that was determined to be accumulative using the sequential method and non-accumulative using the simultaneous method (European Commission, 2006).

3.3. Assessment of k_1 , k_2 and C_w constancy

The OECD 305 model makes several assumptions that C_w , k_1 and k_2 do not change over time. To assess the potential validity of the k_1 constancy assumption in the first instance, k_1 was derived at each time point accordingly (Crookes & Brooke, 2011). It should be noted that a potential limitation to this approach was that the equation to calculate k_1 uses the k_2 estimate from the depuration phase, but this was deemed sufficient to identify any trends in any variation observed. As the lack-of-fit tests of the simultaneous method showed significant lacks-of-fit a direct estimation of k_2 from linear regression is used in Eq. (6) for simplicity and increased reliability. When plotted against time (Fig. 2), a clear reduction in k_1 over the exposure period was observed (especially for imipramine and diclofenac). Some random variance was also observed, such as for propranolol, which resulted in a relatively constant

Table 2
Toxicokinetic parameters and standard errors (SE) for 14 organic micropollutants with bioconcentrations factors and lack-of-fit tests for each compound.

Compound	BAF ^a	Simultaneous BCF						Sequential BCF							
		<i>k</i> ₁ (L kg ⁻¹ day ⁻¹)	SE	<i>k</i> ₂ (day ⁻¹)	SE	<i>p</i> -Value	BCF	<i>k</i> ₁ (L kg ⁻¹ day ⁻¹)	SE	<i>p</i> -Value	<i>k</i> ₂ (day ⁻¹)	SE	<i>p</i> -Value	BCF	
4-Nitrobenzyl chloride	185	666	665.740	4.540	4.540	0.000	147	259	28.324	0.00	1.212	0.149	0.230	214	
2,4-Dichloroaniline	56	140	20.073	2.830	0.465	0.000	50	70	4.689	0.03	0.392	0.092	0.868	179	
2,4-Dichlorophenol	4466	600	34.196	0.066	0.024	0.000	9050	750	48.646	0.00	0.010	0.018	0.400	72,728	
4,6-Dinitro- <i>o</i> -cresol	37	39	2.610	1.146	0.124	0.070	34	37	1.595	1.00	0.729	0.077	0.033	51	
1,2,3-Trichlorobenzene	191	1142	403.773	10.648	3.781	0.513	107	167	32.257	0.06	0.475	0.300	0.986	351	
2,4,5-Trichlorophenol	2635	941	79.280	0.252	0.064	0.001	3729	1091	109.906	0.21	0.131	0.039	0.001	8327	
Aldicarb	2	16	1.421	10.419	0.938	0.000	2	3	0.245	0.00	0.936	0.140	0.003	3	
Carbofuran	65	10	0.355	0.146	0.019	0.026	68	10	0.570	0.11	0.140	0.019	0.025	72	
Diazinon	82	276	24.271	3.569	0.3264	0.005	77	161	10.418	0.00	1.590	0.287	0.259	101	
Ethylacrylate	87	110	12.139	1.594	0.2446	0.000	69	67	5.122	0.00	0.204	0.033	0.000	331	
Hexachlorobenzene	2915	553	36.8714	0.221	0.046	0.000	2505	631	56.518	0.00	0.152	0.031	0.637	4160	
Imidacloprid	7	2	0.093	0.265	0.038	0.000	7	2	0.117	0.02	0.175	0.022	0.000	13	
Malathion	114	86	6.782	0.721	0.113	0.000	120	80	5.059	0.02	0.378	0.072	0.001	212	
Sea nine	1732	755	57.821	0.303	0.065	0.000	2491	950	69.141	0.05	0.123	0.020	0.084	7696	

^a Reported from Ashauer et al. (2010).

average k_1 value of $0.58 (\pm 0.23)$, as would be expected. The simultaneous and sequential models estimated its k_1 value to be 0.54 and $0.62 \text{ L kg}^{-1} \text{ day}^{-1}$, respectively and therefore showed reasonable agreement. This observation is significant as propranolol showed no lack-of-fit in the uptake curve; therefore, the agreement indicates that a lack-of-fit arises from variable k_1 values over time.

This suggests that a decreasing k_1 trend is the cause of the poor model fits although it is possible that this may also be caused by a changing k_2 value (giving an apparent decrease in k_1) or variable exposure concentrations in the water. However, water was monitored during the course of the experiments to account for any losses (Fig. 2 & Table S1) and the only compound that showed any significant loss was imipramine ($>20\%$ nominal concentration). The k_1 and k_2 values should also be independent of pharmaceutical concentrations in the aqueous phase thus the trend observed is not in response to this variable (van Leeuwen & Vermeire, 2007). A change in k_2 is likely to be represented as a decrease over time (unless the compound induces its own metabolism) assuming growth has a negligible effect and therefore would not account for decreases in k_1 . The elimination curves also showed no lack-of-fit for 6 compounds and the linear regression showed no trends of changing k_2 values. The trend observed therefore is not in response to the parameters C_w or k_2 and we therefore suggest the variability in uptake (decreasing k_1) trend is the cause of the poor model fit.

3.4. Performance of OECD models using other micro-pollutant studies in *G. pulex*

G. pulex has been shown to metabolise organic compounds with low bioaccumulation factors previously observed (<1500) (Ashauer et al., 2012). As defined by Annex XIII of the REACH criteria, for a compound to be considered bioaccumulative the BCF/BAF should be >2000 (European Commission, 2006). Other work by Ashauer et al. investigated the toxicokinetics of 14 micro-pollutants in *G. pulex* and presented higher BCFs for three polychlorophenols in particular (Ashauer et al., 2010). As discussed by the authors, correlations showed an observable lack-of-fit in some cases. A slightly different model to the OECD 305 model was used in this work, where changes in C_w were accounted for as well as inclusion of an extra statistical algorithm to select the best parameter combinations of k_1 and k_2 . However, when applying the OECD 305 models to BAF prediction, it is important to understand whether this is likely to be inaccurate and, amongst other reasons, potentially due to variation in k_1 , k_2 or C_w . To determine if any similar trends could be identified in other published *G. pulex* toxicokinetics studies, raw data from Ashauer et al., was re-examined using the OECD 305 modelling approach and presented in Table 2 (Ashauer et al., 2010). Although the authors' experiments were originally designed for determination of bioaccumulation, the report showed this to account for a small percentage of accumulation. Therefore, feeding was not included in any calculations. Similar to our findings for pharmaceuticals, both models displayed a statistically significant lack-of-fit for these organic micro-pollutant compounds. When the sequential method was applied, better fits were obtained for the depuration phase in comparison to the uptake phase. However, despite models used herein not performing as well overall, there was good agreement between the predicted BCF values and those generated by Ashauer et al. The data was then used to plot k_1 versus time (Fig. S2), and again an obvious systematic decrease was observed for 9 out of 14 compounds. Statistical lack-of-fits ($p < 0.05$) were observed in the sequential uptake model especially for 4-nitrobenzylchloride, ethylacrylate, diazinon, aldicarb and hexachlorobenzene ($p < 0.001$). The latter two compounds were notable cases where the spread of replicate k_1 data at each time-point was especially narrow and so the trend in k_1 reduction over time was apparent. Of the remaining five compounds, trends in k_1 were less evident and were coupled with $p > 0.05$ for lack-of-fit for four compounds using the sequential uptake model. The remaining compound, 2,4-

dichlorophenol, showed no obvious trends in k_1 variance as the major reason for the observed lack-of-fit in the uptake phase. In summary, k_1 data could be considered reliable for only 5 of 14 compounds using the OECD 305 sequential model. In addition to the data of these 14 different organic pollutants, we also reassessed data from an exposure study of chlorpyrifos across 15 different invertebrate species to assess the issue more broadly (Rubach et al., 2010). Decreases in k_1 were observed in several species and the trend was somewhat similar, albeit with larger scatter of the data (Fig. S3). This also identifies a further limitation that metabolism is likely to affect k_1 and k_2 values thus the differences in k_1 constancy between organisms may be as a result of biotransformation. The study showed considerable differences between species in uptake and elimination rates showing that species type may affect the constancy of k_1 in particular and further studies are required using more compounds between different species to fully assess this possibility. If the assumption of k_1 constancy varies on a compound-by-compound basis, curve fitting methods to predict BCF are likely to be inherently inaccurate for environmental risk assessment purposes for *G. pulex*. Therefore, it is suggested that the approach taken herein (Crookes & Brooke, 2011) could be used to check the reliability of BCF data where a statistical lack-of-fit exists for this species.

Decreasing k_1 could be explained by several possible mechanisms. The first is that growth dilution could cause an apparent decrease in k_1 due to the mass of the organism increasing while the concentration of substance remains the same. However, this situation is unlikely given the short timescales of this work and that of Ashauer et al. (Ashauer et al., 2010). Therefore, growth of *G. pulex* is assumed to be negligible, particularly as this is regulated in line with their moulting cycle. However, further investigation would be required to fully support this. A second possibility is that *G. pulex* have been shown to alter respiration rates in the presence of a poor diet (Graça et al., 1993). As the animals were not fed during these experiments, it is possible that this slowed uptake. However, the toxicokinetic experiments by Ashauer et al. involved feeding organisms over their uptake period suggesting the uptake trend is not in response to diet induced factors (Ashauer et al., 2010). As these compounds are exposed to non-target animals, it is also possible that toxicodynamic effects could affect uptake, which is more easily interpreted using the dataset by Ashauer et al., where the exposure concentration was between 2 and 88 fold below the 24 h LC_{50} value. However for our dataset, mortality was not significantly higher than in controls for pharmaceuticals at the exposure concentrations used. Another consideration is that instantaneous sorption to the animal cuticle could account for the initially high k_1 constants. However, an examination of the decrease in uptake rate against $\log D$ and $\log P$ revealed no correlation and compounds displayed independent k_1 decreases (Fig. S4). However, $\log D$ only governs sorption to a certain extent and other physicochemical properties including polar/topological surface area, ionic state, amongst others, could influence sorption onto the exoskeleton. Where animals shed their exoskeleton during the exposure period, these were collected, weighed wet and radioactivity measured in a brief experiment. It was found that the maximum concentration of five of the eight pharmaceuticals on the exoskeleton material recovered did not exceed 24% of total compound mass in the animal in these cases (Table S3). Therefore, reduction in k_1 via this mechanism is indeed plausible, but extended measurements across more compounds, conditions and replicates are recommended for full characterisation of this process. The potential for sorption as the reason for changes in k_1 is not based on physiology, but rather on the physicochemical properties of the xenobiotic itself, suggesting that rate constant stability may be compound specific.

4. Conclusions

This work demonstrates the importance of data interpretation using multiple modelling methods to estimate BCFs. Specifically, the comparative assessment of model lack-of-fits for both simultaneous and

sequential models (where k_2 remains constant) is recommended to reliably estimate and to ensure the accuracy of xenobiotic risk assessments. A decreasing trend in the uptake rate constant over time was apparent which disrupted the validity of the standard model assumptions tested, and suggests that more complex models are needed to describe accumulation of xenobiotics in invertebrates, more particularly in *G. pulex*. Kinetic BCF/BAF are an estimate of steady-state values, but it is possible that these models are adequate enough to indicate whether a compound may have a potential to accumulate or not. It is now important to identify whether such trends are also observed more generally across different species as well as a fuller investigation into the roles sorption and metabolism have in these standard models.

Acknowledgements

This work was conducted under funding from the Biotechnology and Biological Sciences Research Council (BBSRC) CASE industrial scholarship scheme (Reference BB/K501177/1) and AstraZeneca Global SHE research programme. AstraZeneca is a biopharmaceutical company specialising in the discovery, development, manufacturing and marketing of prescription medicines, including some products measured here. Funding bodies played no role in the design of the study or decision to publish. The authors declare no financial conflict of interest.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.scitotenv.2015.12.145>.

References

- European Union water framework directive, Official Journal of the European Communities, OJL226.
- Ashauer, R., Escher, B.J., 2010. Advantages of toxicokinetic and toxicodynamic modelling in aquatic ecotoxicology and risk assessment. *J. Environ. Monit.* 12 (11), 2056–2061.
- Ashauer, R., Boxall, A., Brown, C., 2006. Uptake and elimination of chlorpyrifos and pentachlorophenol into the freshwater amphipod *Gammarus pulex*. *Arch. Environ. Contam. Toxicol.* 51 (4), 542–548.
- Ashauer, R., et al., 2010. Bioaccumulation kinetics of organic xenobiotic pollutants in the freshwater invertebrate *Gammarus pulex* modeled with prediction intervals. *Environ. Toxicol. Chem.* 29 (7), 1625–1636.
- Ashauer, R., et al., 2012. Significance of xenobiotic metabolism for bioaccumulation kinetics of organic chemicals in *Gammarus pulex*. *Environ. Sci. Technol.* 46 (6), 3498–3508.
- Browne, J., 2013. 2010 to 2015 Government Policy: Animal Research and Testing. Home Office.
- Bueno, M.J.M., et al., 2013. Fast and easy extraction combined with high resolution-mass spectrometry for residue analysis of two anticonvulsants and their transformation products in marine mussels. *J. Chromatogr. A* 1305, 27–34.
- Carter, L.J., et al., 2014. Minimised bioconcentration tests: a useful tool for assessing chemical uptake into terrestrial and aquatic invertebrates? *Environ. Sci. Technol.* 48 (22), 13497–13503.
- Contardo-Jara, V., et al., 2011. Exposure to human pharmaceuticals Carbamazepine, Ibuprofen and Bezafibrate causes molecular effects in *Dreissena polymorpha*. *Aquat. Toxicol.* 105 (3–4), 428–437.
- Crane, M., Watts, C., Boucard, T., 2006. Chronic aquatic environmental risks from exposure to human pharmaceuticals. *Sci. Total Environ.* 367 (1), 23–41.
- Crookes, M.J., Brooke, D.N., 2011. Estimation of Fish Bioconcentration Factor (BCF) From Depuration Data. Environment Agency, Bristol.
- De Lange, H.J., et al., 2006. Behavioural responses of *Gammarus pulex* (Crustacea, Amphipoda) to low concentrations of pharmaceuticals. *Aquat. Toxicol.* 78 (3), 209–216.
- Erickson, R.J., et al., 2006. Uptake and elimination of ionizable organic chemicals at fish gills: II. Observed and predicted effects of pH, alkalinity, and chemical properties. *Environ. Toxicol. Chem.* 25 (6), 1522–1532.
- European Commission, 2006. Regulation (EC) No 1907/2006 of the European Parliament and of the Council of 18 December 2006 concerning the Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH), establishing a European Chemicals Agency, amending Directive 1999/45/EC and repealing Council Regulation (EEC) No 793/93 and Commission Regulation (EC) No 1488/94 as well as Council Directive 76/769/EEC and Commission Directives 91/155/EEC, 93/67/EEC, 93/105/EC and 2000/21/EC. Off. J. Eur. Union L396, 1–849.
- Fernández, C., Carbonell, G., Babin, M., 2013. Effects of individual and a mixture of pharmaceuticals and personal-care products on cytotoxicity, EROD activity and ROS production in a rainbow trout gonadal cell line (RTG-2). *J. Appl. Toxicol.* 33 (11), 1203–1212.
- Gobas, F.A.P.C., MacKay, D., 1987. Dynamics of hydrophobic organic chemical bioconcentration in fish. *Environ. Toxicol. Chem.* 6 (7), 495–504.
- Graça, M.A.S., Maltby, L., Calow, P., 1993. Importance of fungi in the diet of *Gammarus pulex* and *Asellus aquaticus*. *Oecologia* 96 (3), 304–309.
- Heckmann, L.-H., et al., 2007. Chronic toxicity of ibuprofen to *Daphnia magna*: effects on life history traits and population dynamics. *Toxicol. Lett.* 172 (3), 137–145.
- Hendriks, A.J., Heikens, A., 2001. The power of size. 2. Rate constants and equilibrium ratios for accumulation of inorganic substances related to species weight. *Environ. Toxicol. Chem.* 20 (7), 1421–1437.
- Hilton, M.J., Thomas, K.V., 2003. Determination of selected human pharmaceutical compounds in effluent and surface water samples by high-performance liquid chromatography–electrospray tandem mass spectrometry. *J. Chromatogr. A* 1015 (1–2), 129–141.
- Kell, D.B., Oliver, S.G., 2014. How drugs get into cells: tested and testable predictions to help discriminate between transporter-mediated uptake and lipoidal bilayer diffusion. *Front. Pharmacol.* 5, 231.
- Kenaga, E., 1972. Guidelines for environmental study of pesticides: determination of bioconcentration potential. In: Gunther, F., Gunther, J. (Eds.), *Residue Reviews*. Springer, New York, pp. 73–113.
- Klosterhaus, S.L., et al., 2013. Method validation and reconnaissance of pharmaceuticals, personal care products, and alkylphenols in surface waters, sediments, and mussels in an urban estuary. *Environ. Int.* 54, 92–99.
- Li, W.H., et al., 2012. Investigation of antibiotics in mollusks from coastal waters in the Bohai Sea of China. *Environ. Pollut.* 162, 56–62.
- McNeff, G., et al., 2014. A year-long study of the spatial occurrence and relative distribution of pharmaceutical residues in sewage effluent, receiving marine waters and marine bivalves. *Sci. Total Environ.* 476, 317–326.
- Meador, J.P., et al., 1995. Bioaccumulation of polycyclic aromatic hydrocarbons by marine organisms. In: Ware, G. (Ed.), *Reviews of Environmental Contamination and Toxicology*. Springer, New York, pp. 79–165.
- Meredith-Williams, M., et al., 2012. Uptake and depuration of pharmaceuticals in aquatic invertebrates. *Environ. Pollut.* 165 (0), 250–258.
- Miller, T.H., et al., 2015. Pharmaceuticals in the freshwater invertebrate, *Gammarus pulex*, determined using pulverised liquid extraction, solid phase extraction and liquid chromatography–tandem mass spectrometry. *Sci. Total Environ.* 511 (0), 153–160.
- Norstrom, R.J., McKinnon, A.E., deFreitas, A.S.W., 1976. A bioenergetics-based model for pollutant accumulation by fish. Simulation of PCB and methylmercury residue levels in Ottawa River yellow perch (*Perca flavescens*). *J. Fish. Res. Board Can.* 33 (2), 248–267.
- Nyman, A.-M., Schirmer, K., Ashauer, R., 2014. Importance of toxicokinetics for interspecies variation in sensitivity to chemicals. *Environ. Sci. Technol.* 48 (10), 5946–5954.
- OECD, Test No. 305. Bioaccumulation in Fish: Aqueous and Dietary Exposure: OECD Publishing.
- Oliver, B.G., Niimi, A.J., 1983. Bioconcentration of chlorobenzenes from water by rainbow trout: correlations with partition coefficients and environmental residues. *Environ. Sci. Technol.* 17 (5), 287–291.
- Rubach, M.N., et al., 2010. Toxicokinetic variation in 15 freshwater arthropod species exposed to the insecticide chlorpyrifos. *Environ. Toxicol. Chem.* 29 (10), 2225–2234.
- Solé, M., Livingstone, D.R., 2005. Components of the cytochrome P450-dependent monooxygenase system and 'NADPH-independent benzo[a]pyrene hydroxylase' activity in a wide range of marine invertebrate species. *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* 141 (1), 20–31.
- Stott, L.C., et al., 2015. A primary fish gill cell culture model to assess pharmaceutical uptake and efflux: evidence for passive and facilitated transport. *Aquat. Toxicol.* 159 (0), 127–137.
- Straub, J.O., 2002. Environmental risk assessment for new human pharmaceuticals in the European Union according to the draft guideline/discussion paper of January 2001. *Toxicol. Lett.* 131 (1–2), 137–143.
- Sugano, K., et al., 2010. Coexistence of passive and carrier-mediated processes in drug transport. *Nat. Rev. Drug Discov.* 9 (8), 597–614.
- Tao, S., et al., 2001. Simulation of acid–base condition and copper speciation in the fish gill microenvironment. *Comput. Chem.* 25 (3), 215–222.
- Thomas, K.V., Hilton, M.J., 2004. The occurrence of selected human pharmaceutical compounds in UK estuaries. *Mar. Pollut. Bull.* 49 (5–6), 436–444.
- van Leeuwen, C.J., Vermeire, T., 2007. *Risk Assessment of Chemicals: An Introduction*. Springer Science & Business Media, New York.
- Veith, G.D., Defoe, D.L., Bergstedt, B.V., 1979. Measuring and estimating the bioconcentration factor of chemicals in fish. *J. Fish. Res. Board Can.* 36 (9), 1040–1048.
- Wu, X., et al., 2013. Comparative uptake and translocation of pharmaceutical and personal care products (PPCPs) by common vegetables. *Environ. Int.* 60 (0), 15–22.