

UWL REPOSITORY
repository.uwl.ac.uk

Waterborne protozoan pathogens in environmental aquatic biofilms:
implications for water quality assessment strategies

Masangkay, Frederick, Milanez, Giovanni, Tsiami, Amalia ORCID logoORCID:
<https://orcid.org/0000-0002-1122-4814>, Hapan, Freida, Somsak, Voravuth, Kotepui, Manas
Kotepui, Tangpong, Jitbanjong and Karanis, Panagiotis (2020) Waterborne protozoan pathogens in
environmental aquatic biofilms: implications for water quality assessment strategies.
Environmental Pollution, 259. ISSN 0269-7491

<http://dx.doi.org/10.1016/j.envpol.2019.113903>

This is the Accepted Version of the final output.

UWL repository link: <https://repository.uwl.ac.uk/id/eprint/6661/>

Alternative formats: If you require this document in an alternative format, please contact:
open.research@uwl.ac.uk

Copyright: Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy: If you believe that this document breaches copyright, please contact us at open.research@uwl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Rights Retention Statement:

1 **Waterborne protozoan pathogens in environmental aquatic biofilms:**
2 **Implications for water quality assessment strategies**

3
4 Frederick R. Masangkay^{a,b*}, Giovanni D. Milanez^{a,b}, Amalia Tsiami^c, Freida Z. Hapan^d,
5 Voravuth Somsak^a, Manas Kotepui^a, Jitbanjong Tangpong^{a*}, Panagiotis Karanis^{e,f}
6

7 ^aBiomedical Sciences Program, School of Allied Health Sciences, Walailak University, Nakhon
8 Si Thammarat, 80160, Thailand

9 ^bDepartment of Medical Technology, Institute of Arts and Sciences, Far Eastern University
10 Manila, 1015, Philippines

11 ^cLondon Geller College of Hospitality and Tourism, University of West London, St Mary's
12 Road, Ealing, London, W5 5RF, United Kingdom^d

13 ^dDepartment of Medical Technology, College of Pharmacy, The Pontifical and Royal University
14 of Santo Tomas, España Blvd, Sampaloc, Manila, 1008, Philippines

15 ^eMedical Faculty and University Hospital of Cologne, Cologne, 50923, Germany

16 ^fUniversity of Nicosia Medical School, Nicosia, 2408, Cyprus
17
18
19
20
21
22

23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

Abstract

Biofilms containing pathogenic organisms from the water supply are a potential source of protozoan parasite outbreaks and a general public health concern. The aim of the present study was to demonstrate the simultaneous and multi-spatial occurrence of waterborne protozoan pathogens (WBPP) in substrate-associated biofilms (SAB) and compare it to surface water (SW) and sediments with bottom water (BW) counterparts using manual filtration and elution from low-volume samples. For scenario purposes, simulated environmental biofilm contamination was created from *in-situ* grown one-month-old SAB (OM-SAB) that were spiked with *Cryptosporidium parvum* oocysts. Samples were collected from the largest freshwater reservoirs in Luzon, Philippines and a University Lake in Thailand. A total of 69 samples (23 SAB, 23 SW, and 23 BW) were evaluated using traditional staining techniques for *Cryptosporidium*, and immunofluorescence staining for the simultaneous detection of *Cryptosporidium* and *Giardia*. In the present study, WBPP was found in 43% SAB, 39% SW, and 39% BW samples tested with SAB results reflecting SW and BW results. Further, the potential and advantages of using low-volume sampling for the detection of parasite (oo)cysts in aquatic matrices were also demonstrated. Scanning electron microscopy of OM-SAB revealed a naturally-associated testate amoeba shell, while *Cryptosporidium* oocysts spiked samples provided a visual profile of what can be expected from naturally contaminated biofilms. This study provides the first evidence for the simultaneous and multi-spatial occurrence of waterborne protozoan pathogens in low-volume

45 environmental aquatic matrices and warrants SAB testing along with SW and BW matrices for
46 improved water quality assessment strategies (iWQAS).

47

48 **Capsule**

49 Analysis of biofilms along with surface water and sediments in environmental aquatic systems
50 leads to improved detection and isolation of waterborne protozoan pathogens.

51

52 **Keywords:** Asia; biofilms; *Cryptosporidium*; *Giardia*; lakes

53

54 **1. Introduction**

55 *Cryptosporidium* and *Giardia* are protozoan agents of diarrheal outbreaks worldwide
56 (Baldursson and Karanis, 2011; Efstratiou et al., 2017a; Karanis et al., 2007; Mac Kenzie et al.,
57 1994). Cryptosporidiosis is the second leading global cause of infantile mortality next to rotavirus
58 infection (Checkley et al., 2015), while giardial infections are reported to be at more than 200
59 million cases annually worldwide. Both *Cryptosporidium* and *Giardia* are listed in the ‘Neglected
60 Diseases Initiative’ by the World Health Organization (Savioli et al., 2006). The
61 immunocompromised population is at highest risk in contracting these waterborne protozoan
62 pathogens (WBBP) with low infectious dose; ingestion of as few as 10 (oo)cysts have been
63 reported to cause morbidity even in the immunocompetent population (Dupont et al., 1995;
64 Okhuysen et al., 1999; Ortega et al., 1997; Steiner et al., 1997).

65 Biofilms are formed from the transition of planktonic cells to sessile cells, which leads to
66 the accumulation and establishment of stable interactions between pure and multi-species
67 communities (Matos et al., 2017; Suba and Masangkay, 2013). The complexity of the biofilm

68 community is brought about by the introduction of other unicellular organisms like protozoans
69 (Murphy et al., 2018; Waiser et al., 2016; Xu et al., 2014), which can facilitate the interspecies co-
70 evolutionary processes as recently reported and discussed in the biofilm inhabitation of
71 *Acanthamoeba* spp. infected with bacteria containing bacteriophages with the possibility of
72 virophages as well (Bekliz et al., 2016; Masangkay et al., 2018).

73 Biofilms are potential reservoirs of human and zoonotic pathogens in environmental
74 aquatic systems and contribute to the persistence and transmission of waterborne protozoan
75 pathogens and other microorganisms (Ryan et al., 2016). The biofilm roughness contributes to the
76 attachment of protozoan (oo)cysts (DiCesare et al., 2012a; Wolyniak et al., 2009; 2010) and the
77 biofilm mass as a whole provides UV protection for (oo)cysts trapped within its matrix which
78 contributes to the environmental persistence of potentially-pathogenic protozoans (DiCesare et al.,
79 2012b).

80 The aim of this study was to demonstrate the presence of *Cryptosporidium* and *Giardia*
81 (oo)cysts in environmental aquatic substrate-associated biofilms (SAB) and compare its results
82 against surface water (SW) and bottom water with sediment (BW) counterparts through manual
83 filtration and elution of low-volume samples and to introduce the significance of biofilms as a
84 biological reservoir for WBPP.

85

86 **2. Materials and Methods**

87 *2.1. Study sites*

88 The University Lake of Nakhon Si Thammarat, Thailand (Fig. 1.A) was chosen as one of
89 the study sites due to its recreational use and its importance as a habitat for local amphibians,
90 reptiles, small mammals, livestock, and aquatic birds. Fig. 1.B illustrates the other study sites in

91 the Philippines (Laguna de Bay, Taal Lake, Pantabangan watershed and The Seven Lakes of
92 Laguna, and Ipo watershed), four of which were tested for the presence of WBPP. The Ipo
93 watershed site was selected for the one-month-old SAB (OM-SAB) *in-situ* culture experiment.
94 Laguna de Bay is the largest lake in the Philippines and is the major site for freshwater aquaculture
95 in Luzon (Guzman, 2006; Israel, 2007). Taal Lake is the third largest lake in the country and the
96 primary source of freshwater fish in the surrounding provinces (Martinez, 2011). The Pantabangan
97 watershed was constructed by flooding a town and erecting a dam structure for hydroelectric
98 power. The Seven Lakes of San Pablo Laguna is a system of crater lakes, where Sampaloc and
99 Bunot Lake are used for aquaculture. Palakpakin Lake connects to a river system and Mohicap
100 Lake is enclosed by natural surroundings that offer tourists and local residents a breath-taking view
101 of the area. The remaining Yambo, Pandin, and Kalibato Lakes are mainly used for picnic and
102 water activities. The Ipo watershed has been created by a river system that drains the larger Angat
103 dam, which provides the majority of the water supply to Manila and its nearby metropolis.

104

105 2.2. Sample collection

106 Surface water samples (SW) were collected in 50 mL sterile polyethylene containers by
107 collecting no more than 30 cm below the water surface from each sampling area (Table 1). Bottom
108 water with sediments (BW) were collected from a water depth of one meter along the shoreline
109 that was composed of one-part sediments and four-parts bottom water. Substrate-associated
110 biofilms (SAB) were harvested from aquatic plants whenever present. Short segments of small
111 aquatic plants were collected and washed with sterile distilled water to remove non-adherent cells,
112 cut into smaller portions in order to fit loosely inside 50 mL sterile polyethylene containers with a
113 final volume of 50 mL sterile distilled water. In the absence of aquatic plants, adherent biofilms

114 (approximately 2 g) were scraped from rocks no more than 30 cm below the water surface. Samples
115 were transported to the laboratory and processed within 48 hours after collection. The 50 mL
116 samples in this study aimed to provide initial data on the capability and feasibility of detecting
117 pathogenic (oo)cysts from environmental aquatic matrices using a low-volume sampling
118 technique. The low-volume 50 mL sample was elected to represent the hypothetical volume of a
119 water sample that could be accidentally ingested or inhaled by an individual during water activities,
120 which could lead to the infection of *Cryptosporidium* and/or *Giardia* if present in the water source.

121

122 2.3. Sample processing

123 Each sample was vortexed for one minute to dislodge adherent cells from any larger
124 organic substances and debris in order to distribute the bio-colloids evenly throughout the sample
125 matrix and left to stand for 5 min to settle heavier solids. The SW, BW and SAB sample
126 suspensions were each manually filtered through a 1.2 µm glass microfiber filter fitted inside a 50
127 mL sterile disposable syringe (Masangkay et al., 2016). Glass microfiber filters were recovered
128 and placed in sterile disposable polyethylene plates, where the filtered sediments were scraped
129 using a sterile inoculating loop and 5 mL sterile distilled water as eluent. The 5 mL eluates were
130 transferred to sterile test tubes and centrifuged at 1500 g for 15 minutes (US EPA Method 1623,
131 2005) where 3 mL of the supernatants were discarded and the remaining 2 mL and pellet were
132 mixed to form a suspension and subsequently transferred and stored in microcentrifuge tubes for
133 smearing within 24 hours.

134

135 2.4. Microscopy of *Cryptosporidium* and *Giardia* (oo)cysts

136 For all samples, 25 μ L of the pellet suspension was made into a 1 cm diameter smear on a
137 clean glass slide in duplicate. Screening for *Cryptosporidium* oocysts was performed by staining
138 with modified Kinyoun's (MK), modified Safranin Methylene Blue (SMB), and Auramine (Aura).
139 Microscopic confirmation of *Cryptosporidium* spp. and *Giardia* spp. (oo)cysts were performed by
140 Direct Antibody Fluorescent Testing (IFT) using the Aqua-Glo™ G/C Direct Comprehensive Kit
141 (Waterborne Inc. USA) according to the manufacturer's instructions. Light microscopy of MK
142 and SMB smears was performed by examining 200 oil immersion fields using a Nikon Model
143 Eclipse E100LED light microscope. An upright epifluorescence incident light excitation trinocular
144 UB microscope with a three-megapixel camera was used to examine 200 high power fields of the
145 Aura and IFT smears. Suspected *Cryptosporidium* and *Giardia* (oo)cysts were compared to stained
146 positive controls (A100FLR-1X Aqua-Glo G/C Direct positive control, Waterborne Inc. USA),
147 where MK and SMB stained *Cryptosporidium* (oo)cysts stood out as round bodies measuring 4 to
148 6 μ m in diameter with occasionally visible sporozoites that are bright red against a blue
149 background. Aura and IFT stained *Cryptosporidium* oocysts fluoresced bright apple-green against
150 a black background (CDC DPDx Cryptosporidiosis). *Giardia* cysts stained with IFT were ovoid,
151 measuring 10 to 14 μ m, and fluoresced bright apple-green against a black background (CDC DPDx
152 Giardiasis).

153

154 2.5. OM-SAB grown *in-situ*

155 Naturally grown *in situ* one-month-old substrate-associated biofilms (OM-SAB) were
156 produced from the Ipo Watershed using glass coverslip substrates that were secured to 20 X 3 cm
157 plastic panels held together by a 20 X 5 cm Styrofoam body in the horizontal and vertical

158 orientation on either side (Supplementary ~~1~~-material 1). The constructed substrates were immersed
159 at least 30 cm below the surface of the water from October 1, 2018 to November 1, 2018.

160

161 2.6. OM-SAB microscopy panel

162 To be able to determine the presence of naturally-associated *Cryptosporidium* and *Giardia*
163 (oo)cysts in OM-SAB, coverslip panels with OM-SAB were collected and washed with sterile
164 distilled water to remove non-adherent cells and individually placed into 50 mL sterile
165 polyethylene containers, transported to the laboratory, and harvested by scraping the coverslip
166 substrate-side (in contact with the water column) with sterile scalpel blades, and prepared into 2
167 mL microcentrifuge tube suspensions using sterile distilled water. OM-SAB suspensions were
168 vortexed for one minute and left to stand for five minutes to settle the heavier particles. 25 μ L of
169 the OM-SAB suspension was aspirated and prepared in duplicate into 1-cm diameter smears on
170 clean glass slides and stained with MK, SMB, Aura, and IFT (Aqua-Glo™ G/C Direct
171 Comprehensive Kit) according to the manufacturer's directions.

172

173 2.7. Scanning electron microscopy (SEM) of OM-SAB

174 To be able to provide SEM visualization of the natural architecture and composition of an
175 environmental aquatic substrate-associated biofilm, horizontal and vertical OM-SAB grown on
176 glass coverslips substrates were fixed with absolute methanol and allowed to dry for 24 hours. The
177 coverslip substrates were gently broken into shards of approximately 3 X 3 mm with care not to
178 disrupt the OM-SAB, then attached to a carbon tape secured on the SEM (TM3000 Hitachi
179 Tabletop SEM) metal platform for examination at the University of Santo Tomas, Thomas Aquinas
180 Research Centre.

181

182 2.8. Spiking of OM-SAB for Aura, IFT, and SEM

183 In order to simulate environmental contamination of OM-SAB with *Cryptosporidium*
184 oocysts, 25 μ L of OM-SAB suspensions were spiked with 10 μ L of *C. parvum* oocysts in 1×10^6
185 cells / mL concentration (P102C @ 1×10^6 , *Cryptosporidium parvum* oocysts, 1 million, in 4 mL,
186 Waterborne Inc. USA) and prepared in duplicate into 1 cm diameter smears on clean glass slides
187 and stained with Aura and IFT. For SEM analysis of spiked OM-SAB, one vertical and one
188 horizontal OM-SAB were fixed with absolute methanol, allowed to dry for 24 hours, then
189 manually broken into shards of approximately 3 X 3 mm with care not to disrupt the OM-SAB,
190 then spiked with 5 μ L of *C. parvum* oocysts, and allowed to dry for 24 hours for visualization
191 through SEM.

192

193 3. Results

194 Table 1 outlines the coordinates (geographical location), WBPP, and contributory
195 contamination sources for each sampling area. The majority of the samples tested positive for
196 waterborne protozoan pathogens in Thailand (3/3) and the Philippines (15/20) sampling areas.
197 Overall, 78% (18/23) of the aquatic sample matrices tested in this study contained at least one
198 WBPP in at least one sample matrix. Table 2 highlights the natural association of *Cryptosporidium*
199 and *Giardia* (oo)cysts in SAB relative to SW and BW matrices tested in this study. IFT confirmed
200 22% (5/23) *Cryptosporidium* and 4% (1/23) *Giardia* (oo)cysts in SAB which collates with 39%
201 positivity from both SW and BW counterparts. Table 3 shows a 33% (3/9) agreement between SW
202 and SAB samples positive for WBPP and the increased detection capacity of WBPP by 50% (7/14)
203 as compared to SW negative for WBPP. The comparison of BW and SAB demonstrated similar

204 results. Table 4 provides a panel of microscopy results for OM-SAB, where horizontal and vertical
205 samples were negative for naturally-associated WBPP. The absence of *Giardia* cysts was
206 consistent in both natural and spiked OM-SAB after staining with Aura and IFT, which confirms
207 its absence from the samples tested in the study. Fig. 2. demonstrates positive controls of
208 *Cryptosporidium* and *Giardia* (oo)cysts that were used as a reference for microscopic
209 identification. Fig. 3. shows the microscopic detection of *Cryptosporidium* and *Giardia* (oo)cysts
210 in SW, BW, and SAB using the elected staining techniques. Fig. 3.a-c demonstrates typical
211 *Cryptosporidium* oocysts stained with MK and SMB as round (oo)cysts with internal structures
212 that stained red by carbol fuchsin, while Fig. 3.d-f demonstrates apple-green fluorescence
213 attributed to Auramine and IFT stains. All *Cryptosporidium* oocysts (Fig. 3.a-e) fit within the 4 to
214 6 μm diameter range and had morphologic characteristics similar to the positive control of *C.*
215 *parvum* (as shown in Fig. 2.a-d). Fig 3.f demonstrates the oval to ellipsoid morphology and size
216 range of 10 to 14 μm of *Giardia* cysts, similar to the *G. lamblia* positive control found in Fig. 2.e.
217 Fig. 4. shows pictures of OM-SAB grown *in situ* in Ipo watershed on glass coverslip substrates
218 with microscopic architecture and associated structures visualized through SEM. Thick networks
219 of exopolysaccharide matrix ~~were~~ observed in Fig. 4.d which was responsible for binding the
220 contents of the biofilm microcosm. A naturally-associated testate amoeba shell (Fig. 4.e) was
221 incidentally identified in a vertical OM-SAB. Fig. 4.g demonstrates simulated environmental
222 contamination by spiking with *C. parvum* (oo)cysts for scenario purposes and the relative
223 comparison of the size of *C. parvum* oocysts against contents of the OM-SAB. Evidence of the
224 temporal accumulation and the diversity of organic materials in the biofilm matrix, mainly,
225 freshwater diatoms (Fig. 4.f), contributeds to the variable surface roughness and porosity which
226 can potentially facilitate attachment and trapping of pathogenic (oo)cysts and other organic

227 materials. Horizontal substrates and substrates immersed at ~~122 cm-2-meter~~ depth had denser
228 biofilm growth compared to vertical and 30 cm depth counterparts which were observed by
229 measuring the biofilm mass through light absorbance at optical density of 590 nm using a
230 spectrophotometer (Supplementary† material 2) while~~and~~ dry weight expressed in grams was
231 measured using an analytical balance (Supplementary material -3). All staining methods detected
232 0-1 (oo)cyst per 25 µL smeared suspension thereby providing an approximated maximum (oo)cyst
233 load of 2,000 (oo)cysts per 50 mL of sample matrix in the present study or 40,000 (oo)cysts per 1
234 L of the sample matrix. On very rare occasions, 0-2 (oo)cysts per 25 µL were observed.

235

236 **4. Discussion**

237 *4.1 Microscopy panel and low-volume sampling*

238 The results of this study show that traditional staining methods of MK and SMB are cost-
239 effective but provided lower detection compared to fluorescence techniques. The low-cost MK
240 staining method has been reported to exhibit 66% sensitivity and up to 88% specificity (Elsafi et
241 al., 2014; Johnston et al., 2003), and only requires a light microscope (Current and Garcia, 1991).
242 In the present study, only 4% (1/23) SAB tested positive for *Cryptosporidium* (oo)cysts using MK,
243 which collates with SW and BW results (Tables 1 and 2). In similar studies, MK detected only 2%
244 (3/135) *Cryptosporidium* (oo)cysts from water samples in Turkey, which were identified as *C.*
245 *parvum* after polymerase chain reaction (Aslan et al., 2012). In addition, MK presented with 71%
246 (53/75) positivity for *Cryptosporidium* oocysts in a metropolitan watershed in the Philippines,
247 which suggests heavy parasite contamination during the three-day sampling period (Masangkay et
248 al., 2016). In the present study, the staining characteristics of *Cryptosporidium* were identical for
249 both MK and SMB, where 7% (2/23) positivity for SMB was relatively close to the results obtained

250 from MK but lower than Aura with a 39% (9/23) screening positivity. These results suggest
251 variability in the power of (oo)cysts detection depending on the microscopic method and the utility
252 of Aura as a convenient screening method for detection of *Cryptosporidium* oocysts in water
253 samples (Ahmed and Karanis, 2018; Hanscheid et al., 2008; Smith et al., 1989). In the present
254 study, IFT findings were positive for *Cryptosporidium* and *Giardia* (oo)cysts at 22% and 4%,
255 respectively but IFT kits were expensive and required a fluorescence microscope. However, the
256 cost may be justified by the high degree of sensitivity (99%) and specificity (100%) for
257 *Cryptosporidium* and 96% to 100%, respectively, for *Giardia* (Adeyemo et al., 2018; Pacheco et
258 al., 2013). Researchers in different parts of the globe have extensively contributed to establishing
259 the importance of water analysis methodologies in the effective detection of WBPP (Estratiou et
260 al., 2017a; 2017b; Plutzer and Karanis, 2016). Likewise, the establishment of cost-effective
261 concentration methods for WBPP in water samples like flotation and flocculation being applied to
262 large and lower volume water samples have gained traction over the past two decades (Gallas-
263 Lindemann et al., 2013; 2016; Karanis et al., 2006; Karanis and Kimura, 2002; Koloren et al.,
264 2016; 2018; Kourenti et al., 2003; Kourenti and Karanis, 2004; 2006; Ma et al., 2019; Tsushima
265 et al., 2001; 2003a; 2003b). Low-volume water sampling for the detection of *Cryptosporidium* and
266 *Giardia* is not routinely performed but a number of studies have already confirmed its benefits as
267 reported in the PCR positivity of *Cryptosporidium* in 50 mL raw river water samples in China
268 (Xiao et al., 2012), and a study in the Philippines that reported both MK and PCR confirmation
269 and sequence identification of *C. hominis* directly from 50 mL samples (Masangkay et al., 2016;
270 2019). These methods are cost-effective alternatives to Method 1623 or other methodologies,
271 providing effective detection of *Cryptosporidium* and *Giardia*, particularly in high-turbidity
272 aquatic samples (Bilung et al., 2017; Efstratiou et al., 2017b). In the present study, low-volume

273 water sampling offered the advantages of ease of collection and transport, multiple matrix
274 sampling (SW, BW, and SAB), multiple area sampling per study site, reproducibility, and
275 significantly lower test cost. The (oo)cyst load of approximately 2,000 (oo)cysts per 50 mL or
276 40,000 (oo)cysts per 1 L sample matrix indicated high contamination of the sampling areas with
277 WBPP. These estimates, however, should not be taken as absolute counts as (oo)cysts can be
278 unevenly dispersed and associated with bio-colloids, thereby further complicating its' non-
279 homogeneous distribution in environmental aquatic matrices. The variation of positive and
280 negative results per staining method across each 50 mL sample may have been influenced by the
281 non-homogeneous distribution of (oo)cysts in the aquatic sample matrices.

282

283 4.2 Spatial distribution of WBPP in aquatic matrices

284 Spatial distribution of *Cryptosporidium* and *Giardia* in sample matrices as shown in Table
285 1 and Table 2 demonstrate that BW and SW, and in particular SAB are all suitable environmental
286 aquatic matrices for the detection of WBPP. Out of the 23 combined sampling areas (3 in Thailand
287 and 20 in the Philippines), 78% (18/23) were positive for *Cryptosporidium* and/or *Giardia* in at
288 least one sample matrix, where 39% (9/23) of both SW and BW, and 43% (10/23) SAB, tested
289 positive. Results of this study on the presence of *Cryptosporidium* and/or *Giardia* in environmental
290 drinking and recreational waters can be supported by the investigations done in 2016 in the La
291 Mesa Watershed in the Philippines, where a high incidence of *Cryptosporidium* and *Cyclospora*
292 was documented over a three-day period (Masangkay et al., 2016). Similarly, a study in Malaysia
293 reported 100% (24/24) of SW to be positive for *Cryptosporidium* and *Cyclospora* (Bilung et al.,
294 2017), with *Giardia* cysts reported to contaminate SW as well (Lass et al., 2017; Ramo et al.,
295 2017). Studies investigating pathogenic protozoans from BW are rare; one exemplary study for

296 BW was reported in 2017 from the Yunlong Lake in China where 47% (28/60) BW tested positive
297 for *Cryptosporidium* (Kong et al., 2017). Although *in vitro* extracellular excystation of
298 *Cryptosporidium* has been elaborated in *Pseudomonas aeruginosa* aquatic biofilms (Koh et al.,
299 2013; 2014), there has been no documented case of the natural-association of *Cryptosporidium* in
300 environmental aquatic biofilms until the first report of *C. hominis* in SAB that was isolated from
301 a freshwater sponge in the Philippines in 2019 (Masangkay et al., 2019). In the present study, the
302 value of SAB as a supplemental sample matrix provided close agreement with both SW and BW
303 positivity and increased reporting of WBPP as compared to analyzing SW alone (Table 1 and
304 Table 3). The possible contributory factors for contamination of the sampling areas and the cycling
305 of WBPP between humans and the environment can come from many sources (Table 1) including
306 anthropogenic activities, communities with poor sanitary and living conditions, and improper
307 domestic wastewater sanitization procedures (Adamska, 2014; Bhattachan et al., 2017; Masangkay
308 et al., 2016). The presence of wildlife, domestic, and farm animals in the surroundings of lakes
309 and other water reservoirs plays a significant role in parasite transmission to other animals and
310 water sources thereby contributing to zoonotic transmission of *Cryptosporidium* and *Giardia* (Gil
311 et al., 2017; Wells et al., 2019; Zahedi et al., 2016). In addition, birds, fish, amphibians, and small
312 mammals have been tested and reported to be positive for *Cryptosporidium* and *Giardia* as well
313 (Hublova et al., 2016; Karanis et al., 1996; Ryan, 2010; Yang et al., 2015). Soil run-off, mixed
314 with animal and human excreta can contribute to source water contamination with WBPP (Dai and
315 Boll, 2003; Norman et al., 2013). Aquatic plants and immersed substrates, as demonstrated in this
316 study, can harbor pathogenic (oo)cysts through the temporal accumulation of (oo)cysts in SAB.
317 These contaminating factors are best exemplified in the University Lake of Thailand, which
318 simulates anthropogenic, zoogenic, and environmental cycling of WBPP in a small water

319 catchment (Fig. 3.a-b and Table 1), where all sample matrices were positive for *Cryptosporidium*
320 oocysts and TS1A3 SW was simultaneously positive for both *Cryptosporidium* and *Giardia*.

321

322 4.2. Aquatic biofilms: implications for water quality assessment strategies (WQAS)

323 Biofilms in relation to WQAS have been limited to tap water systems (Wingender and
324 Flemming, 2011; van der Kooij et al., 2017; Zhou et al., 2017), while biofilms in natural freshwater
325 resources are not screened for *Cryptosporidium* and *Giardia* contamination. The natural-
326 association of a testate amoeba shell suspected of being *Diffugia* (Qin et al., 2011) on the OM-
327 SAB surface (Fig. 4.e) demonstrates the potential for interactions between SAB and bio-colloids
328 like *Cryptosporidium* and *Giardia* (oo)cysts (Luo et al., 2016; Searcy et al., 2006). Parasite load
329 and temporal accumulation contribute to the presence of WBPP in SAB as can be observed from
330 Ipo watershed where OM-SAB grown *in-situ* were negative for *Cryptosporidium* and *Giardia*
331 (Table 4) seemingly because of low parasite load where only one *Cryptosporidium* oocyst was
332 detected from all SW and BW from all four sampling sites (results not shown). Fig. 4.c and 4.d
333 demonstrate biofilm roughness and porosity and a network of the exopolysaccharide matrix which
334 can trap *Cryptosporidium* and *Giardia* (oo)cysts (Wolyniak et al., 2010). This is in agreement with
335 the observations presented in Fig. 4.g, where *C. parvum* oocysts spiking experiment revealed the
336 relative size comparison between *Cryptosporidium* oocysts and the size of channels and spaces on
337 the OM-SAB surface, which can potentially permit the temporal accumulation and trapping of
338 WBPP, including larger testate amoeba shells among other bio-colloids. Further, the abundance of
339 organic debris in the OM-SAB permits UV radiation protection, which further contributes to the
340 maintenance of (oo)cyst viability (DiCesare et al., 2012b). As shown in Table 3, additional analysis
341 of SAB generated 33% agreement with positive SW results and enhanced detection of WBPP to

342 50% by not declaring the water samples as not contaminated based on negative SW results alone
343 with similar results observed from SAB and BW comparisons. As opposed to the real-time nature
344 of contamination from SW and BW matrices, SAB is a matrix, which submits results based on the
345 temporal accumulation of WBPP similar to reports of temporal accumulation of *Cryptosporidium*
346 oocysts in marine shellfish (Pagoso and Rivera, 2017) and freshwater sponge in the Philippines
347 (Masangkay et al., 2019). The results of this study support the hypothesis of the natural-association
348 of WBPP in SAB that can be exploited to detect *Cryptosporidium* and *Giardia* (oo)cysts from
349 environmental freshwater resources and lead to improved water quality assessment strategies
350 (iWQAS). The formulation of screening initiatives for the simultaneous testing of SW, BW, and
351 SAB in multiple sampling areas per study site can also provide important data for limiting the
352 exposure of humans and animals to WBPP, stimulate government responses, regulatory actions,
353 and improved accessibility to screening protocols.

354

355 **5. Conclusions**

356 Screening biofilms for waterborne protozoan pathogens has been an underappreciated tool
357 in the detection and mitigation of waterborne infections and outbreaks. The results of this study
358 provide evidence that analysis of aquatic substrate-associated biofilms leads to improved water
359 quality assessment strategies. Employing more than one microscopy method for the detection of
360 waterborne protozoan pathogens in low-volume samples can improve *Cryptosporidium* and
361 *Giardia* detection from surface water, bottom water with sediments, and substrate-associated
362 biofilms. Biofilms can act as biological reservoirs for waterborne protozoan pathogens by
363 associating and protecting (oo)cysts from UV exposure within their matrices and surfaces.

364

365 **Funding**

366 This study was financially supported by The Walailak University Graduate Studies
367 Research Fund Contract No. 24/2562 awarded to FRM.

368

369 **Acknowledgments**

370 We extend our deepest gratitude to Antoinette E. Dasig, Michael Christopher P. Agustin,
371 Rado Paul F. Cruz, Harrison Karl C. De Jesus, Jasmin Aliya L. Ellazar, Enrico B. Gozum, and
372 Paula Marie Isabel F. Inocencio of the Department of Medical Technology, College of Pharmacy,
373 The Pontifical and Royal University of Santo Tomas for being exemplary in their work and as
374 undergraduate research students whose roles played an integral part in the fruition of this
375 international collaboration. We thank Joseph Dionisio, Romel Solomon, Virgilio Bitagcul, Noel
376 Cajuday, Maria Lourdes Policarpio, Ivy Timogtimog, and Ernest Hidalgo of Far Eastern
377 University-Manila for laboratory logistics; Nina Caisido, June Linsangan of the UST Analytical
378 Services Laboratory Research Center for the Natural and Applied Sciences, and Director of facility
379 Dr. Christina Binag for SEM visualizations, Harry Esmaya of Howell Medical supplies for the
380 fluorescence microscopy facility, and Dr. Henry H. Stibbs, Angela Faust, and Molly Rowe of
381 Waterborne Inc. for the comprehensive IFT kit. We thank Chad Schou for the editorial
382 proofreading of the manuscript.

383

384 **Conflict of interests**

385 The authors declare no conflict of interest.

386

387 **References**

388 Adamska, M., 2014. Molecular characterization of *Cryptosporidium* and *Giardia* occurring in
389 natural water bodies in Poland. *Parasitol Res.* 114(2):687–92. doi: 10.1007/s00436-014-
390 4234-9.

391 Adeyemo, F.E., Singh, G., Reddy, P., Stenström, T.A., 2018. Methods for the detection of
392 *Cryptosporidium* and *Giardia*: From microscopy to nucleic acid based tools in clinical and
393 environmental regimes. *Acta Trop.* 184:15-28. doi: 10.1016/j.actatropica.2018.01.011.

394 Ahmed, S.A., Karanis, P., 2018. An overview of methods/techniques for the detection of
395 *Cryptosporidium* in food samples. *Parasitol Res.* 117:629–653. doi.org/10.1007/s00436-
396 017-5735-0.

397 Aslan, G., Bayram, G., Otağ, F., Direkel, S., Taylan Özkan, A., Ceber, K., Emekdaş, G., 2012.
398 Investigation of the presence of *Cryptosporidium* spp. in different water sources in Mersin
399 province, Turkey. *Mikrobiyol Bul.* 46(1):93-100.

400 Baldursson, S., Karanis, P., 2011. Waterborne transmission of protozoan parasites: review of
401 worldwide outbreaks - an update 2004-2010. *Wat Res.* 15;45(20):6603-14. doi:
402 10.1016/j.watres.2011.10.013.

403 Bhattachan, B., Sherchand, J.B., Tandukar, S., Dhoubhadel, B.G., Gauchan, L., Rai, G., 2017.
404 Detection of *Cryptosporidium parvum* and *Cyclospora cayetanensis* infections among
405 people living in a slum area in Kathmandu valley, Nepal. *BMC Res Notes.* 10(1):1–5. doi:
406 10.1186/s13104-017-2779-2.

407 Bekliz, M., Colson, P., La Scola, B., 2016. The expanding family of virophages. *Viruses.* 8(11).
408 pii: E317. doi: 10.3390/v8110317.

409 Bilung, L.M., Tahar, A.S., Yunos, N.E., Apun, K., Lim, Y.A., Nillian, E., Hashim, H.F., 2017.
410 Detection of *Cryptosporidium* and *Cyclospora* oocysts from environmental water for

411 drinking and recreational activities in Sarawak, Malaysia. Biomed Res Int. 2017:4636420.
412 doi: 10.1155/2017/4636420.

413 Current, W.L., Garcia, L.S., 1991. Cryptosporidiosis. Clin Microbiol. 4: 325e358.

414 Checkley, W., White, A.C. Jr., Jaganath, D., Arrowood, M.J., Chalmers, R.M., Chen, X.M., Fayer,
415 R., Griffiths, J.K., Guerrant, R.L., Hedstrom, L., Huston, C.D., Kotloff, K.L., Kang, G.,
416 Mead, J.R., Miller, M., Petri, W.A. Jr., Priest, J.W., Roos, D.S., Striepen, B., Thompson,
417 R.C., Ward, H.D., Van Voorhis, W.A., Xiao, L., Zhu, G., Houpt, E.R., 2015. A review of
418 the global burden, novel diagnostics, therapeutics, and vaccine targets for
419 *Cryptosporidium*. Lancet Infect Dis. 15(1):85–94. doi: 10.1016/S1473-3099(14)70772-8.

420 Dai, X., Boll, J., 2003. Evaluation of attachment of *Cryptosporidium parvum* and *Giardia lamblia*
421 to soil particle. J Environ Qual. 32(1):296–304. doi: 10.2134/jeq2003.2960.

422 DiCesare, E.A., Hargreaves, B.R., Jellison, K.L., 2012a. Biofilm roughness determines
423 *Cryptosporidium parvum* retention in environmental biofilms. Appl Environ Microbiol.
424 78(12):4187–93. doi: 10.1128/AEM.08026-11.

425 DiCesare, E.A., Hargreaves, B.R., Jellison, K.L., 2012b. Biofilms reduce solar disinfection of
426 *Cryptosporidium parvum* oocysts. Appl Environ Microbiol. 78(12):4522–5. doi:
427 10.1128/AEM.08019-11.

428 DuPont, H.L., Chappell, C.L., Sterling, C.R., Okhuysen, P.C., Rose, J.B., Jakubowski, W., 1995.
429 The infectivity of *Cryptosporidium parvum* in healthy volunteers. N Engl J Med.
430 332(13):855-9. doi: 10.1056/NEJM199503303321304.

431 Efstratiou, A., Ongerth, J.E., Karanis, P., 2017a. Waterborne transmission of protozoan parasites:
432 Review of worldwide outbreaks - An update 2011-2016. Wat Res. 114:14-22. doi:
433 10.1016/j.watres.2017.01.036.

434 Efstratiou, A., Ongerth, J., Karanis, P., 2017b. Evolution of monitoring for *Giardia* and
435 *Cryptosporidium* in water. *Wat Res.* 123:96-112. doi: 10.1016/j.watres.2017.06.042.

436 Elsafi, S.H., Al-Sheban, S.S., Al-Jubran, K.M., Abu Hassan, M.M., Al Zahrani, E.M., 2014.
437 Comparison of Kinyoun's acid-fast and immunofluorescent methods detected an
438 unprecedented occurrence of *Cryptosporidium* in the Eastern Region of Saudi Arabia. *J*
439 *Taibah Univ Med Sci.* 9(4): 263e267.

440 Gallas-Lindemann, C., Sotiriadou, I., Plutzer, J., Karanis, P., 2013. Prevalence and distribution of
441 *Cryptosporidium* and *Giardia* in wastewater and the surface, drinking and ground waters
442 in the Lower Rhine, Germany. *Epidemiol Infect.* 141(1):9-21. doi:
443 10.1017/S0950268812002026.

444 Gallas-Lindemann, C., Sotiriadou, I., Plutzer, J., Noack, M.J., Mahmoudi, M.R., Karanis, P., 2016.
445 *Giardia* and *Cryptosporidium* spp. dissemination during wastewater treatment and
446 comparative detection via immunofluorescence assay (IFA), nested polymerase chain
447 reaction (nested PCR) and loop mediated isothermal amplification (LAMP). *Acta Trop.*
448 158:43-51. doi: 10.1016/j.actatropica.2016.02.005.

449 Gil, H., Cano, L., de Lucio, A., Bailo, B., de Mingo, M.H., Cardona, G.A., Fernández-Basterra,
450 J.A., Aramburu-Aguirre, J., López-Molina, N., Carmena, D., 2017. Detection and
451 molecular diversity of *Giardia duodenalis* and *Cryptosporidium* spp. in sheltered dogs and
452 cats in Northern Spain. *Infect Genet Evol.* 50:62–9. doi: 10.1016/j.meegid.2017.02.013.

453 Guzman, M.E.J., 2006. A framework for the sustainability of Laguna de Bay, Philippines. *Wat*
454 *Environ Found.* 1449–59.

455 Hanscheid, T., Cristino, J.M., Salgado, M.J., 2008. Screening of auramine stained smears of all
456 fecal samples is a rapid and inexpensive way to increase the detection of coccidial
457 infections. *Int J Infect Dis* 12(1): 47–50. <https://doi.org/10.1016/j.ijid.2007.04.008>.

458 Holubová, N., Sak, B., Horčíčková, M., Hlásková, L., Květoňová, D., Menchaca, S., McEvoy, J.,
459 Kváč, M., 2016. *Cryptosporidium avium* n. sp. (Apicomplexa: Cryptosporidiidae) in birds.
460 *Parasitol Res.* 115(6):2243–51. doi: 10.1007/s00436-016-4967-8.
461 <https://www.cdc.gov/dpdx/cryptosporidiosis/index.html>.
462 <https://www.cdc.gov/dpdx/giardiasis/index.html>.

463 Israel, D.C., 2007. The current state of aquaculture in Laguna de Bay. Discussion paper series no.
464 2007-20, Philippine Institute for Development Studies. 1-63.

465 Johnston, S.P., Ballard, M.M., Beach, M.J., Causer, L., Wilkins, P.P., 2003. Evaluation of three
466 commercial assays for detection of *Giardia* and *Cryptosporidium* organisms in fecal
467 specimens. *J Clin Microbiol.* 41: 623e626. doi: 10.1128/jcm.41.2.623-626.2003.

468 Karanis, P., Kimura, A., 2002. Evaluation of three flocculation methods for the purification of
469 *Cryptosporidium parvum* oocysts from water samples. *Lett Appl Microbiol.*
470 2002;34(6):444-9. doi: 10.1046/j.1472-765x.2002.01121.x.

471 Karanis, P., Kourenti, C., Smith, H., 2007. Waterborne transmission of protozoan parasites: a
472 worldwide review of outbreaks and lessons learnt. *J Wat Health.* 5(1):1-38. doi:
473 10.2166/wh.2006.002.

474 Karanis, P., Opiela, K., Renoth, S., Seitz, H.M., 1996. Possible contamination of surface waters
475 with *Giardia* spp. through muskrats. *Zentralbl Bakteriol.* 284(2-3):302-6. doi:
476 10.1016/s0934-8840(96)80106-x.

477 Karanis, P., Sotiriadou, I., Kartashev, V., Kourenti, C., Tsvetkova, N., Stojanova, K., 2006.
478 Occurrence of *Giardia* and *Cryptosporidium* in water supplies of Russia and Bulgaria.
479 Environ Res. 102(3):260-71. doi: 10.1016/j.envres.2006.05.005.

480 Koh, W., Clode, P.L., Monis, P., Thompson, R.C., 2013. Multiplication of the waterborne
481 pathogen *Cryptosporidium parvum* in an aquatic biofilm system. Parasit Vectors. 6:270.
482 doi: 10.1186/1756-3305-6-270.

483 Koh, W., Thompson, A., Edwards, H., Monis, P., Clode, P.L., 2014. Extracellular excystation and
484 development of *Cryptosporidium*: tracing the fate of oocysts within *Pseudomonas* aquatic
485 biofilm systems. BMC Microbiol. 14:281. doi: 10.1186/s12866-014-0281-8.

486 Koloren, Z., Seferoğlu, O., Karanis, P., 2016. Occurrence of *Giardia duodenalis* assemblages in
487 river water sources of Black Sea, Turkey. Acta Trop. 164:337-344. doi:
488 10.1016/j.actatropica.2016.09.025.

489 Koloren, Z., Gulabi, B.B., Karanis, P., 2018. Molecular identification of *Blastocystis* sp. subtypes
490 in water samples collected from Black sea, Turkey. Acta Trop. 180:58-68. doi:
491 10.1016/j.actatropica.2017.12.029.

492 Kong, Y., Lu, P., Yuan, T., Niu, J., Li, Z., Yang, B., 2017. *Cryptosporidium* contamination and
493 attributed risks in Yunlong Lake in Xuzhou, China. Can J Infect Dis Med Microbiol.
494 2017:4819594. doi: 10.1155/2017/4819594.

495 Kourenti, C., Heckerroth, A., Tenter, A., Karanis, P., 2003. Development and application of
496 different methods for the detection of *Toxoplasma gondii* in water. Appl Environ
497 Microbiol. 69(1):102-6. doi: 10.1128/aem.69.1.102-106.2003

498 Kourenti, C., Karanis, P., 2004. Development of a sensitive polymerase chain reaction method for
499 the detection of *Toxoplasma gondii* in water. Water Sci Technol. 2004;50(1):287-91.

500 Kourenti, C., Karanis, P., 2006. Evaluation and applicability of a purification method coupled with
501 nested PCR for the detection of *Toxoplasma* oocysts in water. Lett Appl Microbiol.
502 43(5):475-81. doi: 10.1111/j.1472-765X.2006.02008.x.

503 Lass, A., Szostakowska, B., Korzeniewski, K., Karanis, P., 2017. Detection of *Giardia intestinalis*
504 in water samples collected from natural water reservoirs and wells in Northern and North-
505 Eastern Poland using LAMP, real-time PCR and nested PCR. J Wat Health. 15(5):775-
506 787. doi: 10.2166/wh.2017.039.

507 Luo, X., Jedlicka, S., Jellison, K., 2016. Pseudo-second-order calcium-mediated *Cryptosporidium*
508 *parvum* oocyst attachment to environmental biofilms. Appl Environ Microbiol. 83(1). pii:
509 e02339-16. doi: 10.1128/AEM.02339-16.

510 Ma, L., Zhang, X., Jian, Y., Li, X., Wang, G., Hu, Y., Karanis, P., 2019. Detection of
511 *Cryptosporidium* and *Giardia* in the slaughterhouse, sewage and river waters of the
512 Qinghai Tibetan plateau area (QTPA), China. Parasitol Res. 118(7):2041-2051. doi:
513 10.1007/s00436-019-06330-w.

514 Mac Kenzie, W.R., Hoxie, N.J., Proctor, M.E., Gradus, M.S., Blair, K.A., Peterson, D.E.,
515 Kazmierczak, J.J., Addiss, D.G., Fox, K.R., Rose, J.B., et al., 1994. A massive outbreak in
516 Milwaukee of *Cryptosporidium* infection transmitted through the public water supply. N
517 Engl J Med. 331(3):161–7. doi: 10.1056/NEJM199407213310304.

518 Martinez, F.G.I., 2011. Monitoring and evaluation of the water quality of Taal Lake, Talisay,
519 Batangas, Philippines. Acad Res Int. 1(1):229–36.

520 Masangkay, F., Milanez, G., Chua, N., Angulo, F., Aquino, P., Calucin, D., Urtal, G., 2016. Water-
521 borne coccidians in Philippine water sheds: A national inceptive study. Asian J Biol Life
522 Sci. 5(2):149–51.

523 Masangkay, F., Milanez, G., Karanis, P., Nissapatorn, V., 2018. *Vermamoeba vermiformis* —
524 Global trend and future perspective. 2nd ed. Reference Module in Earth Systems and
525 Environmental Sciences. Elsevier Inc. 1-11.

526 Masangkay, F.R., Milanez, G.D., Tsiami, A., Somsak, V., Kotepui, M., Tangpong, J., Karanis, P.,
527 2019. First report of *Cryptosporidium hominis* in a freshwater sponge. Sci Total Environ.
528 700:134447. doi: 10.1016/j.scitotenv.2019.134447.

529 Matos, A.O., Ricomini-Filhom A.P., Beline, T., Ogawa, E.S., Costa-Oliveira, B.E., de Almeida,
530 A.B., Nociti Junior, F.H., Rangel, E.C., da Cruz, N.C., Sukotjo, C., Mathew, M.T., Barão,
531 V.A.R., 2017. Three-species biofilm model onto plasma-treated titanium implant surface.
532 Colloids Surf B Biointerfaces. 152:354-366. doi: 10.1016/j.colsurfb.2017.01.035.

533 Murphy, J.L., Hlavsa, M.C., Carter, B.C., Miller, C., Jothikumar, N., Gerth, T.R., Beach, M.J.,
534 Hill, V.R., 2018. Pool water quality and prevalence of microbes in filter backwash from
535 metro-Atlanta swimming pools. J Wat Health. 16(1):93–101. doi: 10.2166/wh.2017.150.

536 Norman, S.A., Hobbs, R.C., Wuertz, S., Melli, A., Beckett, L.A., Chouicha, N., Kundu, A., Miller,
537 W.A., 2013. Fecal pathogen pollution: Sources and patterns in water and sediment samples
538 from the upper Cook inlet, Alaska ecosystem. Environ Sci Process Impacts. 15(5):1041–
539 51. doi: 10.1039/c3em30930d.

540 Okhuysen, P.C., Chappell, C.L., Crabb, J.H., Sterling, C.R., DuPont, H.L., 1999. Virulence of
541 three distinct *Cryptosporidium parvum* isolates for healthy adults. J Infect Dis.
542 180(4):1275-81. doi: 10.1086/315033.

543 Ortega, Y.R., Adam, R.D., 1997. *Giardia*: overview and update. Clin Infect Dis. 25(3):545-9. doi:
544 10.1086/513745.

545 P102X@1x10/6 Product Description: *Cryptosporidium parvum* oocysts, 1 million, in 4 mL.
546 <http://waterborneinc.com/p102-oocysts-ofcryptosporidium-parvum>.

547 Pacheco, F.T., Silva, R.K., Martins, A.S., Oliveira, R.R., Alcântara-Neves, N.M., Silva, M.P.,
548 Soares, N.M., Teixeira, M.C., 2013. Differences in the detection of *Cryptosporidium* and
549 *Isospora* (*Cystoisospora*) oocysts according to the fecal concentration or staining method
550 used in a clinical laboratory. J Parasitol. 1002-8. doi: 10.1645/12-33.1.

551 Pagoso, E.J.A., Rivera, W.L., 2017. *Cryptosporidium* species from common edible bivalves in
552 Manila Bay, Philippines. Mar Pollut Bull. 119(1):31-39. doi:
553 10.1016/j.marpolbul.2017.03.005.

554 Plutzer, J., Karanis, P., 2016. Neglected waterborne parasitic protozoa and their detection in water.
555 Water Res. 2016 Sep 15;101:318-332. doi: 10.1016/j.watres.2016.05.085.

556 Qin, Y., Xie, S., Smith, H., Swindles, G.T., Gu, Y., 2011. Diversity, distribution and biogeography
557 of testate amoeba in China: Implications for ecological studies in Asia. Eur J Protistol.
558 47(1):1-9. doi: 10.1016/j.ejop.2010.09.004.

559 Ramo, A., Del Cacho, E., Sánchez-Acedo, C., Quílez, J., 2017. Occurrence of *Cryptosporidium*
560 and *Giardia* in raw and finished drinking water in north-eastern Spain. Sci Total Environ.
561 580:1007-1013. doi: 10.1016/j.scitotenv.2016.12.055.

562 Ryan, U., 2010. *Cryptosporidium* in birds, fish and amphibians. Exp Parasitol. 124(1):113–20. doi:
563 10.1016/j.exppara.2009.02.002.

564 Ryan, U., Paparini, A., Monis, P., Hijjawi, N., 2016. It's official – *Cryptosporidium* is a gregarine:
565 What are the implications for the water industry? Wat Res. 105:305–13. doi:
566 10.1016/j.watres.2016.09.013.

567 Savioli, L., Smith, H., Thompson, A., 2006. *Giardia* and *Cryptosporidium* join the ‘neglected
568 diseases initiative’. Trends Parasitol. 22:203-208. doi: 10.1016/j.pt.2006.02.015.

569 Searcy, K.E., Packman, A.I., Atwill, E.R., Harter, T., 2006. Deposition of *Cryptosporidium*
570 oocysts in streambeds. Appl Environ Microbiol. 72(3):1810-6. doi:
571 10.1128/AEM.72.3.1810-1816.2006.

572 Smith, H.V., McDiarmid, A., Smith, A.L., Hinson, A.R., Gilmour, R.A., 1989. An analysis of
573 staining methods for the detection of *Cryptosporidium* spp. oocysts in water-related
574 samples. Parasitol. 99 Pt 3:323-7. doi: 10.1017/s0031182000059023.

575 Steiner, T.S., Thielman, N.M., Guerrant, R.L., 1997. Protozoal agents: what are the dangers for
576 the public water supply? Annu Rev Med. 48:329-40. doi: 10.1146/annurev.med.48.1.329.

577 Suba, S.C., Masangkay, F.R., 2013. Transmission electron microscopy of in vitro biofilms formed
578 by *Candida albicans* and *Escherichia coli*. Asian J Biol Life Sci. 2(1):27–32.

579 Tsushima, Y., Karanis, P., Kamada, T., Nagasawa, H., Xuan, X., Igarashi, I., Fujisaki, K.,
580 Takahashi, E., Mikami, T., 2001. Detection of *Cryptosporidium parvum* oocysts in
581 environmental water in Hokkaido, Japan. J Vet Med Sci. 63(3):233-6. doi:
582 10.1292/jvms.63.233.

583 Tsushima, Y., Karanis, P., Kamada, T., Xuan, X., Makala, L.H., Tohya, Y., Akashi, H., Nagasawa,
584 H., 2003a. Viability and infectivity of *Cryptosporidium parvum* oocysts detected in river
585 water in Hokkaido, Japan. J Vet Med Sci. 65(5):585-9. doi: 10.1292/jvms.65.585.

586 Tsushima, Y., Karanis, P., Kamada, T., Makala, L., Xuan, X., Tohya, Y., Akashi, H., Nagasawa,
587 H., 2003b. Seasonal change in the number of *Cryptosporidium parvum* oocysts in water
588 samples from the rivers in Hokkaido, Japan, detected by the ferric sulfate flocculation
589 method. J Vet Med Sci. 65(1):121-3. doi: 10.1292/jvms.65.121.

590 US EPA, 2005. Method 1623: *Cryptosporidium* and *Giardia* in water by filtration/IMS/FA (PDF),
591 December 2005 Update (EPA 821-R-05-002). Office of Water 4603. U.S. Environmental
592 Protection Agency, Washington. <http://www.epa.gov/microbes/1623de05.pdf>.

593 van der Kooij, D., Bakker, G.L., Italiaander, R., Veenendaal, H.R., Wullings, B.A., 2017. Biofilm
594 composition and threshold concentration for growth of *Legionella pneumophila* on
595 surfaces exposed to flowing warm tap water without disinfectant. *Appl Environ Microbiol.*
596 83(5). doi: 10.1128/AEM.02737-16.

597 Waiser, M.J., Swerhone, G.D., Roy, J., Tumber, V., Lawrence, J.R., 2016. Effects of erythromycin,
598 trimethoprim and clindamycin on attached microbial communities from an effluent
599 dominated prairie stream. *Ecotoxicol Environ Saf.* 132:31–9. doi:
600 10.1016/j.ecoenv.2016.05.026.

601 Wells, B., Paton, C., Bacchetti, R., Shaw, H., Stewart, W., Plowman, J., Katzer, F., Innes, E.A.,
602 2019. *Cryptosporidium* prevalence in calves and geese co-grazing on four livestock farms
603 surrounding two reservoirs supplying public water to mainland Orkney, Scotland.
604 *Microorganisms.* 7(11). pii: E513. doi: 10.3390/microorganisms7110513.

605 Wingender, J., Flemming, H.C., 2011. Biofilms in drinking water and their role as reservoir for
606 pathogens. *Int J Hyg Environ Health.* (6):417-23. doi: 10.1016/j.ijheh.2011.05.009.

607 Wolyniak, E.A., Hargreaves, B.R., Jellison, K.L., 2009. Retention and release of *Cryptosporidium*
608 *parvum* oocysts by experimental biofilms composed of a natural stream microbial
609 community. *Appl Environ Microbiol.* 75(13):4624-6. doi: 10.1128/AEM.02916-08.

610 Wolyniak, E.A., Hargreaves, B.R., Jellison, K.L., 2010. Seasonal retention and release of
611 *Cryptosporidium parvum* oocysts by environmental biofilms in the laboratory. *Appl*
612 *Environ Microbiol.* 76(4):1021–7. doi: 10.1128/AEM.01804-09.

613 Xiao, S., An, W., Chen, Z., Zhang, D., Yu, J., Yang, M., 2012. Occurrences and genotypes of
614 *Cryptosporidium* oocysts in river network of Southern-Eastern China. Parasitol Res.
615 110(5):1701-9. doi: 10.1007/s00436-011-2688-6.

616 Xu, H., Zhang, W., Jiang, Y., Yang, E.J., 2014. Use of biofilm-dwelling ciliate communities to
617 determine environmental quality status of coastal waters. Sci Total Environ. 470–471:511–
618 8. doi: 10.1016/j.scitotenv.2013.10.025.

619 Yang, R., Palermo, C., Chen, L., Edwards, A., Paparini, A., Tong, K., Gibson-Kueh, S., Lymbery,
620 A., Ryan, U., 2015. Genetic diversity of *Cryptosporidium* in fish at the 18S and actin loci
621 and high levels of mixed infections. Vet Parasitol. 214(3–4):255–63. doi:
622 10.1016/j.vetpar.2015.10.013.

623 Zahedi, A., Paparini, A., Jian, F., Robertson, I., Ryan, U., 2016. Public health significance of
624 zoonotic *Cryptosporidium* species in wildlife: Critical insights into better drinking water
625 management. Int J Parasitol Parasites Wildl. 5(1):88-109. doi:
626 10.1016/j.ijppaw.2015.12.001.

627 Zhou, X., Zhang, K., Zhang, T., Li, C., Mao, X., Zhou, X., 2017. An ignored and potential source
628 of taste and odor (T & O) issues - biofilms in drinking water distribution system (DWDS).
629 Appl Microbiol Biotechnol. 101(9):3537–50. doi: 10.1007/s00253-017-8223-7.

HIGHLIGHTS

- Analysis of aquatic biofilms increased the detection of protozoan (oo)cysts.
- Biofilms acted as biological reservoirs for *Cryptosporidium* and *Giardia* (oo)cysts.
- *Cryptosporidium* and *Giardia* (oo)cysts were detected in low-volume samples of 50 mL.
- 50 mL sampling permits multiple sampling, reproducibility, and lower test cost.
- 50 mL is the lowest water volume reported for the detection of *Cryptosporidium*.

Graphical abstract

