1 **The mechanism for improving the flesh quality of grass carp (*Ctenopharyngodon idella*)**
2 **following the micro-flowing water treatment using a UPLC-QTOF/MS based metabolomics**
3 **method**

4

5 Hongying Du^{a, b}; Hao Lv^a; Zeru Xu^a; Siming Zhao^{a, b}; Tianwen Huang^d; Anne Manyande^c; Shanbai

6 Xiong^{a, b, *}

7

8 ^a Key Laboratory of Environment Correlative Dietology, Ministry of Education, College of Food Science
9 and Technology, Huazhong Agricultural University, Wuhan, Hubei, P.R. China

10 ^b National R & D Branch Center for Conventional Freshwater Fish Processing, Wuhan, Hubei 430070,
11 P.R. China

12 ^c School of Human and Social Sciences, University of West London, Middlesex TW8 9GA, UK

13 ^d CAS Key Laboratory of Brain Connectome and Manipulation, the Brain Cognition and Brain Disease
14 Institute (BCBDI), Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences;
15 Shenzhen-Hong Kong Institute of Brain Science-Shenzhen Fundamental Research Institutions,
16 Shenzhen, 518055, China.

17

18 * Corresponding author

19 **Shanbai Xiong**; Email: xiongsb@mail.hzau.edu.cn; Phone: 86-27-87288375;

20 **Abstract:** The micro-flowing water system can improve the flesh quality of freshwater fish using the
21 traditional pond farming method. However, the mechanism of this phenomenon has not yet been
22 explored. This study intends to examine the changes of metabolites in freshwater fish after treatment
23 with the micro-flowing purification system (MFPS). The UPLC-QTOF/MS based metabolomics method
24 was utilized to screen the metabolites and predict the major possible metabolic pathways after MFPS
25 treatment. There were 377 types of metabolites identified in the fish muscle, of which 54-71 represented
26 significant different metabolites identified during different stages of MFPS treatments. The main
27 mechanism of MFPS treatment in improving the quality of grass carp fish muscle was investigated, and
28 the MFPS treatment was shown to improve the flesh quality and the flavor of grass carp fish muscle.
29 This study could provide the theoretical basis for improving the quality of aquatic products.

30

31 **Keywords:** Micro-flowing purification system; Quality; Grass carp; Metabolomics; UPLC-QTOF/MS

1. Introduction

The annual yield of grass carp (*Ctenopharyngodon idella*) is about 25 million tons in the world, with 17.9% produced in China (Qiu, et al., 2020). Grass carp plays a major role in aquaculture, as it is the largest freshwater fish species. It is one of the most popular food items in China, and is often used as fresh fish or processed fish fillet products (Liu, et al., 2013). Because the cultivation of freshwater fish often uses a high-density cultivation mode, the cultivation of microorganisms in aquatic environments could easily cause the accumulation of undesirable flavors in the fish, which in turn would affect, to some extent, the degree of acceptance by consumers of the grass carp culture (Fuentes, et al., 2010; Rincón, et al., 2016).

Depuration treatment or short-term aquaculture can effectively improve the quality and acceptability of market-size fish before selling. For example, depuration improved the muscle quality of atlantic salmon (*salmo salar*), and decreased the content of earth-musty off-flavors (Burr, et al., 2012). By changing nitrate-nitrogen levels in recirculating aquaculture systems, the concentration of off-flavor compounds geosmin and 2- methylisoborneol in rainbow trout was reduced (Schrader, et al., 2013). Common carp had higher levels of polyunsaturated acids after being kept in clear water without feeding for several days (Zajic, et al., 2013). Furthermore, there was a significant improvement in the consumers' acceptability of the flavor of halibut after two weeks of depuration treatment (Drake, et al., 2010). In our previous study, the edible quality of grass carp muscle improved after using treatment of the micro-flowing water depuration system. The texture, taste and flavor of grass carp muscle appeared to significantly change during the purification process (Lv, et al., 2018). But the mechanism of the fish quality improvement through the depuration system treatment has not yet been reported, thus the mechanism of fish quality improvement should be further explored.

Metabolomics is an effective analytical method for the identification and quantification of small molecular endogenous metabolites in organisms (Patti, et al., 2013). It has been widely used in the food industry to investigate the metabolic changes of aquatic products under various physiological or storage conditions, specifically, after discovering the inherent mechanism and potential biomarker of monitoring food quality. Metabolomics has been applied to some aspects of aquatic products processing. For example, the NMR based metabolomics method was used to study changes of metabolites of sea cucumbers and crabs under different stress conditions (Ye, et al., 2016). GC-MS technology was used to characterize the metabolic profiles of crucian carps infected with bacteria, and to screen the important biosynthesis pathway and metabolic biomarkers (Guo, et al., 2014). Owing to its powerful separation ability, high throughput capacity, high resolution and detection sensitivity, HPLC-MS technology has been widely used in the study of aquatic products and metabolomics. For instance, the metabolic changes in zebrafish induced by environmental disturbance were investigated using HPLC-MS technology (De Sotto, et al., 2016). The performance of the HPLC-MS approach is excellent and has been broadly used for quantitative determination of meat quality and related metabolites or nutrients (Gil-Solsona, et al., 2019).

In the present study, market size grass carp were depurated using the micro-flowing water depuration system (MFPS). The UPLC-QTOF/MS based metabolomics method was utilized to explore the metabolites and their contents. The different metabolites were screened out and the key metabolic pathways related to the quality of grass carp muscle were investigated. Thus, the main mechanism of depuration treatment to improve the quality of grass carp fish was explored using UPLC-QTOF/MS based metabolomics and expected to provide the theoretical basis for improving the quality of aquatic products through regulating and controlling the depuration process.

45

46 **2. Materials and Methods**

47 **2.1 Chemical reagents**

48 2-Chloro-L-phenylalanine (Sigma Aldrich, St. Louis, MO, USA) and methanol (A.R.) (Sinopharm
49 Chemical Reagent Co. Ltd, Shanghai, P.R. China) were used in the sample preparation. MS - 222 (3-
50 Aminobenzoic acid ethyl ester methanesulfonate, CAS: 886-86-2, Shanghai yuanye Bio-Technology Co.,
51 Ltd, Shanghai, P.R. China) was used as the anesthetic typically administered to induce anesthesia in
52 fresh grass carp. Ultrapure water was obtained from the Milli-Q system (Millipore, Billerica, MA). The
53 mobile phase was composed of the following components: acetonitrile and methanol (HPLC grade,
54 Merck, Darmstadt, Germany), formic acid (Sigma Aldrich, St. Louis, MO, USA), phosphoric acid
55 (Kemiou Chemical Reagent Co. Ltd, Tianjin) and ultra-pure water (Merck, Darmstadt, Germany).

56 **2.2 Fish sample preparation**

57 All animal procedures were approved by the Animal Care and Use Committee of Huazhong
58 Agricultural University and performed in accordance with the Guidelines for Care and Use of Laboratory
59 Animals of Huazhong Agricultural University. The schematic diagram of the whole experiment is
60 illustrated in Fig. S1. Fresh grass carp (1.5~2.0 kg) from the fish farming in Chidong Lake (115.41°E,
61 30.10°N, Hubei, P.R. China) were transported in troughs (Long×Width×Height: 720cm×200cm×
62 150cm, n=30) in the micro-flowing water purification system (Lv, et al., 2018). There were ~125 kg fish
63 placed in every trough, and the depuration period lasted from October to December (2015). All fish were
64 deprived of food during the whole period. During the period of depuration, five grass carp were collected
65 every 10 days. The sampled fish were anaesthetized using the anesthetic MS - 222 (100 mg/L) and were
66 unconscious before slaughter. Fish were immediately gutted and segmented. At the end, the dorsal

muscle of each fish was sampled and kept at -80°C for further analysis.

2.3 Metabolites extraction

The fish muscle samples (~50 mg) were extracted with 800 µL methanol. Dichlorophenylalanine (10 µL, 2.9mg/mL) was added to the mixture according to the internal standard. All fish samples were grinded for 90s at 65Hz, vortexed for 30s and centrifuged for 15min (12000 rpm, 4°C). At the end, 200 µL supernatant was transferred to sample vials for further detection.

2.4 UPLC-QTOF/MS detection

Chromatographic evaluations were performed using a Waters Acquity UPLC-I Class (Waters Corp., Mil-ford, MA, USA). The UPLC system was equipped with a binary pump, micro degasser, an autosampler and a temperature-controlled column compartment. Chromatographic separations were achieved on an ACQUITY UPLC HSST3 ODS-SP column (1.8 µm, 2.1 mm × 100 mm; Waters, Ireland). The gradient system consisted of 0.1% formic acid in water in mobile phase A and 0.1% formic acid in acetonitrile in mobile phase B (0-2 min, A, 95%; 2-12 min, A, 95–5%; 12-15 min, A, 5%; 15-17 min, A, 5–95%; 17-20 min, A, 95%). The chromatographic separation condition was set as following: oven temperature-40 °C, injection volume-0.6 µL, flow rate-0.35 mL/min.

The separated components were detected with a Vevo G2-S QTOF mass spectrometer (Waters Corp., Mil-ford, MA, USA) equipped with both ESI ion modes (ESI- and ESI+). The detecting parameters were collected as following: Cone gas, 50 L/h; Ion source temperature, 120 °C (+)/110 °C(-); Desolvation gas temperature, 350 °C; Capillary voltage, 1.4 kV (ESI+)/1.3 kV (ESI-); Sample cone, 40V (ESI+) or 23V (ESI-); Gas flow, 600 L/h; Collision energy, 10–40 V; Ion energy: 1V; Scan time: 0.03s; Inter scan time: 0.02s; Scan range 50–1500 m/z. All analyses were performed using lockspray to ensure accuracy and reproducibility, with a concentration of 200 ng/mL of leucine enkephalin (m/z

556.2771) in positive ion mode and (m/z 554.2615) in negative ion mode locking quality.

2.5 Data processing

The raw UPLC-QTOF/MS data were initially transformed to CDF files using CDFbridge (Masslynx 4.1, Waters), then imported into XCMS software for peak alignment, peak detection, peak picking, peak filling and isotope elimination. The data were normalized with retention time, m/z and observation and peak intensity. The data matrix was analyzed using Multivariate Analysis (MVA) SIMCA-P 13.0 software (Umetrics AB, Umea, Sweden) for principal component analysis (PCA) and Orthogonal partial least squares discriminant analysis (OPLS-DA). According to the variable importance plot (VIP), the necessary conditions were set (variables scoring >1.5 and $p < 0.05$) to identify metabolites with significant differences between the control and micro-flowing water system treated groups.

In order to investigate detailed information of these potential significant metabolic biomarkers, the identification and quantification of these biomarkers were investigated using the detailed information detected by UPLC-QTOF/MS. At first, the accurate m/z obtained by multivariate statistical analysis was matched with metabolites from the online databases of Metlin (<https://metlin.scripps.edu/>) and KEGG (<http://www.kegg.com/>), which initially indicated the possible metabolic mechanism. At the end, the identification of the metabolites was validated with ion fragments, parent ion and the retention time. The quantification of metabolites was determined with the area of the peaks in EIC (Extracted ion chromatograms) extracted from TIC (Total ion chromatograms) (Fig. 1), which was automatically completed with the commercial software MassLynx 4.1 (Waters, Milford, MA, USA).

2.5 Determination of hypoxanthine and inosine

For the pretreatment and determination of the content of hypoxanthine and inosine in grass carp muscle, reference was made to a former method (Kuda, et al., 2008). Separation of hypoxanthine and

inosine was conducted with a reverse-phase column (Agilent C18, Agilent Technologies, USA). The mobile phase of phosphate buffer-methanol (85:15, pH 6.0) was used (flow rate: 0.7 ml/min; temperature: 35°C). The detection wavelength of eluent was monitored at 254 nm for hypoxanthine and inosine.

2.6 Statistical analysis:

The chemical analyses were repeated at least three times, and the results expressed as means $\pm$ SEM. Statistical analysis was performed with SPSS 22.0 (IBM, New York, USA) using one-way ANOVA. LSD adjustment was used to determine the significant difference between different groups. Significant differences were declared at $p < 0.05$. The statistical power analysis was used to validate the sample size for significant differences, which was implemented in the freeware GPower (Erdfelder, et al., 1996).

3. Results and discussion

3.1 UPLC-QTOF/MS analysis

The metabolites extracted from fish samples, which were treated with the MFPS under different purified periods (0d, 10d, 20d, 30d, 40d and 50d), were detected with UPLC-QTOF/MS technology in the positive (ESI+) and negative (ESI-) ESI modes. As an example, a typical series of peak intensity chromatogram of treated fish samples in the positive ESI mode are illustrated in Fig. 1. From Fig. 1, most endogenous metabolites could be detected in 18 min using UPLC-QTOF/MS technology.

According to the peak heights in the spectra, about 4420 peaks of positive ions (ESI+) and 4600 peaks of negative ions (ESI-) were detected which were extracted from TIC by the MassLynx (Fig. 1). By comparison with the NIST standard spectrum, there were 377 metabolites identified in the grass carp muscle, including 73 amino acids, 117 lipids, 56 nucleic acids, 123 sugars and carbon sources, and some inorganic acids. From Fig. 1, it is visually evident that there are some differences among these six

different groups in the TIC chromatograms. However, it is rather difficult to screen the potential different metabolites in fish muscle samples except by manually and intuitionally comparing the control and treated chromatograms. Therefore, application of the appropriate pattern recognition approach is very necessary for visualizing characteristic changes.

3.2 Principal component analysis (PCA)

As a non-supervised multivariate data analysis method, PCA is always used to give a comprehensive view of the clustering trend for the multidimensional data (Boyan, et al., 2013). In order to gain an overview of the fish muscle metabolic profiling, PCA was initially used for the analysis of UPLC-QTOF/MS data.

To determinate whether the micro-flowing purification treatment is able to influence metabolic patterns of grass carp muscle and to screen the characteristic metabolites with significant concentration changes (*i.e.* potential biomarkers), the PCA approach was utilized to conduct a model with the ES+ and ES-data, respectively. The score plots of PCA are illustrated in Fig. 2A (ES+) and Fig. S2A (ES-). The validity of the PCA model was evaluated using the correlation coefficient R^2 and the cross-validation correlation coefficient Q^2 . R^2 is defined as the proportion of variance in the data explained by the models and indicates goodness of fit. Q^2 is defined as the proportion of variance in the data predicted by the model and indicates predictability. R^2 and Q^2 of PCA models for the control group and the other five treated groups were 0.514 and 0.384, 0.226 and 0.0198 for the ES+ and ES- modes, respectively. Thus, the PCA method was more useful for the analysis of data in the ES+ mode. It is clearly noticeable that the samples in the control group were completely separated compared with the samples in the micro-flowing purification treated model groups. Thus, the metabolic pattern in the grass carp muscle was

significantly altered after the purification process.

Furthermore, the PCA method also illustrated the separation of the control group and the other five treated groups under the ES+ mode, respectively. The statistical parameters (R^2 and Q^2) of all PCA models are illustrated in Table S1. Results from the PCA model showed that this method could generate good results for the discrimination between samples from the control and treated groups, and the performance from the positive ion mode was better than from the negative ion mode. With various periods of micro-flowing purification system treatment, the different metabolite compositions were further explored between the purified groups and control group.

3.3 OPLS-DA analysis

In order to maximize the separation among the different grass carp groups and focus on metabolic variations at different purification periods, the data for both ESI+ and ESI- acquisition modes were performed using a supervised multivariate data analysis approach OPLS-DA (Aru, et al., 2016) to build more intensive and accurate models and explore the real trend and grouping the multidimensional data.

The score plots of OPLS-DA are shown in Fig. 2B-2F and Fig. S2B-S2F. Compared with the control group, the cluster separations are very good at different micro-flowing purification treated time points. The results mean that the metabolic patterns in grass carp were significantly changed during the purification processing. The performance of OPLS-DA was distinctly better than the PCA models. All models exhibited reasonable separation between groups and acceptable goodness-of-fit (R^2Y) and goodness-of-prediction (Q^2Y). The statistical parameters of the OPLS-DA models (R^2X , R^2Y and Q^2) are summarized in Table S1. The results not only indicated the strong explanatory power of the data and the good predictive ability of all permuted OPLS-DA models, but also demonstrated a robust metabolic

177 difference between the micro-flowing purification systems treated grass carp and the controls.

179 3.4 Identification of different metabolites for micro-flowing system purification treatment

180 Based on the judgement criteria ($VIP > 1.5$, $p < 0.05$) for determining the different metabolites of
181 the OPLS-DA model, there were 145 metabolites identified in fish muscle samples compared with the
182 control group and purified groups under different purification periods. The retention time, molecular
183 weight and relative amount of potential different metabolites are demonstrated in Table 1 and Table S2.
184 The majority of identified metabolites were lipids and secondary metabolites of lipids under different
185 purification times (Fig. 3), which indicates that the micro-flowing purification treatment could alter the
186 lipid metabolism in grass carp. The phenomenon that most metabolites were lipid decomposition
187 products shows that the fat consumption increased during the purification process, which is consistent
188 with our previous study (Lv, et al., 2018). There are many phospholipid components involved in the lipid
189 metabolites, which indicates that a large number of signal molecules (such as phosphatidylserine,
190 phosphatidylethanolamine, *etc.*) contributed to the metabolism caused by the purification treatment.

191 In order to further determine the biomarkers in different metabolites of grass carp during the
192 micro-flowing purification treatment, the repeated metabolites and their corresponding relative contents
193 in different metabolites of each comparable group were searched and analyzed. The selected metabolites
194 were further identified by the molecular weight (m/z) in the Metabolites and Tandem MS database
195 (METLIN) and KEGG database. Based on the variety and content changes of different metabolites of
196 grass carp muscle and the reported metabolic pathway of each metabolite in KEGG database, the
197 metabolic pathway changes of grass carp muscle during purification were obtained (Fig. 4). A total of
198 16 significant differential endogenous metabolites were found which showed an increasing and

decreasing trend from the beginning to the end of the purification process (Table 1). Among these different metabolites, xanthine, hypoxanthine, trimethylamine oxide, alanine and IMP (Inosine monophosphate) are the most common precursor substances. The changes in their amount affected the quality of fish muscle. The causes of these metabolic differences are further discussed. The species and the changes in the amount of the identified metabolites, combined with the metabolic pathway reported from the KEGG database, the potential metabolic pathways of micro-flowing purification system treated grass carp are all illustrated in Fig. 4.

The trend of phospholipid metabolites, such as phosphatidylserine and phosphatidylcholine, indicate that the synthesis and metabolism of signal metabolites are enhanced and kept up-regulating in the early stage of purification. From the decrease in the amount of arachidonic acid, docosanoic acid, leukotriene B4 and the increase in short chain alkyl coenzyme A, it can be seen that the decomposition and metabolism of fatty acids increased in grass carp muscle after purification, and the consumption of fatty acids was the main source of energy. The trend of histidine, homocysteine, alanine and other amino acid metabolites showed that the decomposition rate of amino acid was greater than its synthesis rate during the purification process, which led to the accumulation of alanine in fish muscles. After 20 days of purification, the level of IMP synthesis was enhanced, and its conversion rate of hypoxanthine and xanthine was inhibited. The content of trimethylamine oxide and its related metabolites decreased during the purification process, its metabolic substrate carnitine did not change significantly, while the content of choline increased, indicating that the generation pathway of trimethylamine was inhibited.

In order to validate the results of metabolic analysis, the content of hypoxanthine and inosine in grass carp muscle during purification was measured using the HPLC method. The corresponding results are shown in Fig. 5. After the micro-flowing system purification treatment, the contents of these two

kinds of endogenous metabolites in fish muscle significantly decreased, which is consistent with the results of metabolomics. It indicates that the purification treatment could induce the inhibition of catabolism of IMP to hypoxanthine in the muscle of grass carp. Combined with the trend in different metabolites mentioned above which are shown in Table 1 and Table S1, it can be seen that the main metabolites related to fish flavor changed significantly in 20 days during the purification process. The content of the major metabolites can remain stable for a relatively long time, which indicates that the micro-flowing purification treatment can improve the quality and flavor of fish meat, as long as the period of purification is not less than 20 days.

3.5 Metabolic changes in the grass carp muscle during MFPS treatment

Due to grass carp not feeding during the MFPS treatment period, the fish protein decomposed and stored lipids in tissues in order to maintain the energy supply and daily life activities, hence the contents of protein and lipids in the fish muscle showed a consistent downward trend. Basically, the consumption of protein and lipids in grass carp fish muscle is according to the regulation of energy metabolism under prolonged starvation in fish. In other words, the fish metabolism can be divided into three stages during the MFPS treatment: (I) In the first 20 days of the MFPS treatment, the metabolic rates of both protein and lipids are very fast; (II) During 20d-40d of the MFPS process, the catabolism of grass carp muscle is dominated by lipids until reaching the limit in the body of the fish; (III) In the last 10 days of MFPS treatment, protein catabolism plays an important role in the fish body metabolism in supplying the energy (Bar, 2014).

From Table 1 and Table S2, it can be observed that the content of NADP in the tissues significantly increased at 10 days after purification, which indicates that the energy supply of grass carp

was insufficient during the starvation period. As well as at the end of purification, NADP accumulated due to the decrease in energy supply caused by the reduction in lipid metabolism. This phenomenon is consistent with the expression trend of NADPH related enzymes synthesis found in liver cells of rainbow trout during the starvation phase, as the activities of glucose-6-phosphate dehydrogenase and malic acid enzyme in fish significantly decreased, and the synthesis of NADPH decreased (Nagana Gowda, et al., 2016). After 20-40 days of purification, the degree of metabolic protein and lipid was enhanced, and the synthesis of NADPH returned to normal level. The contents of intermediate products of fatty acid metabolism in fish muscle, such as 3-oxhexyl coenzyme A and 3-oxlauroyl coenzyme A increased significantly after 10 and 20 days' purification, respectively. At the same time, the contents of fatty acids including docosanoic and arachidonic acid decreased, and so did the contents of secondary products of lipid synthesis such as leukotriene, indicating that the consumption of fat was increased and the anabolism weakened. The fat content in the grass carp back muscle was measured but did not significantly change during the first two stages, it however slightly increased in the first 10 days of purification. These results are similar to the purification study of the cod, which reported that during the purification process, the fish consumed the lipid stored in the fish body in the following order, the liver and other internal organs, the abdomen and the digestive tract and the lipid reserved in the muscle (Palmeri, et al., 2008). This style of lipid metabolism also occurred in the carp (*Cyprinus carpio*) and the roach under the prolonged starvation condition, but for the rainbow and brown trout the opposite was true (Regost, et al., 2001).

In this study, the grass carp duration of stage I is similar to that of the striped bass and rainbow trout, while the stage II is shorter (Jönsson, et al., 2007). During the purification process, the content of amino acids such as homocysteine, histidine and homotyrosine decreased (Table 1 and S2), indicating

that the catabolism of protein and amino acids is also very active, and may be related to the synthesis of phosphatidylserine except when involved in supplying energy to the body.

The synthesis of phosphatidylserine is up-regulated to a greater extent during the purification process and accompanied by significant changes in the contents of other phospholipids such as phosphatidylglycerol, phosphatidylcholine, phosphatidylinositol and ceramide metabolites. Phosphatidylserine is an important component of the membrane structure in cells, and it is widely involved in the regulation of the cell signaling pathway, cell proliferation and apoptosis (Fadok & Henson, 2003). In this experiment, the relative content of phosphatidylserine was significantly increased, which is consistent with the phenomenon of grass carp myoblast reported by Sousa et al (Sousa, et al., 2016). In other words, under the condition of diminished nutrient supply, phosphatidylserine is redistributed in the membrane and entered during apoptosis, while proteins, amino acids, nucleic acids and other substances in cells are reused to maintain life activities (Zhou, et al., 2016). The up-regulation of phosphatidylserine indicates that stress starvation during the purification process makes the regulation of cell apoptosis of grass carp more active. Thus, the accumulation of phosphatidylserine can play an important role in the protection of nerve cells (Akbar, et al., 2006). The data in this experiment show that the grass carp body can protect the nervous system by up-regulating the synthesis of phosphatidylserine under stress starvation.

3.6 Differential metabolites related to the flesh quality of grass carp muscle

Flesh quality is defined by the nutritive value and organoleptic properties as determined by consumers (Grigorakis, 2007). The quality and flavor of the fish should be based on the flesh protein (Grigorakis, 2007), peptides (Jürss & Bastrop, 1995), fatty acids (Grigorakis, 2007), and other

metabolites, which have been studied in our study, and protein and amino acids not only belong to the nutritional components, but also influence the flavors of fish products (Jiang, et al., 2016). A period of purging and food deprivation before harvesting is always the common procedure used for improving the quality and flavor of the fish, and the purged fish (2-4 weeks purging) were found to have a consistent and desirable flavor (Palmeri, et al., 2008). Thus, changes in metabolites, especially amino acids, which play important roles in determining the quality and flavor of the fish were investigated in the current study.

3.6.1 Inosine monophosphate (IMP)

The content of IMP was significantly increased after 20 days of purification, while the contents of jaundice and hypoxanthine in fish muscle were considerably reduced. IMP may be degraded into xanthine and hypoxanthine through safrrole and inosinein converted in the fish body, respectively (Fig. 4). The trend of the relative contents of IMP, jaundice and hypoxanthine which changed in the muscle cells of the grass carp indicates that the decomposition of IMP was inhibited after the treatment of MPFS. Therefore, the amount of IMP was accumulated in the fish muscle.

At the same time, IMP can be decomposed by the metabolite AICAR of 5-phosphate- α -ribose-1-diphosphate (PRPP). The other aspect of PRPP metabolism is histidine, and the content of histidine in muscle was decreased (Table S2). It can be seen that the PRPP metabolism during the purification procedure was mainly carried out with the synthesis of AICAR to IMP. When Kuhla et al studied the expression of dairy cow protein, they found that the expression of AICAR methyltransferase/IMP cyclase was up-regulated under starvation, and the synthesis of IMP was enhanced, thereby supplementing and maintaining the total amount of adenine nucleotides with IMP (Kuhla, et al., 2007).

Furthermore, Hypoxanthine-guanine phosphoribosyl transferase (HGPRT) (HGPRT) is one of the

main enzymes that catalyze the synthesis of IMP and XMP from hypoxanthine and xanthine. It has been reported that the increased expression of HGPRT can enhance the reuse of purines in walrus under starvation during the purification process, and maintain a stable amount of purine nucleotides (Sonanez-Organis, et al., 2012). The senegalese sole also showed the same tendency of expressing HGPRT under repeated stress (Cordeiro, et al., 2012). In this study, the increase of IMP content in grass carp muscle might be the result of the joint action between the increased expressions of IMP cyclase and HGPRT, and also the reduced substrate content of hypoxanthine and xanthine.

IMP is the major source of umami in meat products, therefore, the accumulation of IMP in muscle can effectively improve the flavor of food (Hong, et al., 2017). At the same time, xanthine and hypoxanthine are regarded as the sources of bitterness in foods (Bedini, et al., 2013). Therefore, the MFPS treatment is beneficial in improving umami substances and reducing the contents of bitter substances in the fish muscle that may partly explain the improvement of grass carp quality after the purification process.

3.6.2 Trimethylamine N-Oxide (TMAO)

After the MFPS treatment, the content of TMAO in the grass carp muscle was lower than the detection limit, and the contents of secondary metabolites, dimethylamine and methyl coenzyme M, fell by more than 40% and 60%, respectively. The potential substrates of TMAO include carnitine, trimethyllysine and choline (Narasimhan, et al., 2018). It has been reported that TMAO is mainly derived from the metabolism of carnitine and choline in foods through the action of intestinal microbes, further metabolized and accumulated in various tissues of the body (Cho, et al., 2017). Intestinal bacteria could influence the production of TMAO from carnitine, and carnitine the composition of gut bacteria. The gut bacteria environment was seriously influenced by the depuration procedure (Oliveira, et al., 2011),

such as MPFS in the current study. Therefore, while the grass carp was under starvation during the purification process, the major source of TMAO was interrupted. The amount of TMAO and the corresponding downstream metabolites were significantly reduced in the metabolic process. The same phenomenon has also been observed in the crab aquaculture, as starvation contributed to the decrease of TMAO content in muscles (Ye, et al., 2016).

TAMO can generate trimethylamine and dimethylamine catalyzed by TAMO reductase, which are the two main off-flavor components in fish or aquatic products (Fu, et al., 2006). In this study, the content of TAMO and its corresponding metabolites were decreased during the process of MFPS treatment, indicating that the purification process helped to reduce the off flavor in muscle and thereby improved the quality of grass carp.

3.6.3 Alanine

Considered as the primary metabolites of amino acid metabolism, such as glutamate, aspartic acid and cysteine, alanine can be further metabolized into pyruvic acid and enters the TCA cycle and energy metabolism. During the purification process, the degree of amino acid decomposition was high, and the content of alanine obtained enriched (Table S2). The major form of energy supply is fatty acid metabolism and the limit of fatty acids in the organism was reached after 50 days of purification. After that, the style of energy supply changed into protein and amino acid, while the content of alanine recovered in comparison to the control group. Furthermore, the enriched alanine could also be caused by the reduction in the activation of AlaAT (Alanine Aminotransferase) due the starvation and purification procedure by the MPFS treatment, which could catalyze alanine into α -ketoglutarate in the cycle to transform alanine to pyruvate and glutamate (Jürss & Bastrop, 1995).

The glutamic acid, glycine, alanine and histidine in fish muscle contributed directly to the flavor

of fish, thus the higher the content of these four types of amino acids, the better the quality of fish meat (Luo, et al., 2017). Therefore, it is reasonable to conclude that the purification treatment can improve the flavor of fish meat through the enrichment of alanine.

3.6.4 Flesh quality and metabolites

Through analysis of the major differential metabolites and the reported metabolic pathway database, the mechanism of the MFPS treatment to improve flesh quality of grass carp deduced the following: not feeding during purification produced starvation stress which led to the ATP level of grass carp muscle down-regulating and induced significant changes in the purine metabolism pathway in muscle cells, which transferred from hypoxanthine to IMP synthesis and caused the lower concentration of hypoxanthine. At the same time, the decrease in ATP level induced a large number of apoptosis signal molecules to synthesize, thereby accelerating the apoptosis process of some cells, which utilized the stored protein and lipid to generate energy, and finally fed back together with IMP to maintain the normal ATP level. UPLC-QTOF/MS based metabolomics explored whether starvation is the main driving force of metabolic pathway changes which promotes the accumulation of IMP and the results indicate that some of the flavors of amino acids in fish muscle improved significantly. However, the consumption of lipids affected the texture and taste of fish.

Due to the lack of supply of carnitine which is one of TMAO substrates during starvation, the accumulation of TMAO and the corresponding downstream metabolites were reduced, which caused the off flavors of fish muscle to decrease. Among the differential metabolites, hypoxanthine, IMP and alanine showed the greatest influence on the flavor of fish muscle, and their content changed significantly during the micro-flowing purification process, which can be used as the potential biomarkers for grass carp purification treatment.

4. Conclusions

There were 377 metabolites detected in grass carp muscle using UPLC-QTOF/MS, and 54~71 different metabolites were screened out during the MFPS treatment process. During the purification process, the catabolism of protein and fat released energy to the body and provided substrates for various metabolic activities of the body. The starvation stress during purification significantly changed the relative contents of phosphatidylserine and other phospholipid metabolites in grass carp muscle, while enhancing the synthesis of phosphatidylserine. Under the joint action of amino acid catabolism and phosphatidylserine synthesis, the contents of histidine, cysteine and other amino acids in fish muscle decreased, and in the end, alanine was enriched. Alanine contributed directly to the flavor of fish or fish products. Due to the accumulation of IMP in fish muscle after the purification treatment, the content of bitter substances like xanthine and the off-flavor substances such as TMAO decreased, which could have improved the flavor and quality of grass carp. Hypoxanthine, IMP and alanine are potential biomarkers of grass carp in the process of micro-flowing water system purification treatment. These results confirm the potential use of metabolomics as a reliable method to explore the mechanism of fish quality improvement during the purification treatment process.

Acknowledgements

All the authors would like to express our gratitude to Prof. Fang Xu (Faculty of Mathematics and Statistics, Hubei University) for discussing the results of statistical analysis and statistical power analysis. All authors express their appreciation to the Shanghai Sensichip Infotech Co. (Shanghai, China) for the detection of the samples. This research was financially supported by the Fundamental Research Funds

for the Central Universities of China (No. 2662017PY021 and 2662019PY031), the National Natural Science Foundation of China (No. 31772047) and Agriculture Research System of China (CARS-45-27).

References

- Akbar, M., Baick, J., Calderon, F., Wen, Z., & Kim, H.-Y. (2006). Ethanol promotes neuronal apoptosis by inhibiting phosphatidylserine accumulation. *Journal of Neuroscience Research*, 83(3), 432-440.
- Aru, V., Pisano, M. B., Savorani, F., Engelsen, S. B., Cosentino, S., & Cesare, M. F. (2016). Metabolomics analysis of shucked mussels' freshness. *Food Chemistry*, 205, 58-65.
- Bar, N. (2014). Physiological and hormonal changes during prolonged starvation in fish. *Canadian Journal of Fisheries & Aquatic Sciences*, 71(10), 1447-1458.
- Bedini, A., Zanolli, V., Zanardi, S., Bersellini, U., Dalcanale, E., & Suman, M. (2013). Rapid and simultaneous analysis of xanthines and polyphenols as bitter taste markers in bakery products by FT-NIR spectroscopy. *Food Analytical Methods*, 6(1), 17-27.
- Boyan, G., Yingjian, L., Yi, S., Pei, C., & Liangli Lucy, Y. (2013). Differentiating organic and conventional sage by chromatographic and mass spectrometry flow injection fingerprints combined with principal component analysis. *Journal of Agricultural & Food Chemistry*, 61(12), 2957-2963.
- Burr, G. S., Wolters, W. R., Schrader, K. K., & Summerfelt, S. T. (2012). Impact of depuration of earthy-musty off-flavors on fillet quality of Atlantic salmon, *Salmo salar*, cultured in a recirculating aquaculture system. *Aquacultural Engineering*, 50, 28-36.
- Cho, C. E., Taesuan, S., Malysheva, O. V., Bender, E., Tulchinsky, N. F., Yan, J., Sutter, J. L., & Caudill, M. A. (2017). Trimethylamine-N-oxide (TMAO) response to animal source foods varies among healthy young men and is influenced by their gut microbiota composition: A randomized controlled trial. *Molecular Nutrition & Food Research*, 61(1), 1600324.
- Cordeiro, O. D., Silva, T. S., Alves, R. N., Costas, B., Wulff, T., Richard, N., Vareilles, M. d., Conceição, L. E. C., & Rodrigues, P. M. (2012). Changes in liver proteome expression of senegalese sole (*solea senegalensis*) in response to repeated handling stress. *Marine Biotechnology*, 14(6), 714-729.
- De Sotto, R., Medriano, C., Cho, Y., Seok, K.-S., Park, Y., & Kim, S. (2016). Significance of metabolite extraction method for evaluating sulfamethazine toxicity in adult zebrafish using metabolomics. *Ecotoxicology and Environmental Safety*, 127, 127-134.
- Drake, S. L., Drake, M. A., Sanderson, R., Daniels, H. V., & Yates, M. D. (2010). The effect of purging time on the sensory properties of aquacultured southern flounder (*paralichthys lethostigma*). *Journal of Sensory Studies*, 25(2), 246-259.
- Erdfelder, E., Faul, F., & Buchner, A. (1996). GPOWER: A general power analysis program. *Behavior Research Methods, Instruments, & Computers*, 28(1), 1-11.
- Fadok, V. A., & Henson, P. M. (2003). Apoptosis: giving phosphatidylserine recognition an assist - with a twist. *Current Biology*, 13(16), R655-R657.
- Fu, X. Y., Xue, C. H., Miao, B. C., Liang, J. N., Li, Z. J., & Cui, F. X. (2006). Purification and characterization of trimethylamine-N-oxide demethylase from jumbo squid (*Dosidicus gigas*). *Journal of Agricultural & Food Chemistry*, 54(3), 968-972.

- Fuentes, A., Fernández-Segovia, I., Serra, J. A., & Barat, J. M. (2010). Comparison of wild and cultured sea bass (*Dicentrarchus labrax*) quality. *Food Chemistry*, 119(4), 1514-1518.
- Gil-Solsona, R., Caldach-Giner, J. A., Nacher-Mestre, J., Lacalle-Bergeron, L., Sancho, J. V., Hernández, F., & Pérez-Sánchez, J. (2019). Contributions of MS metabolomics to gilthead sea bream (*Sparus aurata*) nutrition. Serum fingerprinting of fish fed low fish meal and fish oil diets. *Aquaculture*, 498, 503-512.
- Grigorakis, K. (2007). Compositional and organoleptic quality of farmed and wild gilthead sea bream (*Sparus aurata*) and sea bass (*Dicentrarchus labrax*) and factors affecting it: A review. *Aquaculture*, 272(1), 55-75.
- Guo, C., Huang, X.-y., Yang, M.-j., Wang, S., Ren, S.-t., Li, H., & Peng, X.-x. (2014). GC/MS-based metabolomics approach to identify biomarkers differentiating survivals from death in crucian carps infected by *Edwardsiella tarda*. *Fish & Shellfish Immunology*, 39(2), 215-222.
- Hong, H., Regenstein, J. M., & Luo, Y. (2017). The importance of ATP-related compounds for the freshness and flavor of post-mortem fish and shellfish muscle: A review. *Critical Reviews in Food Science & Nutrition*, 57(9), 1787-1798.
- Jönsson, E., Forsman, A., Einarsdottir, I. E., Kaiya, H., Ruohonen, K., & Björnsson, B. T. (2007). Plasma ghrelin levels in rainbow trout in response to fasting, feeding and food composition, and effects of ghrelin on voluntary food intake. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 147(4), 1116-1124.
- Jürss, K., & Bastrop, R. (1995). Chapter 7 Amino acid metabolism in fish. In P. W. Hochachka & T. P. Mommsen (Eds.), *Biochemistry and Molecular Biology of Fishes*, vol. 4 (pp. 159-189): Elsevier.
- Jiang, W.-D., Wu, P., Tang, R.-J., Liu, Y., Kuang, S.-Y., Jiang, J., Tang, L., Tang, W.-N., Zhang, Y.-A., Zhou, X.-Q., & Feng, L. (2016). Nutritive values, flavor amino acids, healthcare fatty acids and flesh quality improved by manganese referring to up-regulating the antioxidant capacity and signaling molecules TOR and Nrf2 in the muscle of fish. *Food Research International*, 89, 670-678.
- Kuda, T., Fujita, M., Goto, H., & Yano, T. (2008). Effects of retort conditions on ATP-related compounds in pouched fish muscle. *LWT - Food Science and Technology*, 41(3), 469-473.
- Kuhla, B., Kuhla, S., Rudolph, P. E., Albrecht, D., & Metges, C. C. (2007). Proteomics analysis of hypothalamic response to energy restriction in dairy cows. *Proteomics*, 7(19), 3602-3617.
- Liu, D., Liang, L., Xia, W., Regenstein, J. M., & Zhou, P. (2013). Biochemical and physical changes of grass carp (*Ctenopharyngodon idella*) fillets stored at -3 and 0°C. *Food Chemistry*, 140(1), 105-114.
- Luo, J. B., Feng, L., Jiang, W. D., Liu, Y., Wu, P., Jiang, J., Kuang, S. Y., Tang, L., Tang, W. N., & Zhang, Y. A. (2017). Physical and flavor characteristics, fatty acid profile, antioxidant status and nrf2-dependent antioxidant enzyme gene expression changes in young grass carp (*Ctenopharyngodon idella*) fillets fed dietary valine. *Plos One*, 12(1), e0169270.
- Lv, H., Hu, W., Xiong, S., You, J., & Fan, Q. (2018). Depuration and starvation improves flesh quality of grass carp (*Ctenopharyngodon idella*). *Aquaculture Research*, 49(9), 3196-3206.
- Nagana Gowda, G. A., Abell, L., Lee, C. F., Tian, R., & Raftery, D. (2016). Simultaneous analysis of major coenzymes of cellular redox reactions and energy using *ex vivo* ¹H NMR spectroscopy. *Analytical Chemistry*, 88(9), 4817-4824.
- Narasimhan, R., Coras, R., Rosenthal, S. B., Sweeney, S. R., Lodi, A., Tiziani, S., Boyle, D., Kavanaugh, A., & Guma, M. (2018). Serum metabolomic profiling predicts synovial gene expression in rheumatoid arthritis. *Arthritis research & therapy*, 20(1), 164-164.
- Oliveira, J., Cunha, A., Castilho, F., Romalde, J. L., & Pereira, M. J. (2011). Microbial contamination and purification of bivalve shellfish: Crucial aspects in monitoring and future perspectives – A mini-

- review. *Food Control*, 22(6), 805-816.
- Palmeri, G., Turchini, G. M., Caprino, F., Keast, R., Moretti, V. M., & Silva, S. S. D. (2008). Biometric, nutritional and sensory changes in intensively farmed Murray cod (*Maccullochella peelii peelii*, Mitchell) following different purging times. *Food Chemistry*, 107(4), 1605-1615.
- Patti, G. J., Yanes, O., & Siuzdak, G. (2013). Metabolomics: the apogee of the omic trilogy. *Nature Reviews Molecular Cell Biology*, 13(4), 263-269.
- Qiu, D.-K., Zhao, Z., Ma, R., Guo, Z.-R., Jia, Y.-J., Zhang, C., Wang, G.-X., & Zhu, B. (2020). Antigen epitope screening of grass carp reovirus and its protective immunity assessment for grass carp. *Aquaculture*, 515, 734550.
- Regost, C., Arzel, J., Cardinal, M., Laroche, M., & Kaushik, S. J. (2001). Fat deposition and flesh quality in seawater reared, triploid brown trout (*Salmo trutta*) as affected by dietary fat levels and starvation. *Aquaculture*, 193(3), 325-345.
- Rincón, L., Castro, P. L., Álvarez, B., Hernández, M. D., Álvarez, A., Claret, A., Guerrero, L., & Ginés, R. (2016). Differences in proximal and fatty acid profiles, sensory characteristics, texture, colour and muscle cellularity between wild and farmed blackspot seabream (*Pagellus bogaraveo*). *Aquaculture*, 451, 195-204.
- Schrader, K. K., Davidson, J. W., & Summerfelt, S. T. (2013). Evaluation of the impact of nitrate-nitrogen levels in recirculating aquaculture systems on concentrations of the off-flavor compounds geosmin and 2-methylisoborneol in water and rainbow trout (*Oncorhynchus mykiss*). *Aquacultural Engineering*, 57, 126-130.
- Sonanez-Organis, J. G., Vazquez-Medina, J. P., Zenteno-Savin, T., Aguilar, A., Crocker, D. E., & Ortiz, R. M. (2012). Prolonged fasting increases purine recycling in post-weaned northern elephant seals. *Journal of Experimental Biology*, 215(9), 1448-1455.
- Sousa, B., Melo, T., Campos, A., Moreira, A. S. P., Maciel, E., Domingues, P., Carvalho, R. P., Rodrigues, T. R., Girão, H., & Domingues, M. R. M. (2016). Alteration in phospholipidome profile of myoblast H9c2 cell line in a model of myocardium starvation and ischemia. *Journal of Cellular Physiology*, 231(10), 2266-2274.
- Ye, Y., Xia, M., Mu, C., Li, R., & Wang, C. (2016). Acute metabolic response of *Portunus trituberculatus* to *Vibrio alginolyticus* infection. *Aquaculture*, 463, 201-208.
- Zajic, T., Mraz, J., Sampels, S., & Pickova, J. (2013). Fillet quality changes as a result of purging of common carp (*Cyprinus carpio* L.) with special regard to weight loss and lipid profile. *Aquaculture*, 400-401, 111-119.
- Zhou, H., Wang, q., Yuan, D., Wang, J., Yang, H., Wu, H., Jian, J., Yang, D., Huang, N., Haisch, C., Jiang, Z., & Chen, S. (2016). Early apoptosis real-time detection by label-free SERS based on externalized phosphatidylserine. *Analyst*, 141, 4293-4298.

Figure legends:

Fig. 1. A series examples of chromatograms of representative fish muscle samples detected using UPLC-QTOF/MS under positive ion mode at different depuration processing times (It was arranged as 0d, 10d, 20d, 30d, 40d, 50d in depuration from top to bottom). *Note: TIC: Total ion chromatograms; EIC: Extracted ion chromatograms.*

Fig. 2. Results of PCA analysis (A) and score plots (B-F) of PLS-DA analysis of metabolites detected using UPLC-QTOF/MS equipped with both ESI+ mod which was obtained from grass carp muscle during different depuration periods. *Note: B: 0 day vs 10 days; C: 0 day vs 20 days; D: 0 day vs 30 days; E: 0 day vs 40 days; F: 0 day vs 50 days.*

Fig. 3. Effect of depuration time on the composition of metabolic types in grass carp muscle. *Note: A/B/C/D/E/F represent samples under depuration for 0/10/20/30/40/50 days.*

Fig. 4. Effect of depuration on metabolic pathways of grass carp muscle. *Note: Different colors of metabolites represent change of comparative content, red: increased; blue: not significantly changed; green: decreased. Reference: KEGG database, 2018.*

Fig. 5. Effect of depuration time on content of hypoxanthine (A) and inosine (B) in grass carp muscle