



## **UWL REPOSITORY**

**repository.uwl.ac.uk**

The prevalence, antibiotic resistance and mecA characterization of coagulase negative staphylococci recovered from non-healthcare settings in London, UK

Xu, Zhen, Shah, Haroun N., Misra, Raju, Chen, Jiazhen, Zhang, Wenhong, Liu, Yuting, Cutler, Ronald R. and Mkrтчyan, Hermine (2018) The prevalence, antibiotic resistance and mecA characterization of coagulase negative staphylococci recovered from non-healthcare settings in London, UK. *Antimicrobial Resistance & Infection Control*, 7 (1). ISSN 2047-2994

<http://dx.doi.org/10.1186/s13756-018-0367-4>

This is the Published Version of the final output.

**UWL repository link:** <https://repository.uwl.ac.uk/id/eprint/7306/>

**Alternative formats:** If you require this document in an alternative format, please contact: [open.research@uwl.ac.uk](mailto:open.research@uwl.ac.uk)

**Copyright:** Creative Commons: Attribution 4.0

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy:** If you believe that this document breaches copyright, please contact us at [open.research@uwl.ac.uk](mailto:open.research@uwl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

**Rights Retention Statement:**

RESEARCH

Open Access



# The prevalence, antibiotic resistance and *mecA* characterization of coagulase negative staphylococci recovered from non-healthcare settings in London, UK

Zhen Xu<sup>1,2</sup>, Haroun N. Shah<sup>3</sup>, Raju Misra<sup>4</sup>, Jiazhen Chen<sup>5</sup>, Wenhong Zhang<sup>5</sup>, Yuting Liu<sup>6</sup>, Ronald R. Cutler<sup>2</sup> and Hermine V. Mkrtchyan<sup>2,7\*</sup>

## Abstract

**Background:** Coagulase negative staphylococci (CoNS) are important reservoirs of antibiotic resistance genes and associated mobile genetic elements and are believed to contribute to the emergence of successful methicillin resistant *Staphylococcus aureus* (MRSA) clones. Although, these bacteria have been linked to various ecological niches, little is known about the dissemination and genetic diversity of antibiotic resistant CoNS in general public settings.

**Methods:** Four hundred seventy-nine samples were collected from different non-healthcare/general public settings in various locations ( $n = 355$ ) and from the hands of volunteers ( $n = 124$ ) in London UK between April 2013 and Nov 2014.

**Results:** Six hundred forty-three staphylococcal isolates belonging to 19 staphylococcal species were identified. Five hundred seventy-two (94%) isolates were resistant to at least one antibiotic, and only 34 isolates were fully susceptible. Sixty-eight (11%) *mecA* positive staphylococcal isolates were determined in this study. SCC*mec* types were fully determined for forty-six isolates. Thirteen staphylococci (19%) carried SCC*mec* V, followed by 8 isolates carrying SCC*mec* type I (2%), 5 SCC*mec* type IV (7%), 4 SCC*mec* type II (6%), 1 SCC*mec* type III (2%), 1 SCC*mec* type VI (2%), and 1 SCC*mec* type VIII (2%). In addition, three isolates harboured a new SCC*mec* type 1A, which carried combination of class A *mec* complex and *ccr* type 1.

MLST typing revealed that all *S. epidermidis* strains possess new MLST types and were assigned the following new sequence types: ST599, ST600, ST600, ST600, ST601, ST602, ST602, ST603, ST604, ST605, ST606, ST607 and ST608.

**Conclusions:** The prevalence of antibiotic resistant staphylococci in general public settings demonstrates that antibiotics in the natural environments contribute to the selection of antibiotic resistant microorganisms. The finding of various SCC*mec* types in non-healthcare associated environments indicates the complexity of SCC*mec*. We also report on new MLST types that were assigned for all *S. epidermidis* isolates, which demonstrates the genetic variability of these isolates.

**Keywords:** CoNS, Antibiotic resistance, SCC*mec*, MLST

\* Correspondence: [h.mkrtyan@uel.ac.uk](mailto:h.mkrtyan@uel.ac.uk)

<sup>2</sup>School of Biological and Chemical Sciences, Queen Mary University of London, London, UK

<sup>7</sup>School of Health, Sport and Biosciences, University of East London, E1 4NS, London, UK

Full list of author information is available at the end of the article



## Background

Staphylococci are the most frequently isolated nosocomial pathogens, accounting for 30% of hospital associated infections [1]. Despite, that the high virulence of *S. aureus* has been evidenced in many studies [2], it is believed that coagulase-negative staphylococci (CoNS) act as an important reservoir of antimicrobial resistance genes and resistance-associated mobile genetic elements, which can transfer between staphylococcal species. Among other CoNS, *S. epidermidis*, *S. hominis* and *S. haemolyticus* are often reported to be resistant to multiple antibiotics [3, 4].

The *mecA* gene responsible for methicillin resistance was first determined in *S. aureus*, however, many other staphylococcal species were found to also harbour it [5]. The *mecA* gene encodes an additional penicillin-binding protein 2a (PBP 2a), which mediates cell wall synthesis in the presence of  $\beta$ -lactam antibiotics [6]. Together with its regulators *mecI-mecR1* and site specific recombination genes *ccrA* and *ccrB*, the *mecA* gene, is located on a mobile genetic element known as staphylococcal cassette chromosome *mec* (SCC*mec*) [7]. A number of studies have demonstrated the transfer of *mecA* gene from coagulase-negative staphylococcal species to *S. aureus* in vivo, and thus contributing to more successful *S. aureus* clones [8]. To date 11 SCC*mec* types have been reported based on combinations of *mec* (A, B, C1, C2 and D) and *ccr* (AB1, AB2, AB3, AB4 and *ccrC*) complexes and so called J regions (1, 2, 3) [9].

Traditionally recognised as hospital associated pathogens, methicillin resistant coagulase negative staphylococci (MR-CoNS) have recently been linked with a range of ecological niches (community, wildlife and environmental sources) [10–12]. As a result, today increasing attention is being paid to the rapid spread of MR-CoNS and their role in transmission within the community and non - hospital settings [13].

In this study we demonstrate the dissemination of antibiotic resistance in CoNS isolated from various environmental sites in London, UK. The characterization of *mecA* gene and the SCC*mec* elements provide insights into the diversity of environmental CoNS clones.

## Methods

### Isolation

Four hundred seventy-nine samples were collected from different environmental sites in various locations ( $n = 355$ ) and from the hands of volunteers ( $n = 124$ ) in London UK between April 2013 and Nov 2014. Environmental sites included hotels ( $n = 100$ ), baby care facilities ( $n = 65$ ), handbags ( $n = 43$ ), supermarkets ( $n = 37$ ), restaurants ( $n = 36$ ), public transport ( $n = 54$ ), and a public library ( $n = 20$ ). All specimens were plated on Mannitol Salt Agar (Oxoid, Basingstoke, UK), and then incubated aerobically at 37 °C for 24–72 h. One or two colonies for each site were

selected based on staphylococci morphology [4]. The colonies were then purified on Nutrient Agar (Oxoid, Basingstoke, UK).

### Identification

All isolates were initially screened using Gram staining, catalase and coagulase tests. Those that demonstrated potential staphylococci characteristics were identified by Matrix-assisted laser desorption ionization time flight mass-spectroscopy (MALDI-TOF-MS, Microflex LT, Bruker Daltonics, Coventry, UK) in a positive linear mode (2000–20,000 m/z range) as described previously [12]. The resulting spectra were compared with reference spectra by using the Biotyper 3.0 software (Bruker Daltonics, Coventry, UK). *Escherichia. coli* DH5 $\alpha$  (Bruker Daltonics, Coventry, UK) was used as a standard for calibration and quality control.

### Antimicrobial susceptibility test

A panel of 11 antibiotics was used to determine the antibiotic susceptibility of all the isolates. The standard disk diffusion method was used to test AM: amoxicillin (10  $\mu$ g); CEP: cefepime (30  $\mu$ g); CHL: chloramphenicol (30  $\mu$ g); ERY: erythromycin (5  $\mu$ g); FC: fusidic acid (10  $\mu$ g); GEN: gentamicin (10  $\mu$ g); MUP: mupirocin (20  $\mu$ g); OX: oxacillin (1  $\mu$ g); PEN: penicillin (1 unit); STR: streptomycin (10  $\mu$ g); TET: tetracycline (10  $\mu$ g). The susceptible, intermediate resistant or resistant were determined by the Guidelines for Susceptibility Testing [14]. The Minimum Inhibitory Concentrations (MIC) for oxacillin were additionally evaluated using “M.I.C. evaluators” (Oxoid Ltd., Basingstoke, UK).

### Detection of *mecA* gene and staphylococcal cassette chromosome *mec* (SCC*mec*) typing

The *mecA* gene was determined by using PCR method as described previously [15]. For *mecA* positive isolates, SCC*mec* types were determined by evaluating *mec* and *ccr* complexes [15].

### MLST typing of *Staphylococcus epidermidis*

Multi-locus sequence typing (MLST) was used to determine the sequence types of *S. epidermidis* [16]. Sequence types were assigned using the *S. epidermidis* database ([www.mlst.net](http://www.mlst.net)).

## Results

### Purification of isolates

A total of 643 staphylococci isolates were recovered in this study, including those from hotels ( $n = 74$ ), baby care facilities ( $n = 46$ ), handbags ( $n = 17$ ), supermarkets ( $n = 89$ ), restaurants ( $n = 96$ ), public transportation ( $n = 94$ ), human hands ( $n = 192$ ) and public libraries ( $n = 35$ ) (Additional file 1: Table S1).

### Species determination

Six hundred forty-three staphylococcal isolates belonging to 19 staphylococcal species were identified in this study. This included: *S. epidermidis* (n = 193), *S. hominis* (n = 161), *S. capitis* (n = 77), *S. warneri* (n = 63), *S. haemolyticus* (n = 45), *S. pasteurii* (n = 33), *S. saprophyticus* (n = 20), *S. aureus* (n = 12), *S. simiae* (n = 10), *S. cohnii* (n = 9), *S. sciuri* (n = 5), *S. pettenkoferi* (n = 3), *S. auricularis* (n = 2), *S. caprae* (n = 2), *S. equorum* (n = 2), *S. lugdunensis* (n = 2), *S. xylosum* (n = 2), *S. arlettae* (n = 1), and *S. simulans* (n = 1). *S. epidermidis* was the predominant species, followed by *S. hominis*, *S. capitis*, *S. warneri*, *S. haemolyticus*, *S. pasteurii*, and *S. saprophyticus*. However, the occurrence of the species varied for different sites. *S. epidermidis* was predominant among the isolates recovered from restaurants, public transport, hands and handbags, whereas *S. hominis* was predominant among the isolates recovered from supermarkets, baby care facilities and hotels and *S. haemolyticus* was predominantly isolated from the library (Table 1).

### Antibiotic susceptibility test results

The disc diffusion method was used to test 606 isolates against a panel of 11 antibiotics. 572 (94%) isolates were resistant to at least one antibiotic, and only 34 isolates were fully susceptible. Resistance to penicillin, and fusidic acid was observed in more than 65% of all staphylococcal isolates tested. 202 (33%) isolates were resistant to streptomycin, 190 (31%) to erythromycin, 161 (27%) to amoxicillin, 98 (16%) to tetracycline, 87 (14%) to mupirocin, 59 (10%) to gentamicin, 48 (8%) cefepime, 36 (6%) oxacillin, and 21(3%) chloramphenicol (Table 2).

### mecA gene determination and SCCmec typing results

Sixty-eight (11%) *mecA* positive staphylococcal isolates were determined, however, no MRSA was determined in this study. *S. sciuri* had the highest *mecA* gene carriage

(80%) among all 19 staphylococcal species, followed by *S. cohnii* (33%), *S. haemolyticus* (22%), and *S. saprophyticus* (20%). Other isolates demonstrated relatively lower carriage of *mecA* gene, including *S. hominis* (3%), *S. capitis* (8%), *S. epidermidis* (11%), *S. warneri* (11%), *S. pasteurii* (13%). No *mecA* gene was found in the remaining 10 species, including *S. aureus*, *S. simiae*, *S. equorum*, *S. caprae*, *S. xylosum*, *S. auricularis*, *S. simulans*, *S. arlettae*, *S. pettenkoferi*, and *S. lugdunensis*.

SCCmec types were fully determined in forty-six isolates. Twenty-two out of 68 isolates lacked either the *mec* gene complex or the *ccr* gene complex. Thirteen staphylococci (19%) carried SCCmec type V, followed by 8 isolates carrying SCCmec type I (2%), 5 isolates SCCmec type IV (7%), 4 isolates SCCmec type II (6%), 1 isolate SCCmec type III (2%), 1 isolate SCCmec type VI (2%), and 1 isolate SCCmec type VIII (2%). In addition, three isolates harboured a new SCCmec type 1A, which carried combination of class A *mec* complex and *ccr* type 1. Of the ten isolates that were non-typeable, three carried a combination of class A *mec* complex and *ccrC*, six carried a combination of class B *mec* and *ccrC*, and one carried class B *mec* and *ccr* type 3 (Table 3).

### Multi-locus sequence typing of S. epidermidis

MLST was performed to determine the housekeeping genes of 13 oxacillin resistant and *mecA* positive *S. epidermidis*. MLST typing revealed that all *S. epidermidis* strains possess new MLST types. MLST types of *S. epidermidis* isolates with in house numbers of 279, 133, 134, 135, 126, 259, 124, 127, 234, 187, 308, 153 and 191 were respectively assigned as ST599, ST600, ST600, ST600, ST601, ST602, ST602, ST603, ST604, ST605, ST606, ST607 and ST608 (Table 4). Three *S. epidermidis* isolates shared the same sequence types (ST), including *S. epidermidis* 133, 134 and 135 that were isolated from different sites of a library (DSL) possessed ST600 whereas *S. epidermidis* 259, and *S. epidermidis* 124 that had ST602 sequence type were isolated from the human hands (HH) and different sites of hotels (DSH) respectively.

## Discussion

### Environmental staphylococcal species

Although antibiotic resistance is commonly linked to the clinic, recent studies from different ecological niches revealed multidrug resistant bacteria is widespread in the environment [11, 12, 17].

We have previously reported on high levels of antibiotic resistance in staphylococci isolated from different environmental/public settings [11, 12]. In this study we evaluated the dissemination of antibiotic resistant staphylococci recovered from a wide range of environmental settings, and characterised the carriage of the *mecA* gene and the diversity of SCCmec elements in these isolates.

**Table 1** Predominant and common staphylococcal species recovered from the human hands and different environmental sites

Sites	Predominant species (%)	Commonly isolated species (%)
BCF	<i>S. hominis</i> (17%)	<i>S. warneri</i> (17%)
DSH	<i>S. hominis</i> (30%)	<i>S. haemolyticus</i> (18%)
DSL	<i>S. haemolyticus</i> (29%)	<i>S. epidermidis</i> (26%)
DSR	<i>S. epidermidis</i> (38%)	<i>S. hominis</i> (35%)
DSS	<i>S. hominis</i> (44%)	<i>S. epidermidis</i> (29%)
DST	<i>S. epidermidis</i> (35%)	<i>S. capitis</i> (15%)
HB	<i>S. epidermidis</i> (40%)	<i>S. capitis</i> (27%)
HH	<i>S. epidermidis</i> (36%)	<i>S. hominis</i> (23%)

BCF baby care facilities, DSH different sites of hotels, DSL different sites of a library, DSR different sites of restaurants, DSS different sites of supermarkets; DST different sites of transportation facilities, HB handbags, HH human hand

**Table 2** Antibiotic susceptibility profile of staphylococci isolates recovered from general public settings

Isolates	No of isolates	Resistance to a panel of 11 antibiotics (%)										
		OX	PG	MUP	CEF	GM	FC	S	A	E	T	C
<i>S. epidermidis</i>	176	8	72	16	9	7	64	22	26	43	16	2
<i>S. hominis</i>	152	2	68	9	5	7	66	24	17	38	21	3
<i>S. capitis</i>	73	4	58	15	4	1	60	47	23	14	10	5
<i>S. haemolyticus</i>	40	10	50	13	23	15	68	73	45	0	25	10
<i>S. warneri</i>	63	3	54	17	10	22	59	51	40	27	16	2
<i>S. pasteurii</i>	31	6	69	13	6	9	69	25	31	44	17	3
<i>S. saprophyticus</i>	20	15	90	25	5	10	100	10	25	35	15	10
<i>S. aureus</i>	12	0	83	17	0	58	83	33	50	25	0	8
<i>S. simiae</i>	10	0	10	0	0	0	40	0	0	0	0	0
<i>S. cohnii</i>	9	24	67	11	33	0	78	78	22	56	11	0
<i>S. sciuri</i>	5	60	60	80	0	20	80	80	40	0	0	0
<i>S. pettenkoferi</i>	3	0	33	0	0	0	67	67	33	0	0	0
<i>S. lugdunensis</i>	2	0	50	0	0	0	50	0	0	0	0	0
<i>S. equorum</i>	2	0	50	50	0	0	50	50	50	0	50	0
<i>S. caprae</i>	2	0	100	0	100	100	100	50	50	0	0	0
<i>S. xylosus</i>	2	0	100	50	50	50	100	100	0	0	50	0
<i>S. auricularis</i>	2	0	50	0	50	0	50	0	50	0	0	0
<i>S. arlettae</i>	1	0	100	0	0	0	100	100	100	100	0	0
<i>S. simulans</i>	1	0	100	0	0	0	100	0	0	0	0	0

OX oxacillin (1 µg), PG penicillin G (1 unit), MUP mupirocin (20 µg), CEF cefepime (30 µg), GM gentamicin (10 µg), FC fusidic acid (10 µg), S streptomycin (10 µg), A amoxicillin (10 µg), E erythromycin (5 µg), T tetracycline (10 µg), C chloramphenicol (30 µg)

Six hundred and forty-three staphylococci isolates belonging to 19 species, including *S. epidermidis*, *S. hominis*, *S. haemolyticus*, *S. capitis*, *S. warneri*, *S. pasteurii*, *S. saprophyticus*, *S. cohnii*, *S. aureus*, *S. simiae*, *S. sciuri*, *S. pettenkoferi*, *S. lugdunensis*, *S. equorum*, *S. caprae*, *S. xylosus*, *S. auricularis*, *S. simulans*, and *S. arlettae*, were identified in this study. Interestingly, many of the staphylococci species recovered in our study have previously been associated with the community, preserved food, and wildlife [4, 10].

**Antibiotic resistance**

Antibiotic resistance of staphylococci associated with healthcare settings is well documented, however, little is known about the antibiotic resistance in staphylococci isolated from different ecological niches [4]. In this study, the majority of staphylococci were resistant to penicillin (65%) and fusidic acid (66%) (Fig. 1). Despite that 80% of hospital associated CoNS (across Europe) were reported to be resistant to oxacillin [18], only 6% of CoNS were resistant to oxacillin in this study. In addition, the levels of resistance to chloramphenicol (3%), cefepime (8%), gentamicin (10%), mupirocin (14%), tetracycline (16%), and erythromycin (31%) were lower compared to those reported in clinical settings [19–22]. In contrast, the rates of resistance to fusidic acid (66%), amoxicillin (27%) and streptomycin (33%) in environmental staphylococcal isolates were

higher than those reported in clinical staphylococci isolates [21, 23, 24]. It is widely accepted that higher levels of antibiotic resistance in clinical isolates are due to consistent antibiotic exposure [25]. The environment may also contribute to the development of antibiotic resistance in microorganisms due to human/ animal therapeutics, sewage, agriculture and industrial use of antibiotics [26]. Therefore, the wide dissemination of multidrug resistant CoNS in non-healthcare associated environments is a disturbing finding. In our study, 94% of staphylococcal isolates were phenotypically resistant to at least 1 antibiotic, 18% were resistant to five or more antibiotics and only 6% staphylococcal isolates were fully susceptible. The study also revealed that the number of isolates resistant to multiple antibiotics varied between the different isolation sites. The least number of multiple antibiotic resistant CoNS isolates were recovered from the public transport (58%), the highest was isolated from hotels (78%).

**Methicillin-resistant staphylococci**

Methicillin resistant staphylococci pose a major public health threat, and cause severe economic and health consequences [27]. Methicillin resistance is determined by the *mecA* gene, which encodes for penicillin binding protein 2a (PBP2a) that has a low affinity to β-lactam antibiotics [28]. Hussain et al. assessed the correlation between *mecA*

**Table 3** Molecular characterisation and antibiotic resistance of *mecA* gene positive staphylococci

ID	Sites	Species	PG	MUP	CEF	GM	FC	S	A	E	T	C	<i>mecA</i>	<i>mec</i>	<i>ccr</i>	SCC <i>mec</i>	MIC/OX (mg l <sup>-1</sup> )
71	HH	<i>S. capitis</i>	S	S	S	S	S	R	S	S	S	S	+	-	-	I	0.5
100	DSH	<i>S. cohnii</i>	R	R	S	S	S	S	R	R	S	S	+	Class A	5	5A	1
97	BCF	<i>S. cohnii</i>	R	S	R	S	R	R	S	R	S	S	+	Class B	1	I	<b>0.25</b>
279	HH	<i>S. epidermidis</i>	R	S	S	S	R	S	S	R	S	S	+	Class B	2	IV	2
127	DSH	<i>S. epidermidis</i>	R	S	I	S	S	S	R	R	R	S	+	Class C	5	V	2
139	DSR	<i>S. epidermidis</i>	R	R	R	S	R	S	R	R	R	S	+	Class C	5	V	2
191	DSS	<i>S. epidermidis</i>	R	S	S	S	R	S	R	S	S	S	+	Class B	4	VI	2
153	DSH	<i>S. epidermidis</i>	R	S	S	S	R	S	S	S	S	S	+	Class C	5	V	1
187	DSS	<i>S. epidermidis</i>	R	S	S	S	S	S	S	R	S	S	+	Class C	5	V	1
134	DSL	<i>S. epidermidis</i> *	R	S	S	S	R	R	R	R	R	S	+	Class B	1	I	1
259	HH	<i>S. epidermidis</i>	R	S	R	S	R	R	R	R	S	S	+	Class C	5	V	1
135	DSL	<i>S. epidermidis</i> *	R	S	S	S	R	R	R	R	R	S	+	Class B	2	IV	0.5
124	DSH	<i>S. epidermidis</i>	R	S	R	S	R	R	R	S	R	S	+	Class B	2	IV	0.5
133	DSL	<i>S. epidermidis</i> *	R	S	S	S	R	R	R	R	R	S	+	Class B	3	3B	0.5
126	HH	<i>S. epidermidis</i>	R	S	I	S	S	R	R	R	S	S	+	-	-	III	0.5
119	DSH	<i>S. epidermidis</i>	R	S	S	S	S	S	R	I	R	S	+	Class C	5	V	<b>0.12</b>
111	BCF	<i>S. epidermidis</i>	R	S	S	S	S	S	S	S	S	S	+	Class A	2	II	<b>0.12</b>
202	DST	<i>S. epidermidis</i>	S	R	S	S	R	I	S	S	S	S	+	Class B	5	5B	<b>0.12</b>
264	HH	<i>S. epidermidis</i>	S	R	S	S	R	R	S	S	S	S	+	Class B	2	IV	<b>0.06</b>
129	DSL	<i>S. epidermidis</i>	R	S	S	S	R	R	R	R	R	S	+	Class B	1	I	<b>0.03</b>
362	DSL	<i>S. haemolyticus</i>	R	S	I	S	S	R	R	S	S	S	+	Class C	5	V	2
367	DSL	<i>S. haemolyticus</i>	R	S	R	S	S	R	R	S	S	S	+	Class C	5	V	2
355	DSH	<i>S. haemolyticus</i>	R	S	R	R	R	R	R	S	R	S	+	Class C	5	V	2
384	HH	<i>S. haemolyticus</i>	R	R	S	S	R	R	S	S	S	S	+	Class C	5	V	2
322	DSH	<i>S. haemolyticus</i>	R	S	S	S	S	I	R	S	R	S	+	Class A	1	1A	<b>0.25</b>
382	HH	<i>S. haemolyticus</i>	R	S	I	S	R	R	S	R	S	S	+	-	-	II	<b>0.25</b>
323	DSH	<i>S. haemolyticus</i>	S	S	S	R	S	R	R	I	R	S	+	Class A	2	II	<b>0.12</b>
381	HH	<i>S. haemolyticus</i>	S	S	I	S	S	R	S	S	S	S	+	Class B	5	5B	<b>0.12</b>
360	DSH	<i>S. haemolyticus</i>	S	S	S	S	R	S	S	S	S	R	+	Class B	5	5B	<b>0.06</b>
369	DSL	<i>S. haemolyticus</i>	R	S	S	S	R	R	S	R	S	R	+	Class B	1	I	<b>0.03</b>
413	DSH	<i>S. hominis</i>	S	R	S	S	R	R	S	S	S	S	+	Class C	5	V	2
506	DSS	<i>S. hominis</i>	S	S	S	S	R	S	S	S	S	S	+	Class B	1	I	0.5
400	DSH	<i>S. hominis</i>	R	R	S	S	R	S	R	R	R	S	+	Class A	1	1A	<b>0.12</b>
326	DSH	<i>S. hominis</i>	S	S	S	S	S	S	R	I	S	S	+	Class A	1	1A	<b>0.06</b>
589	HB	<i>S. pasteuri</i>	R	S	S	S	R	R	S	R	S	S	+	Class A	5	5A	<b>0.25</b>
592	HH	<i>S. pasteuri</i>	S	R	S	S	R	R	S	S	S	S	+	Class B	5	5B	<b>0.25</b>
627	HH	<i>S. saprophyticus</i>	R	I	S	S	R	S	S	S	R	S	+	Class B	5	5B	0.5
621	DSS	<i>S. saprophyticus</i>	R	R	R	S	R	S	R	S	R	S	+	Class B	2	IV	<b>0.25</b>
630	HH	<i>S. sciuri</i>	R	R	I	S	R	R	S	S	S	S	+	Class A	4	VIII	2
632	DSH	<i>S. sciuri</i>	R	S	I	S	R	R	R	S	S	S	+	Class A	5	5A	1
633	DSH	<i>S. sciuri</i>	R	R	I	R	R	R	R	S	S	S	+	Class B	5	5B	1
629	HH	<i>S. sciuri</i>	S	R	S	S	S	S	S	S	S	S	+	-	-	II	0.25
704	HH	<i>S. warneri</i>	R	S	I	S	R	R	R	R	S	S	+	Class C	5	V	0.5
662	DSH	<i>S. warneri</i>	R	S	S	R	R	R	R	S	R	S	+	Class C	5	V	<b>0.25</b>

**Table 3** Molecular characterisation and antibiotic resistance of *mecA* gene positive staphylococci (Continued)

ID	Sites	Species	PG	MUP	CEF	GM	FC	S	A	E	T	C	<i>mecA</i>	<i>mec</i>	<i>ccr</i>	SCC <i>mec</i>	MIC/OX (mg l <sup>-1</sup> )
694	HH	<i>S. warneri</i>	R	S	S	S	S	S	S	S	R	S	+	-	-	I	<b>0.25</b>
655	BCF	<i>S. warneri</i>	S	R	S	R	R	R	S	I	S	S	+	Class B	1	I	<b>0.12</b>

Note: \* *S. epidermidis* isolates with similar MLST types

R: resistant, S sensitive, I intermediate

BCF baby care facility, DSH different sites of hotels, DSL different sites of a library, DSR different sites of restaurants, DSS different sites of supermarkets, DST different sites of transportation facilities, HB handbags, HH human hands

A amoxicillin (10 µg), CEF cefepime (30 µg), C chloramphenicol (30 µg), E erythromycin (5 µg), FC fusidic acid (10 µg), GM gentamicin (10 µg), MUP mupirocin (20 µg), OX oxacillin (1 µg), PG penicillin G (1 unit), S streptomycin (10 µg), T tetracycline (10 µg)

gene and oxacillin susceptibility breakpoints (0.5 mg l<sup>-1</sup>) of 493 clinical CoNS belonging to and classified into 4 categories [29]. The *mecA* gene positive staphylococci were categorized into groups I and II, and demonstrated that group I (*S. haemolyticus* (83.3%), *S. epidermidis* (61.9%), *S. hominis* (51.8%)) differs from group II (*S. cohnii* (28.5%), *S. warneri* (27.3%), *S. saprophyticus* (9.0%)) by their high levels of *mecA*-carriage [29]. Interestingly, *S. hominis* (38%), *S. haemolyticus* (22%), and *S. epidermidis* (7%) isolated in this study harboured significantly lower levels of the *mecA* gene. Moreover, in this study *S. cohnii* (33%) and *S. saprophyticus* (10%) showed higher *mecA* gene carriage than clinical isolates reported by Hussain, et al. [29], whereas the levels of *mecA* gene carriage in *S. warneri* (6%) were lower than in clinical isolates. No *mecA* gene was detected in staphylococcal species of groups III and IV, which included *S. xyloso*, *S. lugdunensis*, *S. capitis*, *S. simulans*, and *S. schleiferi* [29]. Similarly, in this study *S. lugdunensis*, *S. xyloso* and *S. simulans* were determined to be susceptible to oxacillin and lacked *mecA* gene. However, in contrast to the reports by Hussain, et al. [29] we found that *mecA* gene was present in 8% of *S. capitis* isolates.

Oxacillin susceptible *mecA* gene positive *S. aureus* (OS-MRSA) has been reported worldwide, and the risk of

induced high levels of oxacillin resistance was determined in OS-MRSA [30, 31]. In this study, 68 (46%) staphylococcal isolates were confirmed by PCR to carry the *mecA* gene, however, they were phenotypically susceptible to oxacillin with the MICs (oxacillin) varying from 0.015 to 2 mg l<sup>-1</sup>. This study demonstrates the prevalence of *mecA* positive but oxacillin susceptible CoNS (OS-CoNS) in the environment. Little is known about OS-CoNS isolates recovered from the environment and their epidemiological data are limited. Additional studies are necessary to further our understanding of the prevalence and molecular epidemiology of OS-CoNS in the environment.

#### SCC*mec* elements


SCC*mec* is a mobile genetic element with two essential components: the *mec* gene complex, and the cassette chromosome recombinase (*ccr*) gene complex [32]. The combination of the *mec* gene complex and *ccr* gene complex confers different SCC*mec* types [32]. SCC*mec* type I, II, III are reported to be associated with MRSA recovered from healthcare settings, whereas SCC*mec* type IV and V are mainly associated with the community [32]. Moreover, it has been shown that the size of SCC*mec* types IV and V are smaller than SCC*mec* types I, II and III, thus conferring

**Table 4** MLST types of 13 oxacillin resistant and *mecA* positive *S. epidermidis*

ID	Sites	Species	<i>arcC</i>	<i>aroE</i>	<i>gtr</i>	<i>mutS</i>	<i>pyrR</i>	<i>tpiA</i>	<i>yqiL</i>	MLST types
279	HH	<i>S. epidermidis</i>	57	17	5	5	3	4	31	ST599
133	DSL	<i>S. epidermidis</i>	57	1	2	2	4	1	4	ST600
134	DSL	<i>S. epidermidis</i>	57	1	2	2	4	1	4	ST600
135	DSL	<i>S. epidermidis</i>	57	1	2	2	4	1	4	ST600
126	HH	<i>S. epidermidis</i>	57	25	9	5	6	1	8	ST601
259	HH	<i>S. epidermidis</i>	57	1	2	2	4	1	1	ST602
124	DSH	<i>S. epidermidis</i>	57	1	2	2	4	1	1	ST602
127	DSH	<i>S. epidermidis</i>	57	10	5	5	10	16	21	ST603
234	HB	<i>S. epidermidis</i>	57	1	1	1	2	41	1	ST604
187	DSS	<i>S. epidermidis</i>	57	1	1	2	2	1	1	ST605
308	HH	<i>S. epidermidis</i>	57	1	2	2	4	7	1	ST606
153	DSH	<i>S. epidermidis</i>	57	1	22	2	2	16	1	ST607
191	DSS	<i>S. epidermidis</i>	57	3	5	5	7	14	11	ST608

HH human hands, DSL different sites of a library, DSH different sites of hotels, DSS different sites of supermarkets

MLST Multi-locus sequence typing



increased mobility by their smaller size and contributing the spread of these smaller SCCmec elements [33]. In this study, SCCmec type I, II or III were found in 19% (n = 13) of mecA-positive CoNS, whereas 27% (n = 18) of CoNS were determined to harbour SCCmec type IV or V. SCCmec type VI and VIII were previously identified in Portugal (2006) and Canada (2009) in hospital associated MRSA (HA-MRSA) [33, 34]. In this study, we identified one of each type, however, we did not detect SCCmec types IX.

Becker et al., have previously summarized the community and livestock associated staphylococcal species and their SCCmec types, which included *S. capitis* (I, IA, II, III, IV, IVa, V, non-typeable: (NT)), *S. cohnii* (NT), *S. epidermidis* (I, IIa, IIb, III, III (variant), IV, IVa, IVb, IVc, IVd, IVe, IVg, V, VI, NT), *S. haemolyticus* (I, II, II.1, III, III (variant), IV, V, NT), *S. hominis* (I, III, IV, NT), *S. pasteurii* (IVc), *S. saprophyticus* (III, NT), *S. sciuri* (I, III, IIIA, V, VII, NT) and *S. warneri* (IV, IV.1, IVb, IVE) [4]. In this study, species associated SCCmec types differed and included the following: *S. capitis* (I, NT), *S. haemolyticus* (I, II, V, NT) and *S. hominis* (I, V, NT), *S. cohnii* (I, V, NT), *S. pasteurii* (NT), *S. saprophyticus* (IV, NT), *S. sciuri* (II, VIII), *S. warneri* (I, V, NT). *S. epidermidis* possessed SCCmec types similar to those reported previously [4].

Thirteen unclassified SCCmec types were determined in this study, including three carrying class A mec complex and ccrC, six had a combination of class B mec and ccrC, one carried class B mec and ccr3, and three had a combination of class A mec complex and ccr type 1. The 1A was previously defined as a new SCCmec type 1A by others [35]. Pseudo (ψ)-SCCmec harbours the mec complex but

lacks ccr, while, SCCmec12263 is reported to carry the ccr complex but lacks mec complex [36, 37]. In this study, 21 isolates (29%) were categorized as (ψ)-SCCmec and SCCmec12263 since they lacked either mec complex or ccr complexes. ψ SCC element is characterized by lacking genes for ccr and mec [4]. One of *S. saprophyticus* isolates in this study was found to possess the ψ SCC element (Table 5).

#### MLST of *S. epidermidis*

Whilst many studies have reported on the changing epidemiology of *S. aureus*, epidemiological data of other staphylococcal species are limited [38, 39]. In this study, 10 new MLST types were determined in 13 *S. epidermidis* isolates. Interestingly, although isolates recovered from human hands (*S. epidermidis* 259/ SCCmec V) and hotels (*S. epidermidis* 124/ SCCmec IV) harboured different SCCmec types, they shared the same MLST type ST602. In addition, three *S. epidermidis* isolates recovered from libraries (*S. epidermidis* 133, *S. epidermidis* 134, *S. epidermidis* 135) shared the same MLST type ST600 (Table 4). However, despite sharing the same MLST type *S. epidermidis* 133, *S. epidermidis* 134 and *S. epidermidis* 135 harbored SCCmec type 3B, I, IV respectively. Others reported that *S. epidermidis* ST2 was associated with type II, III, IV and non-typable SCCmec, and *S. epidermidis* ST22 harboured SCCmec type III, IV and V [40].

#### Conclusions

Systematic analysis of staphylococci isolated from non-healthcare environments provided insights into the diversity and antibiotic susceptibility patterns of these

**Table 5** The diversity of SCCmec types of *mecA* gene positive staphylococci

ID	Sites	Species	PG	MUP	CEF	GM	FC	S	A	E	T	C	<i>mecA</i>	<i>mec</i>	<i>ccr</i>	SCCmec	MIC/OX (mg l <sup>-1</sup> )
75	HH	<i>S. capitis</i>	R	S	S	S	R	R	S	S	S	S	+	Class A	NT	Pseudo (ψ)-SCCmec	0.5
81	HH	<i>S. capitis</i>	R	S	R	S	R	R	R	R	S	S	+	NT	5	SCCmec12263	0.5
70	HH	<i>S. capitis</i>	R	S	S	S	S	R	S	S	R	S	+	NT	5	SCCmec12263	<b>0.25</b>
83	HH	<i>S. capitis</i>	S	R	S	S	R	R	S	S	S	S	+	NT	5	SCCmec12263	<b>0.12</b>
24	DSH	<i>S. capitis</i>	S	S	S	S	R	R	S	S	S	S	+	NT	1	SCCmec12263	<b>0.12</b>
108	HH	<i>S. cohnii</i>	S	S	I	S	R	R	S	R	R	S	+	Class A	NT	Pseudo (ψ)-SCCmec	1
308	HH	<i>S. epidermidis</i>	R	R	S	S	R	S	R	R	S	S	+	Class B	NT	Pseudo (ψ)-SCCmec	2
234	HB	<i>S. epidermidis</i>	S	R	S	S	R	R	S	R	R	S	+	Class A	NT	Pseudo (ψ)-SCCmec	1
249	DSH	<i>S. epidermidis</i>	R	S	S	S	R	R	S	S	R	S	+	NT	2	SCCmec12263	<b>0.12</b>
125	DSH	<i>S. epidermidis</i>	S	S	I	S	S	R	S	S	S	S	+	NT	5	SCCmec12263	<b>0.06</b>
185	DSS	<i>S. epidermidis</i>	R	S	S	S	S	S	S	S	S	S	+	Class C	NT	Pseudo (ψ)-SCCmec	<b>0.06</b>
498	DSS	<i>S. hominis</i>	R	S	S	S	R	S	S	R	S	S	+	Class A	NT	Pseudo (ψ)-SCCmec	0.5
426	DSH	<i>S. hominis</i>	R	S	I	S	R	R	R	R	S	S	+	Class A	NT	Pseudo (ψ)-SCCmec	<b>0.25</b>
412	DSH	<i>S. hominis</i>	R	S	S	S	R	S	R	R	S	S	+	NT	1	SCCmec12263	<b>0.06</b>
391	BCF	<i>S. hominis</i>	R	S	S	S	R	S	S	S	S	S	+	NT	5	SCCmec12263	<b>0.03</b>
593	HH	<i>S. pasteurii</i>	R	S	S	R	R	R	R	S	S	S	+	NT	5	SCCmec12263	0.5
597	HH	<i>S. pasteurii</i>	R	R	I	S	S	R	S	S	S	S	+	NT	5	SCCmec12263	0.5
616	BCF	<i>S. saprophyticus</i>	R	R	S	S	R	I	R	R	R	S	+	NT	5	SCCmec12263	256
612	BCF	<i>S. saprophyticus</i>	R	R	S	S	R	S	S	R	S	S	+	NT	NT	ψ SCC	1
659	DSH	<i>S. warneri</i>	R	R	S	S	R	R	R	S	S	S	+	NT	5	SCCmec12263	0.5
648	BCF	<i>S. warneri</i>	R	S	S	R	R	S	R	S	S	S	+	NT	5	SCCmec12263	<b>0.06</b>
645	BCF	<i>S. warneri</i>	R	S	S	S	R	S	S	S	S	S	+	NT	4	SCCmec12263	<b>0.015</b>

R resistant, S sensitive, I intermediate  
 BCF baby care facility, DSH different sites of hotels, DSL different sites of a library, DSR different sites of restaurants, DSS different sites of supermarkets; DST different sites of transportation facilities, HB handbags, HH human hands  
 A amoxicillin (10 µg), CEF cefepime (30 µg), C chloramphenicol (30 µg), E erythromycin (5 µg), FC fusidic acid (10 µg), GM gentamicin (10 µg), MUP mupirocin (20 µg), OX oxacillin (1 µg), PG penicillin G (1 unit), S streptomycin (10 µg), T tetracycline (10 µg)

isolates. Multi-drug resistance was commonly seen in each staphylococcal species. The prevalence of multiple antibiotic resistant staphylococci in this study provides evidence that antibiotics in the natural environments can contribute to the selection of antibiotic resistance in microorganisms. The finding of various SCCmec types in non-healthcare associated environments emphasizes the complexity of SCCmec elements. In addition to this, we also report on new MLST types that were assigned for all *S. epidermidis* isolates. This highlights the genetic variability of these isolates. In conclusion, the non-healthcare environments may act as a reservoir of multidrug resistant staphylococci, and current infection control measures are ineffective in limiting the spread of these bacteria.

**Additional file**

**Additional file 1: Table S1.** Isolates collected from different environmental sites and human hands (PDF 46 kb)

**Abbreviations**

AM: Amoxicillin; BCF: Baby care facility; CEP: Cefepime; CHL: Chloramphenicol; CoNS: Coagulase-negative staphylococci; DSH: Different sites of hotels; DSL: Different sites of a library; DSR: Different sites of restaurants; DSS: Different sites of supermarkets; DST: Different sites of transportation facilities; ERY: Erythromycin; FC: Fusidic acid; GEN: Gentamicin; HB: Handbags; HH: Human hands; MALDI-TOF-MS: Matrix-assisted laser desorption ionization time flight mass-spectroscopy; MIC: Minimum Inhibitory Concentrations; MLST: Multi-locus sequence typing; MR-CoNS: Methicillin resistant coagulase negative staphylococci; MRSA: Methicillin resistant *Staphylococcus aureus*; MUP: Mupirocin; OX: Oxacillin; PEN: Penicillin; SCCmec: Staphylococcal cassette chromosome mec; ST: Sequence types; STR: Streptomycin; TET: Tetracycline

**Funding**

This work was part of Zhen Xu's PhD study funded by China Scholarship Council.

**Availability of data and materials**

All data generated or analysed during this study are included in this published article and its supplementary information files.

**Authors' contributions**

ZX: samples collection, laboratory work, data analysis, manuscript preparation. HS: study design, critically reviewing the paper. RM: Data analysis, critically reviewing the paper. JC: data analysis, critically reviewing the paper. WZ: data analysis, critically reviewing the paper. YL: data analysis,

critically reviewing the paper. RRC: conception and design of the study. HVM: conception and design of the study; data analysis; writing and critically reviewing the paper. All authors read and approved the final manuscript.

#### Ethics approval and consent to participate

Not applicable.

#### Consent for publication

Not applicable.

#### Competing interests

The authors declare that they have no competing interests.

#### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

#### Author details

<sup>1</sup>National Demonstration Center for Experimental Preventive Medicine Education, Tianjin Medical University, Qixiang Road No. 22, Tianjin 300070, China. <sup>2</sup>School of Biological and Chemical Sciences, Queen Mary University of London, London, UK. <sup>3</sup>Department of Natural Sciences, Middlesex University, The Burroughs Hendon, Middlesex NW4 4BT, UK. <sup>4</sup>Natural History Museum, Cromwell Rd, London SW7 5BD, UK. <sup>5</sup>Department of Infectious Diseases, Huashan Hospital, Fudan University, Shanghai 200040, China. <sup>6</sup>Tianjin Qiqing hospital, Tianjin 300380, China. <sup>7</sup>School of Health, Sport and Biosciences, University of East London, E1 4NS, London, UK.

Received: 19 April 2018 Accepted: 6 June 2018

Published online: 13 June 2018

#### References

- Kloos WE, Bannerman TL. Update on clinical significance of coagulase-negative staphylococci. *Clin Microbiol Rev.* 1994;7(1):117–40.
- Foster TJ, Geoghegan JA, Ganesh VK, Höök M. Adhesion, invasion and evasion: the many functions of the surface proteins of *Staphylococcus aureus*. *Nat Rev Microbiol.* 2014;12(1):49–62.
- Bouchami O, Achour W, Mekni MA, Rolo J, Ben HA. Antibiotic resistance and molecular characterization of clinical isolates of methicillin-resistant coagulase-negative staphylococci isolated from bacteremic patients in oncohematology. *Folia Microbiol.* 2011;56(2):122–30.
- Becker K, Heilmann C, Peters G. Coagulase-negative staphylococci. *Clin Microbiol Rev.* 2014;27(4):870–926.
- Ubukata K, Nonoguchi R, Song MD, Matsuhashi M, Konno M. Homology of *mecA* gene in methicillin-resistant *Staphylococcus haemolyticus* and *Staphylococcus simulans* to that of *Staphylococcus aureus*. *Antimicrob Agents Chemother.* 1990;34(1):170–2.
- Pinho MG, de Lencastre H, Tomasz A. An acquired and a native penicillin-binding protein cooperate in building the cell wall of drug-resistant staphylococci. *Proc Natl Acad Sci.* 2001;98(19):10886–91.
- Hiramatsu K, Cui L, Kuroda M, Ito T. The emergence and evolution of methicillin-resistant *Staphylococcus aureus*. *Trends Microbiol.* 2001;9(10):486–93.
- Harrison EM, Paterson GK, Holden MTG, Ba X, Rolo J, Morgan FJE, et al. A novel hybrid SCC*mec*-*mecC* region in *Staphylococcus sciuri*. *J Antimicrob Chemother.* 2014;69(4):911–8.
- IWG-SCC. Classification of staphylococcal cassette chromosome *mec* (SCC*mec*): guidelines for reporting novel SCC*mec* elements. *Antimicrob Agents Chemother.* 2009;53(12):4961–7.
- Pantucek R. *Staphylococcus simiae* sp. nov., isolated from south American squirrel monkeys. *Int J Syst Evol Microbiol.* 2005;55(5):1953–8.
- Xu Z, Mkrtchyan HV, Cutler RR. Antibiotic resistance and *mecA* characterization of coagulase-negative staphylococci isolated from three hotels in London, UK. *Front Microbiol.* 2015;6:947.
- Mkrtchyan HV, Russell CA, Wang N, Cutler RR. Could public restrooms be an environment for bacterial resistomes? *PLoS One.* 2013;8(1):e54223.
- Boyce JM. Environmental contamination makes an important contribution to hospital infection. *J Hosp Infect.* 2007;65(Suppl 2):50–4.
- Andrews JM, Howe RA. BSAC standardized disc susceptibility testing method (version 10). *J Antimicrob Chemother.* 2011;66:2726–57.
- Kondo Y, Ito T, Ma XX, Watanabe S, Kreiswirth BN, Etienne J, et al. Combination of multiplex PCRs for staphylococcal cassette chromosome *mec* type assignment: rapid identification system for *mec*, *ccr*, and major differences in junkyard regions. *Antimicrob Agents Chemother.* 2007;51(1):264–74.
- Thomas JC, Vargas MR, Miragaia M, Peacock SJ, Archer GL, Enright MC. Improved multilocus sequence typing scheme for *Staphylococcus epidermidis*. *J Clin Microbiol.* 2007;45(2):616–9.
- Leonard FC, Markey BK. Methicillin-resistant *Staphylococcus aureus* in animals: a review. *Vet J.* 2008;175(1):27–36.
- Hanberger H, Diekema D, Fluit A, Jones R, Struelens M, Spencer R, et al. Surveillance of antibiotic resistance in European ICUs. *J Hosp Infect.* 2001;48(3):161–76.
- Agvald-Ohman C, Lund B, Edlund C. Multiresistant coagulase-negative staphylococci disseminate frequently between intubated patients in a multidisciplinary intensive care unit. *Crit Care.* 2004;8(1):R42–7.
- Mohan U, Jindal N, Aggarwal P. Species distribution and antibiotic sensitivity pattern of coagulase negative staphylococci isolated from various clinical specimens. *Indian J Med Microbiol.* 2002;20(1):45–6.
- Akinkunmi E, Lamikanra A. Species Distribution and Antibiotic resistance in coagulase-negative staphylococci colonizing the gastrointestinal tract of children in Ile-Ife, Nigeria. *Trop J Pharm Res.* 2010;9(1):35–43.
- Fritsche TR, Sader HS, Jones RN. Comparative activity and spectrum of broad-spectrum  $\beta$ -lactams (cefepime, ceftazidime, ceftioxone, piperacillin/tazobactam) tested against 12,295 staphylococci and streptococci: report from the SENTRY antimicrobial surveillance program (North America: 2001–2002). *Diagn Microbiol Infect Dis.* 2003;47(2):435–40.
- Ferreira RBR, Nunes APF, Kokis VM, Krepsky N, de Fonseca LS, de Bastos Mdo CF, et al. Simultaneous detection of the *mecA* and *ileS-2* genes in coagulase-negative staphylococci isolated from Brazilian hospitals by multiplex PCR. *Diagn Microbiol Infect Dis.* 2002;42(3):205–12.
- Idriss SHE, Foltys V, Tančin V, Kirchnerová K, Tančinová D, Zaujec K. Mastitis pathogens and their resistance against antimicrobial agents in dairy cows in Nitra, Slovakia. *Slovak J Anim Sci.* 2014;47(1):33–8.
- Antoniadou A, Kanellakopoulou K, Kanellopoulou M, Polemis M, Koratzanis G, Papademetriou E, et al. Impact of a hospital-wide antibiotic restriction policy program on the resistance rates of nosocomial gram-negative bacteria. *Infect Dis (Auckl).* 2013;45(6):438–45.
- Cantas L, Shah SQA, Cavaco LM, Manaia CM, Walsh F, Popowska M, et al. A brief multi-disciplinary review on antimicrobial resistance in medicine and its linkage to the global environmental microbiota. *Front Microbiol.* 2013;4(10):96.
- Stefani S, Varaldo PE. Epidemiology of methicillin-resistant staphylococci in Europe. *Clin Microbiol Infect.* 2003;9(12):1179–86.
- Tulinski P, Fluit AC, Wagenaar JA, Mevius D, van de Vijver L, Duim B. Methicillin-resistant coagulase-negative staphylococci on pig farms as a reservoir of heterogeneous staphylococcal cassette chromosome *mec* elements. *Appl Environ Microbiol.* 2012;78(2):299–304.
- Hussain Z, Stoakes L, Massey V, Diagne D, Fitzgerald V, El Sayed S, et al. Correlation of oxacillin MIC with *mecA* gene carriage in coagulase-negative staphylococci. *J Clin Microbiol.* 2000;38(2):752–4.
- Hosokawa Y, Hanaki H, Endo H, Suzuki Y, Nagasawa Z, Otsuka Y, et al. Characterization of oxacillin-susceptible *mecA*-positive *Staphylococcus aureus*: a new type of MRSA. *J Infect Chemother.* 2007;13(2):79–86.
- Saeed K, Dryden M, Parnaby R. Oxacillin-susceptible MRSA, the emerging MRSA clone in the UK? *J Hosp Infect.* 2010;76(3):267–8.
- Monecke K, Coombs G, Shore AC, Coleman DC, Akpaka P, Borg M, et al. A field guide to pandemic, epidemic and sporadic clones of methicillin-resistant *Staphylococcus aureus*. *PLoS One.* 2011;6(4):e17936.
- Oliveira DC, Tomasz ALH. Secrets of success of a human pathogen: molecular evolution of pandemic clones of methicillin-resistant *Staphylococcus aureus*. *Lancet Infect Dis.* 2002;2(3):180–9.
- Zhang K, McClure J-A, Elsayed S, Conly JM. Novel staphylococcal cassette chromosome *mec* type, tentatively designated type VIII, harboring class A *mec* and type 4 *ccr* gene complexes in a Canadian epidemic strain of methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother.* 2009;53(2):531–40.
- Bouchami O, Ben Hassen A, de Lencastre H, Miragaia M. Molecular epidemiology of methicillin-resistant *Staphylococcus hominis* (MRSHo): low Clonality and reservoirs of SCC*mec* structural elements. *Chaturvedi V, editor PLoS One.* 2011;6(7):e21940.
- Harrison EM, Paterson GK, Holden MTG, Morgan FJE, Larsen AR, Petersen A, et al. A *Staphylococcus xylosus* isolate with a new *mecC* allotype. *Antimicrob Agents Chemother.* 2013;57(3):1524–8.

37. Katayama Y, Takeuchi F, Ito T, Ma XX, Ui-Mizutani Y, Kobayashi I, et al. Identification in methicillin-susceptible *Staphylococcus hominis* of an active primordial mobile genetic element for the staphylococcal cassette chromosome *mec* of methicillin-resistant *Staphylococcus aureus*. *J Bacteriol.* 2003;185(9):2711–22.
38. Herwaldt LA, Geiss M, Kao C, Pfaller MA. The positive predictive value of isolating coagulase-negative staphylococci from blood cultures. *Clin Infect Dis.* 1996;22(1):14–20.
39. Wang XM, Noble L, Kreiswirth BN, Eisner W, McClements W, Jansen KU, et al. Evaluation of a multilocus sequence typing system for *Staphylococcus epidermidis*. *J Med Microbiol.* 2003;52(11):989–98.
40. Miragaia M, Thomas JC, Couto I, Enright MC, de Lencastre H. Inferring a population structure for *Staphylococcus epidermidis* from multilocus sequence typing data. *J Bacteriol.* 2007;189(6):2540–52.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

