



## **UWL REPOSITORY**

**repository.uwl.ac.uk**

Microbiological hazards associated with food products imported from the Asia-Pacific region based on analysis of the rapid alert system for food and feed (RASFF) notifications

Dada, Ayokunle Christopher, Somorin, Yinka M., Ateba, Collins N., Onyeaka, Helen, Anyogu, Amarachukwu ORCID logoORCID: <https://orcid.org/0000-0001-9652-7728>, Kasan, Nor Azman and Odeyemi, Olumide A. (2021) Microbiological hazards associated with food products imported from the Asia-Pacific region based on analysis of the rapid alert system for food and feed (RASFF) notifications. *Food Control*, 129. p. 108243. ISSN 0956-7135

<http://dx.doi.org/10.1016/j.foodcont.2021.108243>

This is the Accepted Version of the final output.

**UWL repository link:** <https://repository.uwl.ac.uk/id/eprint/8240/>

**Alternative formats:** If you require this document in an alternative format, please contact: [open.research@uwl.ac.uk](mailto:open.research@uwl.ac.uk)

**Copyright:** Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy:** If you believe that this document breaches copyright, please contact us at [open.research@uwl.ac.uk](mailto:open.research@uwl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

**Rights Retention Statement:**

# Food Control

## Microbiological hazards associated with food products imported from the Asia-Pacific region based on analysis of the Rapid Alert System for Food and Feed (RASFF) notifications --Manuscript Draft--

<b>Manuscript Number:</b>	FOODCONT-D-21-00745R1
<b>Article Type:</b>	Research Paper
<b>Keywords:</b>	Microbiological safety; Supply chain; Foodborne pathogens; food products
<b>Corresponding Author:</b>	Olumide Adedokun Odeyemi Mowbray, AUSTRALIA
<b>First Author:</b>	Olumide A Odeyemi
<b>Order of Authors:</b>	Olumide A Odeyemi Ayokunle C. Dada Yinka M. Somorin Collins N. Ateba Helen Onyeaka Amarachukwu Anyogu Nor Azman Kasan
<b>Abstract:</b>	<p>The Rapid Alert System for Food and Feeds (RASFF) is a monitoring and notification tool developed by the European Commission to serve as a central database for collating food safety-related information for member states of the European Union (EU). This study evaluated the microbiological safety of foods originating from the Asia-Pacific region in the past two decades (2000-2020) by analysing incidences and trends of notifications and alerts on the RASFF database. The highest number of notifications were for foods exported from India and Thailand as foods exported from these two countries constituted more than half (54%) of the notifications recorded on the RASFF database for the entire Asia-Pacific region from 2000-2020, compared to ANZ (Australia and New Zealand), which had very low notifications (1.2% = 23/1873). Among the 2121 notifications of pathogenic microorganisms, consisting of 14 genera, <i>Salmonella</i> was the most predominant as approximately 7 out of every 10 pathogens isolated from exported foods from the Asia-Pacific Region in the past two decades (74%, 1560/2121) were <i>Salmonella</i> species. More than 95% of pathogen species notifications for fruits and vegetables exported from India and Bangladesh were associated with betel leaves. Among the nuts, nut products and seeds, sesame seeds were the main food item contaminated by <i>Salmonella</i>, and these accounted for 87% of total <i>Salmonella</i> notifications. Across the food categories, there was a statistically significant reduction in the number of notifications associated with seafoods over the years (<math>r = -0.73</math>, <math>p=0.0001</math>). Although a statistically significant increase in notifications for fruits and vegetables (<math>r = 0.66</math>, <math>p=0.008</math>) was recorded between 2000-2014, more recent years (2015-2020) have been associated with reducing trends in the number of notifications associated with this food type (<math>r = -0.85</math>, <math>p=0.03</math>). Results indicate that imported foods may be potential vehicles for the transmission of clinically relevant microorganisms.</p>

Office of Research Services,  
University of Tasmania,  
Australia  
19<sup>th</sup> April 2021

Editor,  
Food Control  
Dear Editor,

**RE: Response to Review Comments [Ref. No.: FOODCONT-D-21-00745]**

Thank you for the review of our manuscript " Microbiological hazards associated with food products exported from the Asia-Pacific region based on analysis of the Rapid Alert System for Food and Feed (RASFF) notifications [Ref. No.: FOODCONT-D-21-00745]".

All the editor's and reviewer's comments have now been addressed. Kindly find tabulated below point-by-point responses to each comment. Locations in the journals where changes have been effected are also stated, with changes in the manuscript set as tracked change, as per journal specification.

We thank you in advance for your time in considering this work for publication and look forward to hearing your response.

Yours sincerely,

Dr Olumide Odeyemi

## Highlights

- RASFF pathogen notifications analysed for Asia Pacific food imports (2000 – 2020).
- 1873 notifications reported over 2 decades across 29 notifying countries.
- Imported foods can potentially transmit clinically relevant microorganisms.
- *Salmonella* was the predominant pathogen (7 out of every 10 pathogens).
- Inconsistent actions in responses to notifications by different EU member states

**Response to Reviewers' comments:**

<b>Reviewer #1</b>		
<b>No</b>	<b>Comment</b>	<b>Response</b>
1	In this study, the authors evaluated the microbiological safety of foods being imported from the Asia-Pacific region from 2000 to 2020 by analyzing incidences and trends of notifications and alerts in the Rapid Alert System for Food and Feeds database. Overall, this is well-organized research article with some interesting findings.	Thank you
2	The scientific names of pathogens in latin should be in italic, no matter they are in the manuscript or the supplementary materials.	In agreement with the reviewer, issues with the pathogen names in the manuscript relate to the supplementary tables. We have now gone through each of the supplementary tables and have corrected the anomalies in the description of pathogen names. The names are now consistent with acceptable norms in literature (e.g. Jiang et al 2021, Food Control, <a href="https://doi.org/10.1016/j.foodcont.2020.107535">https://doi.org/10.1016/j.foodcont.2020.107535</a> and Bangtrakulnonth et al 2004, Emerging infectious diseases <a href="https://doi.org/10.3201/eid1001.02-0781">https://doi.org/10.3201/eid1001.02-0781</a> ), e.g. <i>Salmonella enterica</i> ser. Derby, <i>Salmonella</i> Typhimurium etc
3	It is confusing both "export" and "import" were used. It is more proper to use "import" since this study was conducted from the perspective of who imported foods from the Asia-Pacific region.	In agreement with the reviewer, issues with import/export clarifications have now been made. Given these changes, the manuscript now reflects that it was conducted from the perspective of who imported foods from the Asia-Pacific region. These changes are reflected in Lines 1,34-42,189,201,226,240, 308,369, 381,383,395,404, 412-414, 447-456, 482-484, 496-502,553,566, 583, 641, 667 and 714
<b>Reviewer #2</b>		
5	The MS entitled "Microbiological hazards associated with food products exported from the Asia-Pacific region based on analysis of the Rapid Alert System for Food and Feed (RASFF) notifications" describes microbial hazards associated with exported food items. The subject of the MS is relevant and this type of scientific investigation helps countries to implement policies to improve export quality. Authors have collected data for last 20 years which is significant amount of data analysis.	Thank you

6	Authors may want to break this data into 5 years slots such as 2000-2004, 2005-2009, and so on to see if there is any improvement in number of notifications reported. Alternatively, authors may want to consider analyzing this data over time to see any if number of notifications have gone up or down over 20 years.	Actually, this has already been done in the manuscript and results presented in Table 3. The data was split into distant years (2000-2014) versus most recent years (2015-2020) and assessed for trends in the different data timeframes (i.e. does the pathogen notification increase with years considering the entire period (2000-2020), most recent years (2015-2020) or distant years (2000-2014). However, we agree with the reviewer that this statement does not come out clearly in the methodology section so we have now added this into the methodology section.
<b>Reviewer #3</b>		
7	Table 2, it would be better to provide information on the total number of foods imported or sample for detection from each country or region.	It is difficult to get data on total number of foods imported or sample for detection from each country or region. One limitation of the RASSF database is that it does not record the number of detection per given number of food items tested. We have now included this as a limitation in the study and suggestion for RASSF database operators to consider including in the future (Line 713-716)
8	In the results part, brief analysis of observed results should be provided. More specific analysis should be conducted in the Discussion. For example, the author mentioned that "The highest number of notifications was made by the UK (28%)", why?.	Analysis and discussion of the results is already presented in the discussion section and would be repetitive if these analysis are presented again in the results section. However, as the reviewer as advised, we have now included more specific analysis....in the discussion section e.g. Line 519-527, 499-503, 594-597)
9	The significance of the research should be provided in both abstract and discussion section	This has now been added into the abstract (line 52-53...) and discussion (line 488-490).
<b>Editor</b>		
11	The highlights do not conform to the journal's requirements: maximum 85 characters, including spaces, per bullet point. Please revise.	The highlights have now been edited to conform with the journal's standard.

1 **Microbiological hazards associated with food products ~~imported~~~~exported~~ from the**  
2 **Asia-Pacific region based on analysis of the Rapid Alert System for Food and Feed**  
3 **(RASFF) notifications**

4  
5  
6 Ayokunle C. Dada\*<sup>1</sup>, Yinka M. Somorin<sup>2</sup>, Collins N. Ateba<sup>3</sup>, Helen Onyeaka<sup>4</sup>, Amarachukwu  
7 Anyogu<sup>5</sup>, Nor Azman Kasan<sup>6</sup>  
8 Olumide A. Odeyemi\*<sup>7</sup>  
9

10 <sup>1</sup>*QMRA Data Experts, P.O. Box 37, Waikato Mail Lobby, Hamilton 3204, New*  
11 *Zealand*

12 <sup>2</sup>*School of Pharmacy, Queen's University Belfast, Belfast, United Kingdom*

13 <sup>3</sup>*Food Security and Safety Niche Area, Faculty of Natural and Agricultural Sciences, North-*  
14 *West University, Mmabatho, South Africa*

15 <sup>4</sup>*Department of Chemical Engineering, University of Birmingham, Edgbaston, United*  
16 *Kingdom*

17 <sup>5</sup>*Food, Nutrition and Public Health, School of Life Sciences, University of Westminster,*  
18 *London, United Kingdom*

19 <sup>6</sup>*Higher Institution Centre of Excellence (HiCoE) Institute of Tropical Aquaculture and*  
20 *Fisheries, Universiti Malaysia Terengganu, 21030 Kuala Nerus, Terengganu, Malaysia.*

21 <sup>7</sup>*Office of Research Services, Research Division, University of Tasmania, Launceston,*  
22 *Australia*

23  
24 \*Corresponding ~~Author~~~~authors~~: [chris@qmradata.com](mailto:chris@qmradata.com), [olumide.odeyemi@utas.edu.au](mailto:olumide.odeyemi@utas.edu.au)  
25  
26  
27

28 **Abstract**

29 The Rapid Alert System for Food and Feeds (RASFF) is a monitoring and notification tool and  
30 food safety-related database developed by the European Commission ~~to serve as a central~~  
31 ~~database for collating food safety related information for member states of the European~~  
32 ~~Union (EU)~~. This study evaluated the microbiological safety of foods originating from the Asia-  
33 Pacific region in the past two decades (2000-2020) by analysing incidences and trends of  
34 notifications and alerts on the RASFF database. The highest number of notifications were for  
35 foods ~~imported~~~~exported~~ from India and Thailand as foods ~~imported~~~~exported~~ from these two  
36 countries constituted more than half (54%) of the notifications recorded on the RASFF  
37 database for the entire Asia-Pacific region from 2000-2020, compared to ANZ (Australia and  
38 New Zealand), which had very low notifications (1.2% = 23/1873). Among the 2121  
39 notifications of pathogenic microorganisms, consisting of 14 genera, *Salmonella* was the most  
40 predominant as approximately 7 out of every 10 pathogens isolated from ~~exported~~ foods  
41 imported from the Asia-Pacific Region in the past two decades (74%, 1560/2121) were  
42 *Salmonella* species. More than 95% of pathogen species notifications for fruits and vegetables  
43 ~~imported~~~~exported~~ from India and Bangladesh were associated with betel leaves. Among the  
44 nuts, nut products and seeds, sesame seeds were the main food item contaminated by  
45 *Salmonella*, and these accounted for 87% of total *Salmonella* notifications. Across the food  
46 categories, there was a statistically significant reduction in the number of notifications  
47 associated with seafoods over the years ( $r = -0.73$ ,  $p=0.0001$ ). Although a statistically  
48 significant increase in notifications for fruits and vegetables ( $r = 0.66$ ,  $p=0.008$ ) was recorded  
49 between 2000-2014, more recent years (2015-2020) have been associated with reducing  
50 trends in the number of notifications associated with this food type ( $r = -0.85$ ,  $p=0.03$ ). Results  
51 indicate that imported foods may be potential vehicles for the transmission of clinically relevant  
52 microorganisms. Studies of this nature can potentially encourage -countries to implement  
53 policies that improve export quality.

54

55 **Keywords:** Microbiological safety, supply chain, foodborne pathogens, food products

56

57

58

59

60

61 **1. Introduction**

62  
63 Food safety is everyone's business, and it is pivotal to public health protection. Many factors  
64 make food unsafe for human and animal consumption, ranging from chemical contaminants  
65 such as pesticides, and mercury in seafood, to biological hazards, including parasites and  
66 microorganisms (2010). It is imperative to maintain consumers' health by ensuring optimum  
67 standards for food from production/harvest through the food supply chain up to the point of  
68 consumption. However, the impact of globalization, leading to a long food supply chain  
69 coupled with the rapid distribution of foods across the world, has made it difficult to track the  
70 origin and history of most foods and/or food additives resulting in food spoilage, loss and  
71 waste, as well as foodborne disease outbreaks from the consumption of unsafe food (Rezaei  
72 & Liu, 2017). The major causes of foodborne illnesses include; bacteria (*Salmonella*,  
73 *Campylobacter*, *Listeria*, *Vibrio cholerae*, enterohaemorrhagic *Escherichia coli*), viruses  
74 (*Norovirus*, Hepatitis A), parasites (*Echinococcus spp*, *Taenia solium*, *Ascaris*,  
75 *Cryptosporidium*, *Entamoeba histolytica*, *Giardia*), prions, fungi, and chemical agents  
76 (naturally occurring toxins such as mycotoxins and phycotoxins, persistent organic pollutants  
77 such as dioxins and polychlorinated biphenyls, heavy metals such as lead, mercury and  
78 cadmium) (World Health Organization, 2020a, 2020b). Over the past few years, there has  
79 been a significant increase in the number of documented cases of foodborne diseases  
80 reported by various countries across the globe. Worldwide, the number is estimated to be 600  
81 million cases of foodborne illnesses annually, resulting in 420,000 deaths, with about 30% of  
82 the deaths recorded in children (Food and Agriculture Organization, 2019). In the Asia-Pacific  
83 region, which comprises countries within South, East, Southeast Asia and Oceania, the  
84 number of foodborne illnesses continues to rise, with current estimates pegged at 275 million  
85 cases annually (Food and Agriculture Organization, 2019). The Asia-Pacific region has a  
86 population of about 4.6 billion people, accounting for about 60% of the global population  
87 (Worldometer, 2021). This large, growing population places a lot of demand on water, energy  
88 and food resources. Furthermore, although the region is rapidly becoming one of the largest  
89 economies globally, over 40% of the population live at or below the poverty line (ESCAP,  
90 2020). The high poverty rate coupled with the growing population and other factors, including  
91 poor hygiene, lack of adequate food processing and preservation routines and poor food policy  
92 enforcement, can be considered the main drivers of foodborne disease outbreaks within the  
93 region (Food and Agriculture Organization, 2019).

94  
95 A large proportion of foodborne disease outbreaks are not noticed immediately, traced to the  
96 consumption of a particular food item, or even recognised as a foodborne disease outbreak.  
97 Others go unreported and are not investigated. Furthermore, there are limitations in the  
98 capacity of many countries to undertake food surveillance due to inefficient food regulatory  
99 agencies and unavailability of reference laboratories and analytical tools/expertise to identify  
100 sources of foodborne disease outbreaks and conduct investigations (Wang, et al., 2016). The  
101 Rapid Alert System for Food and Feeds (RASFF), created in 1979, is a monitoring and  
102 notification tool developed by the European Commission to serve as a central database for  
103 collating food safety-related information for member states of the European Union (EU). It  
104 allows the food regulatory bodies of member countries to report and submit information and  
105 notifications about unsafe foods and unauthorised/illegal foods/food additives in circulation  
106 within the market, ensuring the rapid transmission and sharing of information regarding food  
107 safety risks as they occur in real time. This allows actions, including product  
108 recalls/withdrawals, to prevent adverse health and economic consequences to consumers  
109 within the European Union and beyond (European Commission, 2020a). RASFF features a  
110 consumer portal that is interactive and has been available to consumers since 2014. Its  
111 interactive online database can be used for customised searches of food safety-related  
112 information using criteria including Notification, Type, Hazard, Date, Product and Keywords.  
113 In addition to monitoring food safety, RASFF is also used to track food fraud, substitutions,  
114 and other economically motivated adulterations (EMAs), which may not be considered direct  
115 food hazards, but are still of public health and economic significance (Tähkääpää, Majjala,

116 Korkeala, & Nevas, 2015). The system operates on a 24-hour basis, allowing for a seamless  
117 flow of information and ensuring regulatory compliance. Thus, the RASFF serves the very  
118 important function of mitigating risks associated with foods and continuously ensuring public  
119 health safety (Kowalska & Manning, 2020).

120  
121  
122 The RASFF has recently been used in tracing the origin of food contaminations and in  
123 foodborne outbreak investigations such as the *Salmonella* Poona infections reported in France  
124 in 2019 and the multi-country outbreak of *Listeria monocytogenes* in cold-smoked fish  
125 products in five member EU states of Denmark, Estonia, Finland, France and Sweden  
126 (European Commission, 2020b). Similarly, there have been RASFF notifications from  
127 countries within the Asia-Pacific region, including Australia, Hong Kong, Japan and Nepal,  
128 amongst others (European Commission, 2020a). It is imperative that food exporters,  
129 especially those exporting from developing countries to the more developed countries in  
130 Europe, meet the food standards of the destination country. The presence of microbiological  
131 and chemical hazards in foods destined for Europe, when flagged by RASFF, can lead to  
132 import rejection with attendant economic losses (Somorin, Odeyemi, & Ateba, 2021). The  
133 European Food Safety Authority (EFSA) and the European Center for Disease Prevention and  
134 Control (ECDC) have identified *Salmonella enteritidis*, *Campylobacter* spp., shiga-toxin  
135 producing *Escherichia coli*, *Yersinia* spp. and *Listeria monocytogenes* as the major bacteria  
136 species implicated in human cases of foodborne illnesses (European Food Safety Authority,  
137 2021) and are actively monitored for by the RASFF. These species have previously been  
138 identified in foods of Asian-Pacific origin (Food and Agricultural Organisation, 2018;  
139 Kaakoush, Castaño-Rodríguez, Mitchell, & Man, 2015; Rivas, Strydom, Paine, Wang, &  
140 Wright, 2021; Sodagari, Wang, Robertson, Habib, & Sahibzada, 2020; Sugiri, et al., 2014),  
141 highlighting the need for deliberate measures to eliminate these pathogens from foods,  
142 especially those destined for export to Europe.

143  
144 Although systems such as the RASFF are in operation worldwide in both the public and private  
145 sector to assure food safety, there continue to be incidences of consumer foodborne illnesses  
146 that are directly related to lapses in food regulations, inadequate monitoring and enforcement  
147 or the deliberate attempts by Food Business Operators (FBOs) to increase profits through  
148 opportunistic malpractices (Tähkäpää, et al., 2015). Furthermore, the Covid-19 pandemic, with  
149 its disruption of the food supply chain both domestically and internationally, has worsened the  
150 food security risks, especially in the Asia-Pacific region. The pandemic undermined food  
151 accessibility and availability, while the global lockdown of 2020 disrupted food production,  
152 distribution operations, logistics and trading (Kim, Kim, & Park, 2020). Also, the disruption in  
153 the food supply chain and food production and trade led to an increase in food spoilage rates  
154 due to longer wait times, thus increasing the risks of foodborne illnesses and economically  
155 motivated fraud (Kim, et al., 2020). Therefore, it is evident that continued vigilance is required  
156 to assure food safety, which continues to be a challenge globally, especially regarding the  
157 presence of microorganisms in foods. This article evaluates the microbiological safety of foods  
158 originating from the Asia-Pacific region by analysing incidences and trends of notifications and  
159 alerts on the RASFF database.

160

161

162 **2. Materials and Methods**

163

164 **2.1. RASFF database**

165

166 Six sections (Notification, Type, Hazard, Date, Product and Keywords), presented as fill-in  
167 forms, constitute the RASFF portal search page. In the "Notification" section, users can use  
168 four sub-sections to enter details related to a food item for which notification information is  
169 being sought. These sub-sections include 1) 'Reference' – a unique reference number  
170 associated with each notification; 2) 'Subject' – a field wherein specific subjects can be  
171 searched using the "AND","OR" Boolean operators; 3) 'Notified by' – a drop-down list of  
172 notifying countries (EU member states); and 4) 'Open alert' – a field where users can select  
173 for information that includes or excludes open alerts. The "Type" section consists of three  
174 subsections; 1) 'Type'- allows users to select their preference for information related to food,  
175 feed or food contact material; 2) 'Classification'- where users can select the notification  
176 classification; and 3) 'Basis'- where users can select the notification basis for which information  
177 is being sought. The 'Date' section allows users to select dates or periods of the year for which  
178 notification data is required. The 'Product' section consists of four subsections; 'Category' (a  
179 drop-down list of types of food included in the RASFF database); 'Country' (a drop-down list  
180 of all notifying and originating countries); 'action taken' (a drop-down list of action taken  
181 following notification); and 'flagged as' (with two sublevels – distribution and origin). The  
182 'Keyword' section allows users to search by keywords.

183

184 **2.2 Data collection**

185

186 The dataset used for this study was obtained from the RASFF database as described by Yinka  
187 M. Somorin, Olumide A. Odeyemi, and Collins N. Ateba (2021) with slight modifications based  
188 on the three search criteria that focused on the product country of origin, product type and  
189 hazard category. In terms of the product's country of origin, the database was searched for  
190 information relating to all Asia-Pacific countries. Notification data was only available for imports  
191 from 20 ~~exporting~~ countries in the Asia-Pacific region: Afghanistan, Australia, Bangladesh,  
192 Cambodia, China, Hong Kong, India, Indonesia, Laos, Malaysia, Nepal, New Zealand,  
193 Pakistan, Philippines, Singapore, South Korea, Sri Lanka, Taiwan, Thailand and Vietnam. In  
194 terms of product type, the search was restricted to food only, excluding feed and food contact  
195 materials. In terms of hazard category, the focus was placed on "pathogenic microorganisms"  
196 included in the RASFF database. The initial notification date was not restricted to enable  
197 complete capturing of all the available notifications; however, considering the paucity of data  
198 preceding the year 2000, notification data preceding 2000 were subsequently excluded from  
199 the dataset. The dataset used therefore spans two decades.

200

201 **2.2. Data analysis**

202

203 The notifications obtained for the imports from 20 ~~exporting~~ countries in the Asia-Pacific region  
204 were exported as an Excel file. Data sorting, filtering, and formatting were completed using  
205 Microsoft Excel spreadsheets. Missing data instances were replaced with the term "not  
206 specified". Descriptive statistics were collated to determine the predominant pathogens  
207 associated with seafood originating from countries in the Asia-Pacific region (the most  
208 contaminated food category); period of lowest and highest notifications; number of notifying  
209 countries; and the country of origin. To determine trends in the yearly notifications, Pearson  
210 correlations of year versus number of notifications per sub-region and per food category were  
211 conducted using IBM® Statistical Package for the Social Sciences (SPSS®) software version  
212 24. The data was split into distant years (2000-2014) versus most recent years (2015-2020)  
213 and assessed for trends in the different data timeframes (i.e., does the pathogen notification  
214 increase with years considering the entire period (2000-2020), most recent years (2015-2020)  
215 or distant years (2000-2014). A statistically significant increasing trend of notifications was

216 reported when the Pearson correlation statistics (r) were >0.5 with an associated p-value  
217 <0.05. A statistically significant decreasing trend of notifications was reported when the  
218 Pearson correlation statistics (r) were < -0.5 with an associated p-value <0.05. A map of  
219 originating Asia-Pacific countries was processed in Tableau, and Chord diagrams were  
220 generated using Rstudio, a free and open-source environment for the statistical language R  
221 (<https://www.r-project.org>). A Sankey Diagram was developed using Python to display  
222 notification flows and their quantities in proportion to one another, with the originating country  
223 designated as the source node, while the product type and action taken were designated as  
224 target nodes.

225

### 226 3. Results

227

228

#### 229 3.1. Notifications - originating and notifying countries.

230


231 A total of 1873 notifications from 2000 to 2020 were obtained, based on our search criteria,  
232 ~~for food items imported~~ from 20 ~~countries in the exporting countries~~ in the Asia-Pacific region:  
233 Afghanistan, Australia, Bangladesh, Cambodia, China, Hong Kong, India, Indonesia, Laos,  
234 Malaysia, Nepal, New Zealand, Pakistan, Philippines, Singapore, South Korea, Sri Lanka,  
235 Taiwan, Thailand and Vietnam (**Fig.1**).

236

#### 237 3.2 Notification categories

238

239 **Presented in Table 1** are three levels of notification categories associated with the RASFF  
240 database (i.e., Notification basis, Notification type and Action taken). was observed in this  
241 study. In more than half of the notifications (51%), the basis of notification was border control,  
242 with the consignment detained (**Table 1**). Only 1% of notifications were due to food poisoning  
243 and < 1% due to consumer complaints. The notification type was mainly border rejection (43%)  
244 while the most prevalent action taken was re-dispatching (17.6%) and destruction of the  
245 consignments (17.5%, **Table 1**). The highest number of notifications were for foods  
246 ~~imported~~~~exported~~ from India (31% = 588/1873), followed by Thailand (23% = 428/1873, **Table**  
247 **2**). The categories of foods categories originating from Asia-Pacific Region (n=1873) by  
248 country of origin from 2020 – 2020 are shown in **Fig. 2**. Together, foods ~~imported~~~~exported~~  
249 from these two countries constituted more than half (54%) of the notifications recorded on the  
250 RASFF database for the entire Asia-Pacific region from 2000-2020 (**Fig 3**), compared to ANZ  
251 (Australia and New Zealand), which had very low notifications (1.2% = 23/1873). Also, the  
252 prevalent actions taken associated with notifications from India and Thailand were re-  
253 dispatching and destruction of the consignments (**Fig. 4**).



254

255 **Fig. 1:** Map of originating country and number of notifications listed in the *RASFF* database

256 [N.B. Additional countries not on the map: Hong Kong (Notifications: 6, % of total notifications:  
 257 0.32%) and Singapore (Notifications: 3, % of total notifications: 0.16%)]


258

259

260

261

262




263  
 264 **Fig. 2** Food categories of RASFF notifications (n=1873) originating from Asia-Pacific Region  
 265 (n=1873) by country of origin from 2020 - 2020. Meat includes meat and meat products (other  
 266 than poultry), poultry meat and poultry meat products. Seafoods include bivalve molluscs and  
 267 products thereof, cephalopods and products thereof, crustaceans and products thereof, fish  
 268 and fish products. Other foods include prepared dishes and snacks, soups, broths, sauces  
 269 and condiments, cereals and bakery products, cocoa and cocoa preparations, coffee and tea,  
 270 confectionery, eggs and egg products, food additives and flavourings, milk and milk products  
 271 and other food product / mixed dietetic foods, food supplements, fortified foods. ANZ=  
 272 Australia and New Zealand, IDN = Indonesia, MYS=Malaysia. Other\_O = Other originating  
 273 countries, i.e., Pakistan, Philippines, Hong Kong, Taiwan, Sri Lanka, Singapore, South Korea,  
 274 Afghanistan, Chile, Nepal and Cambodia. The length of the arc on the circumference of the  
 275 circle represents the number of notifications in each food category.

276



277  
 278 **Fig. 3.** Types of RASFF notifications (n=1873) originating from Asia-Pacific Region by country of origin from 2020 - 2020. Inform\_ATTN= Information for attention, Inform\_FLP = Information  
 279 for follow-up. ANZ= Australia and New Zealand, IDN = Indonesia, MYS=Malaysia. Other\_O =  
 280 Other originating countries, i.e., Pakistan, Philippines, Hong Kong, Taiwan, Sri Lanka,  
 281 Singapore, South Korea, Afghanistan, Chile, Nepal and Cambodia. The length of the arc on  
 282 the circumference of the circle represents the number of notifications in each food category.  
 283

284  
 285  
 286  
 287  
 288  
 289  
 290



291

292 **Fig. 4.** Sankey diagram of originating country versus food type and action taken.

293 N.B. The width of the arrows or lines show their magnitudes, so the bigger the arrow, the larger  
 294 the flow (i.e., the number of pathogen notifications associated with that source-target node).  
 295 Colours have been used to divide the diagram into different categories in a way that shows  
 296 different source-target node connections.

297

298 **Table 1**

299 RASFF notifications of pathogenic microorganisms in seafood originating from the Asia-  
 300 Pacific Region (n=1873) from 2020 – 2020 based on notification category.

Classification category	Notifications	
	Number	Percentage (%)
<b>Notification basis</b>		
border control - consignment detained	949	50.67%
official control on the market	349	18.63%
border control - consignment released	239	12.76%
not specified	216	11.53%
company's own check	70	3.74%
food poisoning	24	1.28%
border control - consignment under customs	17	0.91%
consumer complaint	8	0.43%
<b>Notification type</b>		
border rejection	810	43.25%
information	509	27.18%
information for attention	283	15.11%
alert	246	13.13%
information for follow-up	25	1.33%
<b>Action taken</b>		
re-dispatch	329	17.57%
destruction	327	17.46%
import not authorised	275	14.68%
official detention	130	6.94%
not specified	104	5.55%
withdrawal from the market	97	5.18%
informing recipient(s)	83	4.43%
informing authorities	76	4.06%
recall from consumers	67	3.58%
product recall or withdrawal	56	2.99%
no action taken	51	2.72%
no stock left	43	2.30%
prohibition to trade - sales ban	37	1.98%
re-dispatch or destruction	29	1.55%
return to consignor	27	1.44%
placed under customs seals	21	1.12%
physical/chemical treatment	21	1.12%
detained by operator	19	1.01%
reinforced checking	16	0.85%
withdrawal from recipient(s)	13	0.69%
public warning - press release	11	0.59%
Other actions	41	2.19%

301 Other actions: Screening sample, Product destruction or return after official permission,  
 302 physical treatment - heat treatment, relabelling, Product recalled / Product to be destroyed,  
 303 destination of the product identified, informing consignor and seizure.

304 **Table 2**

305 RASFF notifications of pathogenic microorganisms in foods originating from the Asia-Pacific Region (n=1873) from 2000 - 2020 based on  
 306 notifying country.

Notifying country	Country of origin										Grand Total
	India	Thailand	Vietnam	Bangladesh	China	Malaysia	Indonesia	Laos	ANZ	Other_O	
United Kingdom	216	80	5	150	28	11	3	16	2	19	<b>528</b>
Italy	51	17	29	1	22	58	27	0	3	16	<b>221</b>
Netherlands	29	92	35	2	10	1	1	6	1	3	<b>179</b>
Norway	27	54	36	19	6	1	9	1	8	15	<b>168</b>
Germany	46	30	39	1	9	3	2	1	3	14	<b>145</b>
Finland	13	72	7	4	10	1	0	0	1	8	<b>115</b>
Spain	15	2	30	0	18	8	1	0	0	0	<b>74</b>
Greece	64	0	2	1	2	0	3	0	0	1	<b>73</b>
Sweden	6	26	9	1	10	0	1	0	1	7	<b>60</b>
Poland	48	0	6	0	2	0	0	0	0	0	<b>56</b>
France	16	5	5	8	5	2	7	0	3	6	<b>54</b>
Denmark	3	27	10	1	8	1	1	0	0	2	<b>53</b>
Portugal	1	0	26	0	0	2	1	0	0	1	<b>31</b>
Belgium	4	8	4	2	3	1	0	0	0	2	<b>24</b>
Other_N	49	15	6	1	7	0	8	0	1	6	<b>92</b>
<b>Grand Total</b>	<b>588</b>	<b>428</b>	<b>249</b>	<b>191</b>	<b>140</b>	<b>89</b>	<b>64</b>	<b>24</b>	<b>23</b>	<b>100</b>	<b>1873</b>

307

308 **Other\_N: Other notifying countries:** Cyprus, Austria, Slovenia, Ireland, Lithuania, Iceland, Romania, Switzerland, Bulgaria, Croatia, Estonia,  
 309 Hungary, Latvia, Luxembourg, Slovakia, Sri Lanka

310


311 **ANZ=** Australia and New Zealand. **Other\_O: Other originating countries:** Cambodia, Pakistan, Phillipines, Hong Kong, Taiwan, Singapore,  
 312 South Korea, Afghanistan, Nepal.

313

314 During the two decades studied, very few notifications were recorded for foods ~~imported~~exported from  
315 Cambodia, Pakistan, Philippines, Hong Kong, Taiwan, Singapore, South Korea, Afghanistan, and Nepal  
316 (**Table 2**). Notifications were recorded by a total of 29 notifying countries. Among these, the highest  
317 number of notifications were from the United Kingdom (28% = 528/1873), Italy (12% = 221/1873) and  
318 the Netherlands (~10% = 179/1873), while the least number of notifications were reported from Cyprus,  
319 Austria, Slovenia, Ireland, Lithuania, Iceland, Romania, Switzerland, Bulgaria, Croatia, Estonia, Hungary,  
320 Latvia, Luxembourg, Slovakia and Sri Lanka (a total of only 92 notifications were reported in these 16  
321 countries over the two decades studied, **Table 2**).

### 322 3.3. Trends in RASFF notifications of pathogenic microorganisms across sub-regions and food 323 categories

324  
325 **Fig. 5** presents RASFF notifications of pathogenic microorganisms in food originating from the Asia-  
326 Pacific Region (n=1873) from 2000–2020, based on the notifying year. In the two decades considered,  
327 notifications were greatest in 2015 and 2011. Consistently over the years, food originating from South  
328 Asia was associated with the highest number of notifications while the least notifications were reported  
329 for Oceania (**Fig. 5**). Over the entire two decades of notifications data recorded in the RASFF database,  
330 there was no observable trend across sub-regions, with the notable exception of Oceania. Pearson  
331 correlation of year versus RASFF notifications of pathogenic microorganisms in food sub-regions  
332 indicates that notifications from Oceania have significantly reduced over the years ( $r=-0.68$ ,  $p=0.006$ ). In  
333 more recent years, however, (2015-2020), notifications from the sub-regions have reduced significantly,  
334 with the notable exception of East Asia (**Table 3a**). Across the food categories, there has been a  
335 statistically significant reduction in the number of notifications associated with seafoods over the years ( $r$   
336 =  $-0.73$ ,  $p=0.0001$ , **Table 3b**). In contrast, there has been a statistically significant increase in the number  
337 of notifications associated with nuts, nut products and seeds over the years ( $r = 0.57$ ,  $p=0.007$ , **Table**  
338 **3b**). Although a statistically significant increase in notifications for fruits and vegetables ( $r = 0.66$ ,  
339  $p=0.008$ , **Table 3b**) was recorded between 2000-2014, more recent years (2015-2020) have been  
340 associated with reducing trends in the number of notifications associated with this food type ( $r = -0.85$ ,  
341  $p=0.03$ , **Table 3b**).



345

346 **Fig. 5.** RASFF notifications of pathogenic microorganisms in food originating from Asia-Pacific Region  
 347 (n=1873) from 2000 - 2020 based on notifying year.

348 **Table 3**

349 Pearson correlation of year versus RASFF notifications of pathogenic microorganisms in various (a) Asia-Pacific sub-regions and (b) food  
 350 categories.

**a) Yearly notification versus sub-region**

Period	Statistics	East Asia	Oceania	South Asia	Southeast Asia	Grand Total
2000-2020	Pearson Correlation	.486*	-.676**	.452*	-0.188	0.353
	Sig. (2-tailed)	0.025	0.006	0.040	0.415	0.116
	N	21	15	21	21	21
2000-2014	Pearson Correlation	0.205	-.731*	.653**	0.025	.562*
	Sig. (2-tailed)	0.464	0.011	0.008	0.929	0.029
	N	15	11	15	15	15
2015-2020	Pearson Correlation	.821*	. <sup>b</sup>	-.873*	-.957**	-.918**
	Sig. (2-tailed)	0.045		0.023	0.003	0.010
	N	6	4	6	6	6

**b) Yearly notification versus food category**

Period	Statistics	nuts, nut products and seeds	seafoods	fruits and vegetables	herbs and spices	meat	others	Grand Total
2000-2020	Pearson Correlation	.567**	-.725**	0.387	0.004	0.354	0.419	0.294
	Sig. (2-tailed)	0.007	0.000	0.083	0.988	0.115	0.059	0.197
	N	21	21	21	19	21	21	21
2000-2014	Pearson Correlation	.655**	-.727**	.658**	0.187	-0.258	0.496	0.483
	Sig. (2-tailed)	0.008	0.002	0.008	0.541	0.353	0.060	0.068
	N	15	15	15	13	15	15	15
2015-2020	Pearson Correlation	-0.681	-0.709	-.849*	-0.332	-.878*	-0.427	-.915*
	Sig. (2-tailed)	0.136	0.115	0.032	0.520	0.021	0.399	0.011
	N	6	6	6	6	6	6	6

\*. Correlation is significant at the 0.05 level (2-tailed). \*\*. Correlation is significant at the 0.01 level (2-tailed). b- too few samples, blue cells indicate a downward trend, and red cells indicate an upward trend

351

352

353  
354 **3.4. Pathogens associated with notifications**  
355

356 Although the total number of notifications was 1873, a higher total number of pathogens (n=2121) was  
357 recorded because of co-occurrence of pathogens in some food products (**Table 4**). Among the 29  
358 notifying countries, more than a quarter (28%) of pathogen notifications for foods originating from the  
359 Asia-Pacific Region were reported by the United Kingdom (28%, i.e., 587/2121, see **Supplementary**  
360 **Table 1**). Cumulatively, the United Kingdom, Italy, Norway, the Netherlands, Germany and Finland  
361 accounted for more than 70% (i.e., 1556/2121) of pathogen notifications in the two-decade study period  
362 (**Supplementary Table 1**). The least number of pathogen notifications was observed for France, Poland,  
363 Portugal, Belgium, Cyprus, Austria, Slovenia, Ireland, Lithuania, Iceland, Romania, Switzerland,  
364 Bulgaria, Croatia, Estonia, Luxembourg, Hungary, Latvia and Slovakia, as these 19 countries accounted  
365 for less than 15% (i.e., 277/2121) of pathogen notifications in the two-decade study period  
366 (**Supplementary Table 1**).  
367

368 Among the 2121 notifications of pathogenic microorganisms from 2000-2020, the following 14 genera  
369 consisting of *Salmonella*, *Vibrio*, *Bacillus*, *Escherichia*, *Norovirus*, *Clostridium*, *Campylobacter*,  
370 *Staphylococcus*, *Enterococcus*, *Hepatitis*, *Pseudomonas*, *Cronobacter*, and *Plesiomonas*, were  
371 observed in this study (**Table 4**). Among these, the least commonly isolated pathogen genera were  
372 observed to be *Cronobacter*, Fungi (moulds - no specific genus was identified), *Plesiomonas*, *Shigella*,  
373 *Yersinia* and *Streptococci* (**Table 4**). Conversely, *Salmonella* was the most predominant as  
374 approximately 7 out of every 10 pathogens isolated from ~~foods imported from exported foods from~~ the  
375 Asia-Pacific Region in the past two decades (74%, 1560/2121) were *Salmonella* species (**Table 4**).  
376 Foods exported from India had the highest pathogen notifications (642/2121, **Table 4**).  
377

378 **Table 4**

379 RASFF notifications of pathogenic microorganisms in food originating from the Asia-Pacific Region (n=1873) from 2000 - 2020 based on  
380 originating country.

Pathogen genus	Country of origin										Grand Total (%)
	India	Thailand	Vietnam	Bangladesh	China	Malaysia	Indonesia	Laos	ANZ	Others	
<i>Salmonella</i>	551	411	182	165	69	30	38	30	23	61	1560(73.55%)
<i>Vibrio</i>	61	40	59	35	12	62	31	0	1	8	309(14.57%)
<i>Bacillus</i>	16	5	14	2	42	0	3	0	0	10	92(4.34%)
<i>Escherichia</i>	3	16	5	0	2	4	0	15	0	2	47(2.22%)
Norovirus	0	0	30	0	11	0	0	0	0	1	42(1.98%)
Not specified	6	2	5	0	4	0	0	0	0	11	28(1.32%)
<i>Clostridium</i>	3	5	2	0	12	0	0	0	0	0	22(1.04%)
<i>Campylobacter</i>	0	5	1	0	0	0	0	0	0	0	6(0.28%)
<i>Staphylococcus</i>	0	1	2	0	0	0	0	0	0	0	3(0.14%)
<i>Enterococcus</i>	1	0	0	0	0	0	1	0	0	0	2(0.09%)
Hepatitis	0	0	0	0	1	0	0	0	1	0	2(0.09%)
<i>Pseudomonas</i>	1	0	0	0	0	0	0	0	0	1	2(0.09%)
<i>Cronobacter</i>	0	0	0	0	1	0	0	0	0	0	1(0.05%)
Fungi	0	0	0	0	1	0	0	0	0	0	1(0.05%)
<i>Plesiomonas</i>	0	0	0	0	0	0	1	0	0	0	1(0.05%)
<i>Shigella</i>	0	1	0	0	0	0	0	0	0	0	1(0.05%)
<i>Yersinia</i>	0	1	0	0	0	0	0	0	0	0	1(0.05%)
<i>Streptococci</i>	0	0	0	0	1	0	0	0	0	0	1(0.05%)


381

382 **Other\_N: Other notifying countries:** Cyprus, Austria, Slovenia, Ireland, Lithuania, Iceland, Romania, Switzerland, Bulgaria, Croatia, Estonia,  
383 Hungary, Latvia, Luxembourg, Slovakia, Sri Lanka. **Other\_O: Other originating countries:** Australia, Cambodia, New Zealand, Pakistan,  
384 Philippines, Hong Kong, Taiwan, Singapore, South Korea, Afghanistan, Nepal.

385 The total number of pathogens (n=2119) exceeds the total number of notifications (n=1873) because of the co-occurrence of pathogens.

386 Among the pathogens identified in **imported/exported** foods originating from the Asia-Pacific Region, India  
 387 also had the highest *Salmonella* notifications (551/1560). On the whole, more than 80% (i.e. 1309/1560)  
 388 of *Salmonella* notifications were from foods **imported/exported** from India, Thailand, Vietnam and  
 389 Bangladesh. Forty percent (i.e., 625/1560) of the time, pathogen identification did not exceed the genus  
 390 level for RASFF notifications of *Salmonella* in food originating from the Asia-Pacific Region during the  
 391 two-decade study period (**Table 5**). Hence in these instances, the species or serovar implicated in the  
 392 *Salmonella* notification was not identified. In the remaining 935 instances of *Salmonella* notifications in  
 393 food originating from the Asia-Pacific Region, a total of 178 serovars of *Salmonella enterica* were  
 394 identified (**Table 5**). The top 10 most frequently isolated serovars in the RASFF *Salmonella* notifications  
 395 are serovars Weltevreden, Enteritidis, Hvittingfoss, Agona, Stanley, Senftenberg, Augustenborg, Brunei,  
 396 Typhimurium, Lexington, Newport and Mbandaka (**Table 5**).

398 Apart from *Salmonella*, *Vibrio* was the next most common, albeit at a reported frequency far lower than  
 399 that of *Salmonella* (14.6% = 309/1562). The most commonly isolated *Vibrio* in food products  
 400 **imported/exported** from the Asia-Pacific Region from 2000 – 2020 were *V. cholerae* (46.6%, i.e., 144/309)  
 401 and *V. parahaemolyticus* (46%, i.e., 142/309) while the least isolated *Vibrio* were *V. fluvialis* and *V.*  
 402 *mimicus* (0.32% each, i.e., 1/309) (**Fig. 6**).




405  
 406 **Fig. 6.** Species identified in 309 RASFF notifications of *Vibrio* in food originating from Asia-Pacific  
 407 Region from 2000 - 2020.

408 Apart from hepatitis A and norovirus, viruses were generally not detected in the foods  
 409 **imported/exported** from the Asia-Pacific region (**Table 4**). When Hepatitis A was reported in  
 410 food items, both instances were in 2012 when frozen green mussels (*Perna canaliculus*) from  
 411

412 New Zealand and frozen strawberry cubes from China tested positive for the virus. The action  
413 taken in each instance was different. In the case of contaminated frozen green mussels from  
414 New Zealand, the notifying country (Italy) withdrew the product from the market. In  
415 contaminated frozen strawberry cubes from China, the notifying country (Belgium) only  
416 informed recipients. Norovirus detection in foods imported from the Asia-Pacific region was  
417 reported 42 times in the two-decade study period (**Table 4**). Eleven of these notifications  
418 were from raspberries, seaweed salad and strawberries ~~imported~~ exported by China and thirty  
419 other instances were from other fruits ~~imported~~ exported by Vietnam between 2011 and 2019.  
420 In 6 out of the 11 norovirus notifications for fruits and vegetables from China, notifications were  
421 informed by the occurrence of ongoing food poisoning or consumer complaints.  
422 There was no consistency in response to ~~N~~orovirus notifications as the action taken varied  
423 between countries. For instance, actions taken included rejection at the border (Denmark and  
424 Lithuania). Other actions included recall from consumers (Denmark), withdrawal from the  
425 market (Netherlands, Germany), product recall or withdrawal (Sweden), withdrawal from  
426 recipients or destruction (Germany), public warning - press release (Norway), and informing  
427 recipients (Spain). The two instances when ~~h~~Hepatitis A was reported in food items were in  
428 2012 when frozen green mussels (*Perna canaliculus*) from New Zealand and frozen  
429 strawberry cubes from China tested positive to the virus. The action taken in each instance  
430 was different. In the case of contaminated frozen green mussels from New Zealand, the  
431 notifying country (Italy) withdrew the product from the market while in the case of contaminated  
432 frozen strawberry cubes from China, the notifying country (Belgium) only informed recipient.  
433 *Campylobacter* notifications in foods originating from the Asia Pacific region were made only  
434 in six instances (between 2009 and 2012) for asparagus, corn, ground peppers and spring  
435 onions from Thailand and Vietnam. Nearly half (46%, i.e., 42/92) of *Bacillus* notifications in  
436 food items from the Asia Pacific were from China.

#### 437 3.4. Pathogens associated with specific product categories

438 **Fig. 7** presents RASFF notifications of pathogenic microorganisms in food originating from  
439 Asia-Pacific Region (n=1873) from 2020 - 2020 based on product category. The most  
440 important product categories with *Salmonella* contamination were fruits and vegetables, herbs  
441 and spices, nuts, nut products and seeds, while the most important product category with  
442 *Vibrio* contamination was crustaceans and related products (**Fig. 7**).  
444



445

446 **Fig. 7** RASFF notifications of pathogenic microorganisms in food originating from the Asia-Pacific Region (n=1873) from 2020 - 2020 based on  
 447 food type.

448 A further analysis of pathogen diversity associated with notifications (2000 - 2020) revealed  
449 that crustaceans and related products from India, Malaysia and Vietnam showed the highest  
450 number of pathogen species (n >60) (**Supplementary Table 2**). Most of the pathogen species  
451 in crustaceans and related products in these countries were associated with shrimps (Table  
452 6). Sri Lanka and Pakistan presented with the least (n=7) pathogen species in crustaceans  
453 and related products ~~imported/exported~~ from the Asia-Pacific Region (**Supplementary Table**  
454 **2**). Among the Asia-Pacific countries, the highest counts of pathogen species in fruits and  
455 vegetables were observed among ~~import/export~~ from India and Bangladesh  
456 (**Supplementary Table 3**). More than 95% (i.e., India=184/193, Bangladesh = 149/152) of  
457 pathogen species notifications for fruits and vegetables ~~imported/exported~~ from India and  
458 Bangladesh were associated with betel leaves (**Supplementary Table 3**). All the 346  
459 pathogen notifications in fruits and vegetables ~~imported/exported~~ from India and Bangladesh  
460 were *Salmonella* serovars (**Supplementary Table 3**). Pathogen notifications for mushrooms  
461 ~~imported/exported~~ from the Asia-Pacific countries were all due to *Salmonella*, with the notable  
462 exception of mushrooms ~~imported/exported~~ from China, which had additional pathogen  
463 notifications due to *Bacillus cereus*, *Clostridium perfringens*, *Clostridium* spp, aerobic  
464 mesophiles and moulds. Among the Asia-Pacific countries, spinach imported from Thailand  
465 had the highest number of reported *Salmonella* serovars (*Salmonella* Ndolo, *Salmonella*  
466 Aberdeen, *Salmonella* Bovismorbificans, *Salmonella enterica*, *Salmonella* group B,  
467 *Salmonella* group D, *Salmonella* Hvittingfoss, *Salmonella* Mbandaka, *Salmonella* Newport -  
468 see **Supplementary Table 3**).

469  
470 Among the nuts, nut products and seeds, sesame seeds were the main food item  
471 contaminated by *Salmonella* (261 notifications), and these accounted for 87% of total  
472 *Salmonella* notifications (261/299, **Supplementary Table 4**). *Salmonella*-contaminated  
473 sesame seeds originated from India (n=260; 99.6%) and Thailand (n=1; 0.4%)  
474 (**Supplementary Table 5**). Fifty serovars of *Salmonella enterica* were isolated from sesame  
475 seeds (**Supplementary Table 4**). Among the herbs and spices, basil, mint and coriander from  
476 Thailand had the highest number of pathogen notifications. For instance, basil was associated  
477 with at least 8 *Salmonella enterica* serovars and 65 notifications, coriander with at least 12  
478 *Salmonella enterica* serovars and 24 notifications, and mint with at least 2 *Salmonella enterica*  
479 serovars and 25 notifications (**Supplementary Table 6**).

480

481

482

483

484 **4. Discussion**

485 The presence of pathogenic microorganisms in food products, particularly in minimally  
486 processed and ready-to-eat foods, constitutes a significant public health risk to consumers. In  
487 this study, we analysed notifications associated with microbiological hazards (MH) in foods  
488 ~~imported~~~~exported~~ from the Asia Pacific region into the EU between 2000 and 2020. Studies  
489 of this nature can potentially encourage countries to implement policies that improve quality  
490 of exported food products.

491 Unlike a previous report (Somorin, et al., 2021) which noted a growing trend in notifications  
492 from food products ~~imported~~~~exported~~ from Africa, there was no overall trend in the number of  
493 notifications from the Asia Pacific region during the same study period. This difference could  
494 be attributed to the significant decrease in MH notifications from the South and Southeast Asia  
495 subregions between 2015 and 2020. Interestingly, the countries responsible for the highest  
496 number of notifications during the study period are located in the South and Southeast Asia  
497 subregions.  
498

499 Of the 1873 notifications, most (n=1456/1873) of them concerned food originating from India,  
500 Thailand, Vietnam and Bangladesh. A contributory factor could be the volume of trade  
501 between these countries and the EU. In 2018/19, India, Thailand and Vietnam ranked 10th,  
502 14th and 17th, respectively, in terms of ~~for the export~~ value of food products traded with the  
503 EU (European Commission, 2020a). However, this appears not to be the sole reason.  
504 Compared to other countries in the Asia-Pacific region, EU food imports from China and  
505 Indonesia are the largest, with a combined value of about 10 billion euros (European  
506 Commission, 2020a), but only contributed 11% of notifications observed during the study  
507 period. New Zealand is the 19th highest exporter of food to the EU but contributed the least  
508 number of notifications from the region.  
509

510 A more likely explanation for our observations is the nature of the microbial hazard and  
511 implicated foods. The highest number of MH notifications from the Asia-Pacific region was  
512 recorded in 2015. Aflatoxin contamination of nuts, nut products and seeds imported from  
513 China and *Salmonella* contamination in fruits and vegetables and nuts, nut products and seeds  
514 imported from India produced the three highest MH notifications from all regions for 2015  
515 (European Commission 2016). In addition, *Salmonella* contamination of herbs and spices,  
516 fruits and vegetables, nuts, nut products, and seeds were responsible for the highest number  
517 of notifications overall. These ~~products~~ were predominantly products ~~importeds~~ from India,  
518 Thailand, Vietnam and Bangladesh. Food safety remains a significant challenge in the Asia-  
519 Pacific region and a hurdle to expanding agricultural and food product exports (Pham & Dinh,  
520 2020). Food safety challenges include inadequate sanitation infrastructure, poor food safety  
521 knowledge and practices, and insufficient implementation of national food safety management  
522 systems (Food and Agricultural Organisation, 2018; Jaffee, Henson, Unnevehr, Grace, &  
523 Cassou, 2018; Mangla, et al., 2021; Ortega, 2017).

524 The highest number of notifications was made by the UK (28%). This is understandable  
525 because the standard for food quality, in terms of compliance requirement, is generally high  
526 in the UK. With about half of UK's food produced locally (DEFRA, 2020), a recent survey  
527 showed that a majority (97.5%) of food manufacturers within the UK had an integrated food  
528 safety management system in place (Mensah & Julien, 2011). Only about 5% of UK's food are  
529 imported from Asia and Australasia (DEFRA, 2020), yet exporters wishing to supply the UK  
530 are expected to have monitoring systems that ensure compliance with retail (product quality)  
531 and legislative (due diligence) requirements in the UK (Dolan & Humphrey, 2000). Notifications  
532 made by the UK (28%) mainly concerned foods originating from India and Bangladesh. Food  
533 safety has been a major challenge in South Asia and Southeast Asia, with huge impacts on  
534 public health, economics, and international trade (World Health Organization, 2020a).  
535 Southeast Asia has the second highest annual burden of foodborne diseases among the six

536 World Health Organization (WHO) regions, with 150 million foodborne illnesses and 175,000  
537 deaths in 2010 (World Health Organization, 2015). In contrast to South Asia and Southeast  
538 Asia, there were significantly fewer notifications of pathogenic microorganisms in foods  
539 **imported** from Australia and New Zealand (n=23/1873; 1.2%), and this could be due to the  
540 strong food control systems in Australia and New Zealand (Ghosh, 2014).

541  
542 Data received through the RASFF are a vital source of information on emerging health risks  
543 associated with specific food products. Import border rejections were the most common  
544 notification type in this study, while the most common actions taken were product redispach,  
545 destruction and non-authorisation of imports. Besides constituting a significant revenue loss  
546 for exporting countries, and causing reduced consumer confidence, these actions may lead to  
547 more stringent responses from the EU to safeguard the health of consumers (Chaoniruthisai,  
548 Punnakitkashem, & Rajchamaha, 2018; Somorin, et al., 2021). For example, peppers  
549 (*Capsicum* spp. except sweet) from India and Pakistan were subjected to increased official  
550 controls in 2018 due to concerns about the levels of pesticide residues (European  
551 Commission, 2020b). Data from these official controls and RASFF notifications provided  
552 evidence of continuous non-compliance, and these food products were assessed as  
553 constituting a serious risk to human health. In 2020, the EU legislated that these products  
554 should now include an official certificate confirming that maximum pesticide levels have not  
555 been exceeded (European Commission, 2020b).

556  
557 Increased official controls are the first stage of a continuum that can end in the suspension of  
558 imports of a specific product. A good example is the suspension of imports into the EU of  
559 foodstuff containing betel leaves from Bangladesh due to the presence of *Salmonella*. This  
560 suspension has been in place since 2014 as no action plan for managing this hazard has been  
561 provided (European Commission, 2014, 2020b). A national monitoring programme for betel  
562 leaves imported into the United Kingdom between 2011 and 2017 showed that *Salmonella*  
563 contamination remains a significant hazard in this product (McLauchlin, et al., 2019). Betel is  
564 an important cash crop in Bangladesh and the principal source of livelihood in many rural  
565 communities (Nath & Inoue, 2009; Ullah, Tani, Tsuchiya, Rahman, & Rahman, 2020).  
566 **Import/Export** bans can have devastating consequences on income generation and food  
567 security. The poor food safety record in South Asia and Southeast Asia is due to several  
568 factors including climatic factors, poor sanitary conditions, and food habits. For example,  
569 extreme weather conditions and natural disasters, such as flooding, have been frequently  
570 occurring in countries Southeast Asia (A. Chen, Giese, & Chen, 2020), and since flood waters  
571 are often contaminated with sewage, this leads to increased contamination risk in the food  
572 supply chain and exposure to faecal pathogens (Yu, et al., 2018). Consumption of street-foods  
573 is common in many Asian countries and it is the only affordable form of nutrition for most  
574 people in low-income groups. However, inadequate observation of food hygiene practices  
575 significantly increases the exposure of consumers to foodborne pathogens (Reddy, Ricart, &  
576 Cadman, 2020).

577  
578 There was a significant ( $p < 0.006$ ) downward trend in the numbers of pathogenic  
579 microorganism notifications from foods **imported/exported** from Oceania to the EU from 2000-  
580 2020, unlike South Asia and East Asia. In more recent years (2015-2020), East Asia was the  
581 only sub-region with a significantly increasing trend in RASFF notifications due to  
582 contamination by pathogenic microorganisms. When considered by food categories, RASFF  
583 notifications regarding “nut, nut products and seeds” significantly increased over the years  
584 from 2000 - 2020. However, notifications of pathogenic microorganisms in “fruits and  
585 vegetables” significantly reduced in recent years (2015 – 2020). To address the food safety  
586 concerns in the affected regions, it is important that appropriate and effective methods to  
587 minimise contamination of foods by pathogenic microorganisms are implemented across the  
588 value chains in the originating countries. For instance, a framework has been recently  
589 established in the WHO Southeast Asia region (including India, Thailand and Bangladesh) to  
590 strengthen food control systems across the value chain and protect consumer health, building

591 on previous regional multi-actor interventions (World Health Organization, 2020b). These  
592 actions could have contributed to the downward trend in notifications of pathogenic  
593 microorganisms from South Asia and Southeast Asia in recent years (2015-2020) (Table 3).  
594 It is however important to take these RASFF database trend results cautiously As argued in  
595 published literature (Kowalska & Manning, 2020) iterative changes in food law potentially  
596 impacts on the frequency of regulatory sampling associated with border and inland regulatory  
597 checks in a way that could affect the generalisability of the trends noted.

598  
599 Border rejection was the most frequent type of notification received (42.3%) in this study.  
600 When food ~~importsexports~~ do not meet the standards set by the EU, a variety of actions can  
601 be taken, which have huge economic implications for the originating countries. About half of  
602 the notifications from the Asia-Pacific region resulted in importation not being authorized,  
603 leading to destruction or re-dispatch of the contaminated food product. Frequent non-  
604 compliance of some food products with EU food safety requirements has led to increased  
605 official control at border control posts and the imposition of suspension of entry into the EU  
606 (European Commission, 2020b). Increased identity and physical checks at EU border control  
607 points is in place for betel leaves and sesame seeds from India, and sweet peppers from China  
608 due to *Salmonella* contamination (European Commission, 2020b). Similar official controls  
609 exist for black pepper from Brazil as well as sesame seeds from Ethiopia, Nigeria, Sudan and  
610 Uganda due to *Salmonella* contamination (European Commission, 2020a, 2020b; Somorin, et  
611 al., 2021). Continuous non-compliance has led to the suspension of importation of betel leaves  
612 from Bangladesh to the EU since 2014 due to *Salmonella* contamination (European  
613 Commission, 2015). A similar ban is in place for dried beans from Nigeria due to pesticide  
614 residues (European Commission, 2020a, 2020b).

615  
616 The most frequently reported genera of pathogenic microorganisms in foods from the Asia-  
617 Pacific region were *Salmonella* (73.6%), *Vibrio* (14.6%), *Bacillus* (4.3%) and *Escherichia*  
618 (2.2%). *Salmonella* spp. are the most reported pathogens associated with foods imported into  
619 the EU from the Asia-Pacific region. This corroborates the report by Somorin et al., (2021) on  
620 products imported into the EU from Africa. The predominance of *Salmonella*, an enteric  
621 bacterium, could be attributed to unhygienic practices during food production processes and  
622 this is corroborated by a previous study that reported a significant association between the  
623 presence of *Salmonella* spp. and *E. coli* in imported edible leaves retailed in the England  
624 (McLauchlin, et al., 2018). Foods reported in *Salmonella* notifications were mainly “fruits and  
625 vegetables”; “herbs and spices” and “nuts, nut products and seeds”. India accounted for over  
626 35% of the *Salmonella* notifications in this study and were mostly from “nuts, nut products and  
627 seeds”, “fruits and vegetables” and seafoods. In our study, about 178 serovars of *Salmonella*  
628 *enterica* were reported in foods from Asia-Pacific from 2000-2020 with serovars Weltevrede,  
629 Enteritidis and Hvittingfoss being the most frequently reported. These serovars are significant  
630 causes of diarrhoea, and sometimes, invasive infections in tropical low-income Asian  
631 countries (Bangtrakulnonth, et al., 2004; Kantama & Jayanetra, 1996; Makendi, et al., 2016)  
632 as well as in Europe (ECDC/EFSA, 2021; Emberland, et al., 2007).

633  
634 Over 95% of notifications regarding “fruit/vegetables” were from India and Bangladesh.  
635 Several *Salmonella* serovars have been shown to contaminate ready-to-eat betel leaves in  
636 Asia and those imported into the UK. Fakruddin *et al.* (2017) reported that 77% of betel leaves  
637 collected in local markets in Bangladesh were contaminated with *Salmonella* spp., with some  
638 of them imported from India. *Salmonella* from betel leaves also harbour multidrug resistance  
639 genes (Singh, et al., 2006), which could be disseminated to other countries through  
640 international trade. The majority of the notifications regarding *Salmonella* in betel leaves were  
641 from the UK, affecting products originating from India and Bangladesh. With South Asians  
642 being the largest ethnic minority group in the UK, and accounting for about 5% of the UK  
643 population (Office of National Statistics, 2011), there is a ready demand for betel leaves and  
644 other indigenous foods. Analysis of betel leaf consignments at Border Inspection Posts in  
645 England between 2011 and 2017 showed that 57% had *Salmonella* contamination

646 (McLauchlin, et al., 2019). In a separate study by McLauchlin *et al.* (2018), 14% of betel leaves  
647 collected from retailers in England had unsatisfactory levels of *Salmonella* including  
648 *Salmonella* Bareilly, *Salmonella* Hvittingfoss, *Salmonella* Lichfield, and *Salmonella* Newport.  
649 The frequency and high contamination rates of *Salmonella* in betel leaves pose a public health  
650 concern for importing countries. While the UK is no longer part of the EU, it is expected to  
651 remain an important market for produce from South Asia and Southeast Asia and retain  
652 stringent food safety regulations.

653  
654 *Salmonella* is a principal aetiological agent of foodborne disease and is the third leading cause  
655 of death due to diarrhoeal disease globally (Ferrari, et al., 2019; World Health Organization,  
656 2019, 2020c). *Salmonella* is the most common cause of foodborne outbreaks in the EU and  
657 in 2018, was implicated in almost 30% of outbreaks in the region (EFSA/ECDC, 2019). Fresh  
658 produce, including fruits, vegetables, herbs, and spices ~~imported~~~~exported~~ from the Asia-  
659 Pacific region, were the food products most associated with *Salmonella*. Fruit and vegetables  
660 are increasingly recognised as vehicles for transmitting pathogenic organisms and have been  
661 implicated in global foodborne outbreaks (Bisht, et al., 2021; Carstens, Salazar, & Darkoh,  
662 2019; Wadamori, Gooneratne, & Hussain, 2017).

663 Contamination of fresh produce can occur at different stages of the supply chain. *Salmonella*  
664 is widespread in the environment and is commonly found in materials susceptible to faecal  
665 contamination (Machado-Moreira, Richards, Brennan, Abram, & Burgess, 2019).  
666 Contaminated soil and irrigation water are important sources of preharvest contamination.  
667 Unhygienic practices during harvest, processing and transport are the main contributors to  
668 post-harvest contamination (Dos Santos, et al., 2020; Park, et al., 2012). Fresh produce is  
669 usually consumed with minimal processing, making it challenging to eradicate pathogenic  
670 microorganisms via conventional methods like cooking. Several studies have noted  
671 unacceptable levels of *Salmonella* contamination in fresh produce available for retail sale in  
672 the Asia-Pacific region (Abatcha, Effarizah, & Rusul, 2018; Nguyen, et al., 2021; Vital,  
673 Dimasuay, Widmer, & Rivera, 2014). These reports are of concern and highlight the need for  
674 improved food safety management practices along the fresh produce supply chain. To our  
675 knowledge, no outbreaks linked to *Salmonella* in fresh produce from the Asia-Pacific region  
676 have been reported in the EU. However, it is not always possible to identify the potential source  
677 and vehicle of infection during an outbreak (EFSA/ECDC, 2018).

678 The most frequently reported *Vibrio* spp. in the notifications were *V. cholerae* (46.6%) and *V.*  
679 *parahaemolyticus* (46%) while *V. vulnificus*, *V. alginolyticus*, *V. fluvialis* and *V. mimicus* were  
680 less frequently isolated. The main food category contaminated by *Vibrio* was “crustaceans and  
681 related products” particularly shrimps and prawns. Precious studies have reported that most  
682 of the foods with *Vibrio* contamination originated from India, Malaysia and Vietnam. For  
683 instance, *V. parahaemolyticus* was reported in 76% of shrimps ~~imported~~~~exported~~ from  
684 Thailand (Wong, Chen, Liu, & Liu, 1999) and 43.4% of shrimps in China (Xu, Cheng, Wu,  
685 Zhang, & Xie, 2016). A *V. parahaemolyticus* pandemic serotype O3:K6 originating from India  
686 have been detected in many Asian countries (Bag, et al., 1999) as well as in Europe (Martinez-  
687 Urtaza, et al., 2005). predominantly linked to crustaceans such as shrimps and prawns and  
688 related products. *Vibrio* spp. are important causes of diarrhoea and cholera associated with  
689 the consumption of raw or undercooked seafood and are frequently detected in seafood  
690 products from countries in the Asia-Pacific region (X. Chen, et al., 2018; Fu, et al., 2020;  
691 Kumar & Lalitha, 2013; Lee, et al., 2019; Nakaguchi, 2013). Seafood imported from Asia was  
692 associated with an outbreak in France in 1997 (Lemoine, Germanetto, & Giraud, 1999).  
693 Similarly, Briet et al. (2018) reported the presence of a carbapenemase-producing strain of *V.*  
694 *parahaemolyticus* isolated from shrimps imported into France from Vietnam. This study  
695 provides evidence that imported foods may be potential vehicles for the transmission of  
696 clinically relevant microorganisms.

697  
698 The predominant *Bacillus* spp. reported in the notifications was *Bacillus cereus* and it was  
699 reported in cheese, soy cheese, tofu, garlic powder, ground pepper, pumpkin seeds,  
700 mushroom, sesame seeds, desiccated coconut, cinnamon powder, curry powder, ginger  
701 powder and ground cumin across East-, South- and Southeast Asia. Production and  
702 processing of these food products involves drying and/or grinding, where they could come in  
703 contact with soil harbouring *Bacillus* spores (Vilain, Luo, Hildreth, & Brözel, 2006). *B. cereus*  
704 has been previously reported in soybean products/tofu (Ananchaipattana, et al., 2012;  
705 Keisam, Tuikhar, Ahmed, & Jeyaram, 2019), spice and herbs (Banerjee & Sarkar, 2003) and  
706 cheese (Kumari & Sarkar, 2014).

707  
708 Furthermore, foodborne viruses, such as Hepatitis A and Norovirus, were reported in green  
709 mussels and berries (strawberry and raspberry) mainly originating from Vietnam and China.  
710 Norovirus has been reported in whole fruits and vegetable leaves in France (EFSA/ECDC,  
711 2021) and the UK (Cook, Williams, & D'Agostino, 2019) and the majority of foodborne viral  
712 outbreaks are associated with frozen fruits (Nasheri, Vester, & Petronella, 2019). A large  
713 outbreak in Germany in 2012 was associated with strawberries imported from China (Bernard,  
714 et al., 2014). Norovirus was associated with 22.5% of all outbreak-related illnesses in the EU  
715 in 2019, with contaminated shellfish and fish products playing a significant role (EFSA/ECDC,  
716 2021). Although there were few notifications regarding *Campylobacter* and *Yersinia*, and no  
717 report of *Listeria monocytogenes* in foods from the Asia-Pacific region, foodborne diseases  
718 caused by these pathogens are among the most common in Europe (EFSA/ECDC, 2021),  
719 hence requiring continuous surveillance of imported foods from other parts of the world. The  
720 lack of consistency in the actions taken by different EU member states in response to  
721 notifications arising from the presence of norovirus and Hepatitis A is of interest. This suggests  
722 a lack in uniformity across Member States with respect to pathogens detected in food items.  
723 To safeguard public health, it may be necessary for stakeholders to collate a guidance action  
724 document with respect to specific actions that should be taken if a particular pathogen is  
725 detected in imported food products. This will ensure consistency across notifying countries.  
726 One limitation with the current study is the lack of comparison between the numbers of  
727 pathogen notifications and the total number of foods or food product category from each  
728 country that is tested for pathogens. This is partly because this data is not recorded in the  
729 RASSF database. It is hoped that these considerations will be included in future improvements  
730 in the RASSF database.

## 731 732 **5. Conclusion**

733 The analysis of RASFF notifications for foods from the Asia-Pacific region from 2000 – 2020  
734 showed that foods from India, Thailand, Vietnam and Bangladesh had the most notifications  
735 due to pathogenic microorganisms. *Salmonella* was the most predominant foodborne  
736 pathogen reported in foods ~~imported~~~~exported~~ from Asia-Pacific to the EU, with India  
737 accounting for over 35% of *Salmonella* notifications (fresh produce) and *Vibrio* spp.  
738 contamination of seafood products such as shrimps and prawns from India, Malaysia, and  
739 Vietnam. There were very few notifications regarding *Campylobacter* spp. and *Yersinia* spp.,  
740 and no notification of *L. monocytogenes* was reported. Increasingly, food producers are  
741 connected to consumers via food supply chains that cross international borders. These value  
742 chains offer a stable supply of food that meet consumers varied needs and preferences and  
743 provide a source of income to producers (OECD, 2020). An improved understanding of the  
744 food safety challenges along these international supply chains, and capacity building to  
745 support implementation of good manufacturing practices and food management systems, is  
746 required to safeguard consumer health and sustain the economic development of food-  
747 producing nations. Similarly, improved hygienic practices should be implemented across the  
748 value chains of imported food products from the region to ensure continuous access to  
749 international markets and sustained income.

750  
751

752

753 **CRedit authorship contribution statement**

754 **Ayokunle C. Dada:** Conceptualization, Methodology, Formal analysis, Writing - original draft,  
755 Review and Editing, Writing - review & editing. **Yinka M. Somorin:** Writing - original draft,  
756 Review and Editing, Writing - review & editing. **Collins N. Ateba:** Writing - original draft,  
757 Review and Editing, Writing - review & editing. **Helen Onyeaka:** Review and Editing, Writing  
758 - review & editing **Amarachukwu Anyogu:** Review and Editing, Writing - review & editing.  
759 **Nor Azman Kasan:** Writing - original draft, Review and Editing, Writing - review & editing.  
760 **Olumide A. Odeyemi:** Conceptualization, Methodology, Formal analysis, Writing - original  
761 draft, Review and Editing, Writing - review & editing.

762

763

764

765

766

767

768

769

770

771 **References**

772

- 773 [Abatcha, M.G., Effarizah, M.E., Rusul, G. \(2018\). Prevalence, antimicrobial resistance,](#)  
774 [resistance genes and class 1 integrons of \*Salmonella\* serovars in leafy vegetables,](#)  
775 [chicken carcasses and related processing environments in Malaysian fresh food](#)  
776 [markets. \*Food Control\*, 91, 170-180. <https://doi.org/10.1016/j.foodcont.2018.02.030>.](#)  
777 [Ananchaipattana, C., Hosotani, Y., Kawasaki, S., Pongewat, S., Latiful, B., Isobe, S. and Inatsu,](#)  
778 [Y. \(2012\); Bacterial contamination of soybean curd \(tofu\) sold in Thailand. \*Food Sci.\*  
779 \[Technol. Res.\]\(#\) 18\(6\), 843–848](#)
- 780 [Bag, P. K., Nandi, S., Bhadra, R. K., Ramamurthy, T., Bhattacharya, S. K., Nishibuchi, M.,](#)  
781 [Hamabata, T., Yamasaki, S., Takeda, Y. and Nair, G. B. \(1999\). Clonal diversity](#)  
782 [among recently emerged strains of \*Vibrio parahaemolyticus\* O3:K6 associated with](#)  
783 [pandemic spread. \*J Clin Microbiol\* 37\(7\): 2354–2357.](#)
- 784 [Balali, G.I., Yar, D.D., Dela, G.B.A., Adjoi Kusi, P. \(2020\). Microbial Contamination, an](#)  
785 [Increasing Threat to the Consumption of Fresh Fruits and Vegetables in Today's](#)  
786 [World. \*International Journal of Microbiology\*, 3029295,](#)  
787 [https://doi.org/10.1155/2020/3029295](#)
- 788 [Banerjee, M. and Sarkar, P. K. \(2003\). Microbiological quality of some retail spices in India.](#)  
789 [\*Food Research International\* 36: 469–474.](#)
- 790 [Bangtrakulnonth, A., Pornreongwong, S., Pulsrikarn, C., Sawanpanyalert, P., Hendriksen, R.](#)  
791 [S., Wong, D. M. and Aarestrup, F. M. \(2004\). \*Salmonella\* Serovars from Humans and](#)  
792 [Other Sources in Thailand, 1993–2002. \*Emerging Infectious Diseases\*, 10\(1\), 131–](#)  
793 [136.](#)
- 794 [Bernard, H., Faber, M., Wilking, H., Haller, S., Höhle, M., Schielke, A., Ducomble, T., Sifczyk,](#)  
795 [C., Merbecks, S. S., Fricke, G., Hamouda, O., Stark, K., Werber, D. a on behalf of](#)  
796 [the Outbreak Investigation Team. \(2014\). Large multistate outbreak of norovirus](#)  
797 [gastroenteritis associated with frozen strawberries, Germany, 2012. \*Euro Surveill.\*](#)  
798 [19\(8\): 20719.](#)
- 799 [Bisht, A., Kamble, M.P., Choudhary, P., Chaturvedi, K., Kohli, G., Juneja, V.K., Sehgal, S.,](#)  
800 [Taneja, N.K. \(2020\). A surveillance of foodborne disease outbreaks in India: 2009–](#)  
801 [2018. \*Food Control\*, 121, 107630. <https://doi.org/10.1016/j.foodcont.2020.107630>.](#)

802 Borchers, A., Teuber, S. S., Keen, C. L., & Gershwin, M. E. (2010). Food safety. Clinical  
803 reviews in allergy & immunology, 39(2), 95-141.

804 Briet A, Holsens N, Delannoy S, Debuiche S, Brisabois A, Midelet G, Granier SA. (2018).  
805 NDM 1-producing *Vibrio parahaemolyticus* isolated from imported seafood. J  
806 Antimicrob Chemother. 73, 2578-2579. <https://doi.org/10.1093/jac/dky200>.

807 Carstens, C. K., Salazar, J. K., & Darkoh, C. (2019). Multistate Outbreaks of Foodborne Illness  
808 in the United States Associated With Fresh Produce From 2010 to 2017. Frontiers in  
809 Microbiology, 10, 2667. <https://doi.org/10.3389/fmicb.2019.02667>

810 Chaoniruthicai, P., Punnakitikashem, P., Rajchamaha, K. (2018). Challenges and difficulties  
811 in the implementation of a food safety management system in Thailand: A survey of  
812 BRC certified food productions, Food Control, 93, 274-282,  
813 <https://doi.org/10.1016/j.foodcont.2018.06.004>.

814 Chen, A., Giese, M. and Chen, D. 2020. Flood impact on Mainland Southeast Asia between  
815 1985 and 2018—The role of tropical cyclones. J Flood Risk Management 13: e12598.

816 Chen, X., Zhu, Q., Yu, F., Zhang, W., Wang, R., Ye, X., Jin, L., Liu, Y., Li, S., & Chen, Y.  
817 (2018). Serology, virulence and molecular characteristics of *Vibrio parahaemolyticus*  
818 isolated from seafood in Zhejiang province. PloS one, 13(10), e0204892.  
819 <https://doi.org/10.1371/journal.pone.0204892>

820 Commission, E. (2020a). The Rapid Alert System for Food and Feed Annual Report 2019.  
821 Retrieved from [https://op.europa.eu/en/publication-detail/-/publication/2c5c7729-  
822 0c31-11eb-bc07-01aa75ed71a1/language-en/format-PDF/source-174742448](https://op.europa.eu/en/publication-detail/-/publication/2c5c7729-0c31-11eb-bc07-01aa75ed71a1/language-en/format-PDF/source-174742448)

823 Commission, E. (2020b). RASFF Food and Feed Safety Alerts. Retrieved from  
824 [https://ec.europa.eu/food/safety/rasff\\_en](https://ec.europa.eu/food/safety/rasff_en)

825 Cook, N., Williams, L. and D'Agostino, M. 2019. Prevalence of norovirus in produce sold at  
826 retail in the United Kingdom. Food Microbiology 79: 85-89.

827 D'Ortenzio, E., Weill, F. X., Ragonneau, S., Lebon, J. A., Renault, P. and Pierre, V. 2008. First  
828 report of a *Salmonella enterica* serovar Weltevreden outbreak on Réunion Island,  
829 France, August 2007. Euro Surveill. 13(32): 18949.

830 ECDC and EFSA (European Centre for Disease Prevention and Control/European Food  
831 Safety Authority) (2018). Multi-country outbreak of *Salmonella* Agona possibly linked  
832 to ready-to-eat food—26 July 2018. Stockholm and Parma.

833 EFSA and ECDC (European Food Safety Authority and European Centre for Disease  
834 Prevention and Control), (2019). The European Union One Health 2018 Zoonoses  
835 Report. EFSA Journal, 17:5926. <https://doi.org/10.2903/j.efsa.2019.5926>

836 EFSA. (2019). The European Union one health 2018 zoonoses report. European Centre for  
837 Disease Prevention, Control. European Food Safety Authority Journal, 17(12),  
838 e05926.

839 Emberland, K. E., Ethelberg, S., Kuusi, M., Vold, L., Jensvoll, L., Lindstedt, B. A., Nygård, K.,  
840 Kjelsø, C., Torpdahl, M., Sørensen, G., Jensen, T., Lukinmaa, S., Niskanen, T. and  
841 Kapperud, G. 2007. Outbreak of *Salmonella* Weltevreden infections in Norway,  
842 Denmark and Finland associated with alfalfa sprouts, July-October 2007. Euro  
843 Surveill. 12(48): 3321.

844 ESCAP. (2020). Economic and Social Survey of Asia and the Pacific 2020: Towards  
845 sustainable economies. Retrieved from  
846 [https://www.unescap.org/sites/default/files/publications/Economic%20and%20Social  
847 I%20Survey%20of%20Asia%20and%20the%20Pacific%202020%20Towards%20Sustainable%20Economies.pdf](https://www.unescap.org/sites/default/files/publications/Economic%20and%20Social%20Survey%20of%20Asia%20and%20the%20Pacific%202020%20Towards%20Sustainable%20Economies.pdf)

848

849 European Centre for Disease Prevention and Control (ECDC) and European Food Safety  
850 Authority (EFSA). 2021. Multi-country outbreak of *Salmonella* Enteritidis sequence  
851 type (ST)11 infections linked to poultry products in the EU/EEA and the United  
852 Kingdom—25 February 2021. Stockholm: ECDC/EFSA; 2021.

853 European Commission Health & Food Safety (2016). RASFF The Rapid Alert System for  
854 Food and Feed 2015 annual report. Available from:

855 [https://op.europa.eu/en/publication-detail/-/publication/98e04cfc-aa54-11e6-aab7-](https://op.europa.eu/en/publication-detail/-/publication/98e04cfc-aa54-11e6-aab7-01aa75ed71a1/language-en/format-PDF/source-174743070)  
856 [01aa75ed71a1/language-en/format-PDF/source-174743070](https://op.europa.eu/en/publication-detail/-/publication/98e04cfc-aa54-11e6-aab7-01aa75ed71a1/language-en/format-PDF/source-174743070) Accessed 19.02.2021

857 European Commission. (2020a). Monitoring EU Agri Food Trade: Developments in 2019.  
858 Available from: [https://ec.europa.eu/info/news/2019-record-year-eu-agri-food-trade-](https://ec.europa.eu/info/news/2019-record-year-eu-agri-food-trade-2020-mar-27-en#moreinfo)  
859 [2020-mar-27-en#moreinfo](https://ec.europa.eu/info/news/2019-record-year-eu-agri-food-trade-2020-mar-27-en#moreinfo) Accessed 12.02.2021.

860 European Commission, 2020. Commission implementing regulation (EU) 2020/625 of 7 May  
861 2020. *Official Journal of the European Union* 144: 13-33

862 European Commission. (2014). Commission implementing decision of 13 February 2014  
863 suspending temporarily imports from Bangladesh of foodstuffs containing or  
864 consisting of betel leaves ('Piper betle') Available from: [https://eur-](https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32014D0088&from=DA)  
865 [lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32014D0088&from=DA](https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32014D0088&from=DA)  
866 Accessed 25.02.2021.

867 European Commission. (2020b). Commission implementing regulation (EU) 2020/625 of 6  
868 May 2020. *Official Journal of the European Union*, 63, 13.

869 European Commission. 2015. Commission implementing regulation (EU) 2015/943 of 18 June  
870 2015. *Official Journal of the European Union* 154: 8-9

871 European Food Safety Authority (EFSA) and European Centre for Disease Prevention and  
872 Control (ECDC). 2021. The European Union One Health 2019 Zoonoses Report.  
873 *EFSA Journal* 19(2): 6406.

874 Fakruddin, M., Sultana, R., Hossain, M. N., Rahaman, M. M., Islam, M. K. and Ahmed, M. M.  
875 2017. Occurrence of ingression of *Salmonella* spp. in Betel leaf (*Piper betle* L.).  
876 *International Journal of Food Contamination* 4: 6.

877 FAO. (2018). Shiga toxin-producing *Escherichia coli* (STEC) and food: attribution,  
878 characterization, and monitoring. Retrieved from  
879 [https://apps.who.int/iris/bitstream/handle/10665/272871/0789241514279-](https://apps.who.int/iris/bitstream/handle/10665/272871/0789241514279-eng.pdf?ua=1)  
880 [eng.pdf?ua=1](https://apps.who.int/iris/bitstream/handle/10665/272871/0789241514279-eng.pdf?ua=1)

881 FAO. (2019). Foodborne illnesses in Asia-Pacific 'needlessly' sicken 275 million annually and  
882 threaten trade. Retrieved from [http://www.fao.org/asiapacific/news/detail-](http://www.fao.org/asiapacific/news/detail-events/en/c/1197008/#:~:text=In%20the%20Asia%2DPacific%20region,of%20Zero%20Hunger%20by%202030.)  
883 [events/en/c/1197008/#:~:text=In%20the%20Asia%2DPacific%20region,of%20Zero](http://www.fao.org/asiapacific/news/detail-events/en/c/1197008/#:~:text=In%20the%20Asia%2DPacific%20region,of%20Zero%20Hunger%20by%202030.)  
884 [%20Hunger%20by%202030.](http://www.fao.org/asiapacific/news/detail-events/en/c/1197008/#:~:text=In%20the%20Asia%2DPacific%20region,of%20Zero%20Hunger%20by%202030.)

885 Ferrari, R.G., Rosario, D.K.A., Cunha Neto, A., Mano, S.B., Figueiredo, E.E.S., Conte Junior,  
886 C.A. (2019). Worldwide Epidemiology of *Salmonella* Serovars in Animal-Based  
887 Foods: a Meta-analysis. *Applied and Environmental Microbiology*, 85, e00591-19,  
888 <https://doi.org/10.1128/AEM.00591-19>

889 Food and Agricultural Organisation of the United Nations (2018). Addressing the food safety  
890 challenges of the Asia-Pacific region. 34<sup>th</sup> Session of the FAO Regional Conference  
891 for Asia and the Pacific. Available from: <http://www.fao.org/3/MV819en/mv819en.pdf>  
892 Accessed 24.02.2021

893 Fu, H., Yu, P., Liang, W., Kan, B., Peng, X., & Chen, L. (2020). Virulence, Resistance, and  
894 Genomic Fingerprint Traits of *Vibrio cholerae* Isolated from 12 Species of Aquatic  
895 Products in Shanghai, China. *Microbial drug resistance* (Larchmont, N.Y.), 26(12),  
896 1526-1539. <https://doi.org/10.1089/mdr.2020.0269>

897 Ghosh, D. 2014. Food safety regulations in Australia and New Zealand Food Standards. *J Sci*  
898 *Food Agric* 94: 1970-1973

899 Jaffee, S., Henson, S., Unnevehr, L., Grace, D., Cassou, Emilie. (2018). The Safe Food  
900 Imperative: Accelerating Progress in Low- and Middle-Income Countries. Agriculture  
901 and Food Series; Washington, DC: World Bank. © World Bank. License: CC BY 3.0  
902 IGO. Available from: <https://openknowledge.worldbank.org/handle/10986/30568>  
903 Accessed 10.02.2021.

904 *Journal of Cleaner Production*, 284, 124709, <https://doi.org/10.1016/j.jclepro.2020.124709>.

905 Kaakoush, N. O., Castañó Rodríguez, N., Mitchell, H. M., & Man, S. M. (2015). Global  
906 epidemiology of *Campylobacter* infection. *Clinical microbiology reviews*, 28(3), 687-  
907 720.

908 Kantama, L. and Jayanetra, P. 1996. *Salmonella enteritidis* outbreak in Thailand: study by  
909 random amplified polymorphic DNA (RAPD) analysis. *Southeast Asian J Trop Med*  
910 *Public Health*, 27(1): 119-125.

911 Keisam, S., Tuikhar, N., Ahmed, G., and Jeyaram, K. 2019. Toxigenic and pathogenic  
912 potential of enteric bacterial pathogens prevalent in the traditional fermented foods  
913 marketed in the Northeast region of India. *International Journal of Food Microbiology*  
914 296: 21-30.

915 Kim, K., Kim, S., & Park, C. Y. (2020). Food Security in Asia and the Pacific amid the COVID-  
916 19 Pandemic.

917 Kowalska, A., & Manning, L. (2020). Using the rapid alert system for food and feed: potential  
918 benefits and problems on data interpretation. *Critical Reviews in Food Science and*  
919 *Nutrition*, 1-14.

920 Kumar R, Lalitha KV. Prevalence and molecular characterization of *Vibrio cholerae* O1, non-  
921 O1 and non-O139 in tropical seafood in Cochin, India. (2013). *Foodborne Pathog Dis*.  
922 10, 278-83. <https://doi.org/10.1089/fpd.2012.1310>.

923 Kumari, S. and Sarkar, P. K. 2014. Prevalence and characterization of *Bacillus cereus* group  
924 from various marketed dairy products in India. *Dairy Sci. & Technol.* 94: 483-497

925 Lee Y, Choi Y, Lee S, Lee H, Kim S, Ha J, Lee J, Oh H, Kim Y, Yoon Y. Occurrence of  
926 pathogenic *Vibrio parahaemolyticus* in seafood distribution channels and their  
927 antibiotic resistance profiles in S. Korea. *Lett Appl Microbiol.* 2019 Feb;68(2):128-  
928 133. <https://doi.org/10.1111/lam.13099>.

929 Lemoine, T., Germanetto, P. and Giraud, P. 1999. Toxi-infection alimentaire à *V*.  
930 *parahaemolyticus*. *Bull Epidemiol Hebdo*. 10: 37-38.

931 Machado-Moreira B, Richards K, Brennan F, Abram F, Burgess CM. (2019). Microbial  
932 Contamination of Fresh Produce: What, Where, and How? *Comprehensive Reviews in*  
933 *Food Science and Food Safety*, 6, 1727-1750. <https://doi.org/10.1111/1541-4337.12487>.

934 Makendi C, Page AJ, Wren BW, Le Thi Phuong T, Clare S, et al. 2016. A phylogenetic and  
935 phenotypic analysis of *Salmonella enterica* Serovar Weltevreden, an emerging agent  
936 of diarrheal disease in tropical regions. *PLOS Neglected Tropical Diseases* 10(2):  
937 e0004446.

938 Mangla, S.K., Bhattacharya, A., Yadav, A.K., Sharma, Y.K., Ishizaka, A., Luthra, S.,  
939 Chakraborty, R. (2021). A framework to assess the challenges to food safety  
940 initiatives in an emerging economy.

941 Martinez-Urtaza, J., Simental, L., Velasco, D., DePaola, A., Ishibashi, M., Nakaguchi, Y.,  
942 Nishibuchi, M., Carrera-Flores, D., Rey-Alvarez, C. and Pousa, A. 2005. Pandemic  
943 *Vibrio parahaemolyticus* O3:K6, Europe. *Emerg Infect Dis*. 11(8): 1319-1320.

944 McLauchlin, J., Aird, H., Andrews, N., Chattaway, M., de Pinna, E., Elviss, N., Jørgensen, F.,  
945 Larkin, L., Willis, C. (2019). Public health risks associated with *Salmonella*  
946 contamination of imported edible betel leaves: Analysis of results from England,  
947 2011-2017. *International Journal of Food Microbiology*, 298, 1-10,  
948 <https://doi.org/10.1016/j.ijfoodmicro.2019.03.004>.

949 McLauchlin, J., Aird, H., Andrews, N., Chattaway, M., de Pinna, E., Elviss, N., Jørgensen, F.,  
950 Larkin, L. and Willis, C. 2019. Public health risks associated with *Salmonella*  
951 contamination of imported edible betel leaves: Analysis of results from England,  
952 2011-2017. *International Journal of Food Microbiology* 298: 1-10

953 McLauchlin, J., Aird, H., Charlett, A., Chattaway, M., Elviss, N., Hartman, H., Jenkins, C.,  
954 Jørgensen, F., Larkin, L., Sadler-Reeves, L. and Willis, C. 2018. Imported edible  
955 leaves collected at retail sale in England during 2017 with an emphasis on betel and  
956 curry leaves: Microbiological quality with respect to *Salmonella*, Shiga toxin-  
957 producing *E. coli* (STEC) and levels of *Escherichia coli*. *J. Appl. Microbiol.*, 125:  
958 1175-1185

959

960 Nakaguchi, Y. (2013). Contamination by *Vibrio parahaemolyticus* and Its Virulent Strains in  
961 Seafood Marketed in Thailand, Vietnam, Malaysia, and Indonesia. *Tropical Medicine*  
962 *and Health*, 71, 95–102. <https://doi.org/10.2149/tmh.2011-06>

963 Nasheri, N., Vester, A. and Petronella, N. 2019. Foodborne viral outbreaks associated with  
964 frozen produce. *Epidemiology and Infection* 147: e291.

965 Nath, T.K., Inoue, M. (2009). Sustainability Attributes of a Small-Scale Betel-Leaf Agroforestry  
966 System: A Case Study in North-eastern Hill Forests of Bangladesh. *Small-scale*  
967 *Forestry*, 8, 289–304. <https://doi.org/10.1007/s11842-009-9084-4>.

968 Nguyen, T.K., Bui, H.T., Truong, T.A., Lam, D.N., Ikeuchi, S., Ly, L.K.T., Hara-Kudo, Y.,  
969 Taniguchi, T., Hayashidani, H. (2021). Retail fresh vegetables as a potential source  
970 of *Salmonella* infection in the Mekong Delta, Vietnam. *Int J Food Microbiol.*  
971 341:109049. <https://doi.org/10.1016/j.ijfoodmicro.2021.109049>

972 North China. *BMC Microbiology* 16: 32

973 Office of National Statistics, 2011. UK Census 2011.  
974 <https://www.ons.gov.uk/census/2011census>

975 Organisation for Economic Cooperation and Development (2020). Global value chains  
976 connect producers to consumers across the world Available from:  
977 <https://www.oecd.org/agriculture/topics/global-value-chains-and-agriculture/>

978 Ortega, D.L. and Tschirley, D.L. (2017). Demand for food safety in emerging and developing  
979 countries: A research agenda for Asia and Sub-Saharan Africa. *Journal of*  
980 *Agribusiness in Developing and Emerging Economies*, 7, 21–34.  
981 <https://doi.org/10.1108/JADEE-12-2014-0045>.

982 Park, S., Szonyi, B., Gautam, J., Nightingale, K., Anciso, J., Ivanek, R. (2012). Risk Factors for  
983 Microbial Contamination in Fruits and Vegetables at the Preharvest Level: A  
984 Systematic Review. *Journal of Food Protection*, 75, 2055–2081.  
985 <https://doi.org/10.4315/0362-028X-JFP-12-160>

986 Pham, H.V. and Dinh, T.L. (2020). The Vietnam's food control system: Achievements and  
987 remaining issues. *Food Control*, 108, 106862.  
988 <https://doi.org/10.1016/j.foodcont.2019.106862>.

989 Reddy, A. A., Ricart, S. and Cadman, T. 2020. Driving factors of food safety standards in  
990 India: learning from street food vendors' behaviour and attitude. *Food Security* 12:  
991 1201–1217

992 Rezaei, M., & Liu, B. (2017). Food loss and waste in the food supply chain. *International Nut*  
993 *and Dried Fruit Council: Reus, Spain*, 26–27.

994 Rivas, L., Strydom, H., Paine, S., Wang, J., & Wright, J. (2021). Yersiniosis in New Zealand.  
995 *Pathogens*, 10(2), 191.

996 Singh, B. R., Singh, M., Singh, P., Babu, N., Chandra, M., Agarwal, R. K. 2006. Prevalence of  
997 multidrug-resistant *Salmonella* on ready-to-eat betel leaves (Paan) and in water used  
998 for soaking betel leaves in North Indian cities. *J Food Prot.* 69(2): 288–292

999 Sodagari, H. R., Wang, P., Robertson, I., Habib, I., & Sahibzada, S. (2020). Non-Typhoidal  
1000 *Salmonella* at the Human Food-of-Animal-Origin Interface in Australia. *Animals*,  
1001 10(7), 1192.

1002 Somorin, Y. M., Odeyemi, O. A., & Ateba, C. N. (2021). *Salmonella* is the most common  
1003 foodborne pathogen in African food exports to the European Union: Analysis of Rapid  
1004 Alert System for Food and Feed (1999–2019). *Food Control*, 107849.  
1005 doi:<https://doi.org/10.1016/j.foodcont.2020.107849>

1006 Somorin, Y. M., Odeyemi, O. A., & Ateba, C. N. (2021). *Salmonella* is the most common  
1007 foodborne pathogen in African food exports to the European Union: Analysis of the  
1008 Rapid Alert System for Food and Feed (1999–2019). *Food Control*, 123, 107849.

1009 Somorin, Y.M., Odeyemi, O.A. and Ateba, C.N. 2021. *Salmonella* is the most common  
1010 foodborne pathogen in African food exports to the European Union: Analysis of Rapid  
1011 Alert System for Food and Feed (1999–2019). *Food Control* 123: 107849.

1012 Sugiri, Y. D., Geelz, G., Meeyam, T., Baumann, M. P., Kloor, J., Chaisowwong, W., & Alter,  
1013 T. (2014). Prevalence and antimicrobial susceptibility of *Listeria monocytogenes* on  
1014 chicken carcasses in Bandung, Indonesia. *Journal of food protection*, 77(8), 1407–  
1015 1410.

1016 Tähkääpää, S., Majjala, R., Korkeala, H., & Nevas, M. (2015). Patterns of food frauds and  
1017 adulterations reported in the EU rapid alert system for food and feed and in Finland.  
1018 *Food Control*, 47, 175–184.

1019 Ullah, A.S.M., Tani, M., Tsuchiya, J., Rahman, A.M., Rahman, Z.M. (2020). Impact of Betel  
1020 Leaf Cultivation on the Protected Forest Area of Teknaf Peninsula, Bangladesh.  
1021 *Small-scale Forestry*, 19, 335–355. <https://doi.org/10.1007/s11842-020-09441-w>

1022 Vilain, S., Luo, Y., Hildreth, M.B. and Brözel, V.S. 2006. Analysis of the life cycle of the soil  
1023 saprophyte *Bacillus cereus* in liquid soil extract and in soil. *Applied and  
1024 Environmental Microbiology* 72(7): 4970–4977.

1025 Vital, P.G., Dimasuay, G.B., Widmer, K.W., Rivera, W.L. (2014). Microbiological Quality of  
1026 Fresh Produce from Open Air Markets and Supermarkets in the Philippines. *The  
1027 Scientific World Journal*, Article ID 219534, <https://doi.org/10.1155/2014/219534>

1028 Wadamori, Y., Gooneratne, R., Hussain, M.A. (2016). Outbreaks and factors influencing  
1029 microbiological contamination of fresh produce. *Journal of the Science of Food and  
1030 Agriculture*, 97, 1396–1403. <https://doi.org/10.1002/jsfa.8125>

1031 Wang, S., Weller, D., Falardeau, J., Strawn, L. K., Mardones, F. O., Adell, A. D., & Switt, A. I.  
1032 M. (2016). Food safety trends: from globalization of whole genome sequencing to  
1033 application of new tools to prevent foodborne diseases. *Trends in Food Science &  
1034 Technology*, 57, 188–198.

1035 WHO. (2020). Food Safety. Retrieved from [https://www.who.int/news-room/fact-](https://www.who.int/news-room/fact-sheets/detail/food-safety)  
1036 [sheets/detail/food-safety](https://www.who.int/news-room/fact-sheets/detail/food-safety)

1037 Wong, H. C., Chen, M. C., Liu, S. H. and Liu, D. P. 1999. Incidence of highly genetically  
1038 diversified *Vibrio parahaemolyticus* in seafood imported from Asian countries.  
1039 *International Journal of Food Microbiology* 52: 181–188

1040 World Health Organisation (2020) *Salmonella* non-typhoidal. Available from:  
1041 [https://www.who.int/news-room/fact-sheets/detail/salmonella-\(non-typhoidal\)](https://www.who.int/news-room/fact-sheets/detail/salmonella-(non-typhoidal))  
1042 Accessed 25.02.2024.

1043 World Health Organization, 2015. WHO estimates of the global burden of foodborne diseases.  
1044 Foodborne Disease Burden Epidemiology Reference Group 2007–2015. World  
1045 Health Organization, Geneva.

1046 World Health Organization, 2020. Framework for action on food safety in the WHO South-East  
1047 Asia Region. New Delhi: World Health Organization, Regional Office for South-East  
1048 Asia.

1049 Worldometer. (2021). Current World Population. Retrieved from  
1050 <https://www.worldometers.info/world-population/#ref-1>

1051 Xu, X., Cheng, J., Wu, Q., Zhang, J. and Xie, T. 2016. Prevalence, characterization, and  
1052 antibiotic susceptibility of *Vibrio parahaemolyticus* isolated from retail aquatic  
1053 products in

1054 Yu, P., Zaleski, A., Li, Q., He, Y., Mapili, K., Pruden, A., Alvarez, P. J. J. and Stadler, L. B.  
1055 2018. Elevated levels of pathogenic indicator bacteria and antibiotic resistance genes  
1056 after Hurricane Harvey's flooding in Houston. *Environ. Sci. Technol. Lett.* 5(8): 481–  
1057 486

1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113

## References

- [Abatcha, M. G., Effarizah, M. E., & Rusul, G. \(2018\). Prevalence, antimicrobial resistance, resistance genes and class 1 integrons of Salmonella serovars in leafy vegetables, chicken carcasses and related processing environments in Malaysian fresh food markets. \*Food Control\*, 91, 170-180.](#)
- [Ananchaipattana, C., Hosotani, Y., Kawasaki, S., Pongswat, S., Latiful, B. M., Isobe, S., & Inatsu, Y. \(2012\). Bacterial contamination of soybean curd \(tofu\) sold in Thailand. \*Food Science and Technology Research\*, 18\(6\), 843-848.](#)
- [Bag, P. K., Nandi, S., Bhadra, R. K., Ramamurthy, T., Bhattacharya, S., Nishibuchi, M., Hamabata, T., Yamasaki, S., Takeda, Y., & Nair, G. B. \(1999\). Clonal diversity among recently emerged strains of \*Vibrio parahaemolyticus\* O3: K6 associated with pandemic spread. \*Journal of clinical microbiology\*, 37\(7\), 2354-2357.](#)
- [Banerjee, M., & Sarkar, P. K. \(2003\). Microbiological quality of some retail spices in India. \*Food Research International\*, 36\(5\), 469-474.](#)
- [Bangtrakulnonth, A., Pornreongwong, S., Pulsrikarn, C., Sawanpanyalert, P., Hendriksen, R. S., Wong, D. M. L. F., & Aarestrup, F. M. \(2004\). \*Salmonella\* serovars from humans and other sources in Thailand, 1993–2002. \*Emerging infectious diseases\*, 10\(1\), 131.](#)
- [Bernard, H., Faber, M., Wilking, H., Haller, S., Höhle, M., Schielke, A., Ducomble, T., Sifczyk, C., Merbecks, S., & Fricke, G. \(2014\). Large multistate outbreak of norovirus gastroenteritis associated with frozen strawberries, Germany, 2012. \*Eurosurveillance\*, 19\(8\), 20719.](#)
- [Bisht, A., Kamble, M. P., Choudhary, P., Chaturvedi, K., Kohli, G., Juneja, V. K., Sehgal, S., & Taneja, N. K. \(2021\). A surveillance of food borne disease outbreaks in India: 2009–2018. \*Food Control\*, 121, 107630.](#)
- [Borchers, A., Teuber, S. S., Keen, C. L., & Gershwin, M. E. \(2010\). Food safety. \*Clinical reviews in allergy & immunology\*, 39\(2\), 95-141.](#)
- [Briet, A., Helsens, N., Delannoy, S., Debuiche, S., Brisabois, A., Midelet, G., & Granier, S. \(2018\). NDM-1-producing \*Vibrio parahaemolyticus\* isolated from imported seafood. \*J Antimicrob Chemother\* doi, 10.](#)
- [Carstens, C. K., Salazar, J. K., & Darkoh, C. \(2019\). Multistate outbreaks of foodborne illness in the United States associated with fresh produce from 2010 to 2017. \*Frontiers in microbiology\*, 10, 2667.](#)
- [Chaoniruthisai, P., Punnakitikashem, P., & Rajchamaha, K. \(2018\). Challenges and difficulties in the implementation of a food safety management system in Thailand: A survey of BRC certified food productions. \*Food Control\*, 93, 274-282.](#)
- [Chen, A., Giese, M., & Chen, D. \(2020\). Flood impact on Mainland Southeast Asia between 1985 and 2018—The role of tropical cyclones. \*Journal of Flood Risk Management\*, 13\(2\), e12598.](#)
- [Chen, X., Zhu, Q., Yu, F., Zhang, W., Wang, R., Ye, X., Jin, L., Liu, Y., Li, S., & Chen, Y. \(2018\). Serology, virulence and molecular characteristics of \*Vibrio parahaemolyticus\* isolated from seafood in Zhejiang province. \*PLoS one\*, 13\(10\), e0204892.](#)
- [Cook, N., Williams, L., & D'Agostino, M. \(2019\). Prevalence of Norovirus in produce sold at retail in the United Kingdom. \*Food microbiology\*, 79, 85-89.](#)
- [DEFRA. \(2020\). Food Statistics in your pocket: Global and UK supply. Department for Environment Food & Rural Affairs.](#)
- [Dolan, C., & Humphrey, J. \(2000\). Governance and trade in fresh vegetables: the impact of UK supermarkets on the African horticulture industry. \*Journal of development studies\*, 37\(2\), 147-176.](#)

Formatted: Justified

Formatted: Font: (Default) Arial

Field Code Changed

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

1114 [Dos Santos, S. F., Cardoso, R. d. C. V., Borges, Í. M. P., e Almeida, A. C., Andrade, E. S.,](#)  
1115 [Ferreira, I. O., & do Carmo Ramos, L. \(2020\). Post-harvest losses of fruits and](#)  
1116 [vegetables in supply centers in Salvador, Brazil: Analysis of determinants, volumes](#)  
1117 [and reduction strategies. \*Waste Management\*, 101, 161-170.](#)

1118 [ECDC/EFSA. \(2021\). Multi-country outbreak of Salmonella Enteritidis sequence type \(ST\)11](#)  
1119 [infections linked to poultry products in the EU/EEA and the United Kingdom – 25](#)  
1120 [February 2021. Stockholm: ECDC/EFSA; 2021. European Centre for Disease](#)  
1121 [Prevention and Control \(ECDC\) and European Food Safety Authority \(EFSA\).](#)

1122 [EFSA/ECDC. \(2019\). Scientific report on the European Union one health 2018. Zoonoses](#)  
1123 [report. European Centre for Disease Prevention and Control \(ECDC\) and European](#)  
1124 [Food Safety Authority \(EFSA\). \*EFSA J.\*, 17, 5926.](#)

1125 [EFSA/ECDC. \(2021\). The European Union One Health 2019 Zoonoses Report. European](#)  
1126 [Food Safety Authority \(EFSA\) and European Centre for Disease Prevention and](#)  
1127 [Control \(ECDC\). \*EFSA Journal\* 19, 6406.](#)

1128 [EFSA/ECDC. \(2018\). Multi-country outbreak of Salmonella Agona infections possibly linked](#)  
1129 [to ready-to-eat food. European Food Safety Authority \(EFSA\) and European Centre](#)  
1130 [for Disease Prevention Control \(ECDC\). In: Wiley Online Library.](#)

1131 [Emberland, K., Ethelberg, S., Kuusi, M., Vold, L., Jensvoll, L., Lindstedt, B., Nygård, K., Kjelsø,](#)  
1132 [C., Torpdahl, M., & Sørensen, G. \(2007\). Outbreak of \*Salmonella\* Weltevreden](#)  
1133 [infections in Norway, Denmark and Finland associated with alfalfa sprouts, July-](#)  
1134 [October 2007. \*Weekly releases \(1997–2007\)\*, 12\(48\), 3321.](#)

1135 [European Commission. \(2014\). Commission implementing decision of 13 February 2014](#)  
1136 [suspending temporarily imports from Bangladesh of foodstuffs containing or](#)  
1137 [consisting of betel leaves \('Piper betle'\) Available from: \[https://eur-\]\(https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32014D0088&from=DA\)](#)  
1138 [lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32014D0088&from=DA](#)  
1139 [Accessed 25.02.2021.](#)

1140 [European Commission. \(2015\). Commission implementing regulation \(EU\) 2015/943 of 18](#)  
1141 [June 2015. \*Official Journal of the European Union\*\(154\), 8-9.](#)

1142 [European Commission \(2016\). RASFF — The Rapid Alert System for Food and Feed — 2015](#)  
1143 [annual report. Available from: \[https://op.europa.eu/en/publication-detail/-\]\(https://op.europa.eu/en/publication-detail/-/publication/98e04cfc-aa54-11e6-aab7-01aa75ed71a1/language-en/format-PDF/source-174743070\)](#)  
1144 [/publication/98e04cfc-aa54-11e6-aab7-01aa75ed71a1/language-en/format-](#)  
1145 [PDF/source-174743070](#) Accessed 19.02.2021.

1146 [European Commission. \(2020a\). Commission implementing regulation \(EU\) 2020/625 of 6](#)  
1147 [May 2020. \*Official Journal of the European Union\*\(63\), 13.](#)

1148 [European Commission. \(2020b\). Monitoring EU Agri-Food Trade: Developments in 2019.](#)  
1149 [Available from: \[https://ec.europa.eu/info/news/2019-record-year-eu-agri-food-trade-\]\(https://ec.europa.eu/info/news/2019-record-year-eu-agri-food-trade-2020-mar-27\_en#moreinfo\)](#)  
1150 [2020-mar-27\\_en#moreinfo](#) Accessed 12.02.2021.

1151 [European Food Safety Authority. \(2021\). The European Union One Health 2019 Zoonoses](#)  
1152 [Report. \*EFSA Journal\*, 19\(2\).](#)

1153 [Fakruddin, M., Sultana, R., Hossain, M. N., Rahaman, M. M., Islam, M. K., & Ahmed, M. M.](#)  
1154 [\(2017\). Occurrence of ingestion of \*Salmonella\* spp. in Betel leaf \(Piper betle L.\).](#)  
1155 [International Journal of Food Contamination, 4\(1\), 1-10.](#)

1156 [Ferrari, R. G., Rosario, D. K., Cunha-Neto, A., Mano, S. B., Figueiredo, E. E., & Conte-Junior,](#)  
1157 [C. A. \(2019\). Worldwide epidemiology of \*Salmonella\* serovars in animal-based foods:](#)  
1158 [a meta-analysis. \*Applied and environmental microbiology\*, 85\(14\).](#)

1159 [Food and Agricultural Organisation. \(2018\). Addressing the food safety challenges of the Asia-](#)  
1160 [Pacific region. 34th Session of the FAO Regional Conference for Asia and the Pacific.](#)  
1161 [Available from: <http://www.fao.org/3/MV819en/mv819en.pdf> Accessed 24.02.2021.](#)

1162 [Food and Agriculture Organization. \(2019\). Foodborne illnesses in Asia-Pacific 'needlessly'](#)  
1163 [sicken 275 million annually and threaten trade. Retrieved from](#)  
1164 [http://www.fao.org/asiapacific/news/detail-](#)  
1165 [events/en/c/1197008/#:~:text=In%20the%20Asia%2DPacific%20region,of%20Zero%20Hunger%20by%202030,.](#)

Formatted: Font: (Default) Cambria Math

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Cambria Math

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Cambria Math

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

1167 [Fu, H., Yu, P., Liang, W., Kan, B., Peng, X., & Chen, L. \(2020\). Virulence, Resistance, and](#)  
1168 [Genomic Fingerprint Traits of \*Vibrio cholerae\* Isolated from 12 Species of Aquatic](#)  
1169 [Products in Shanghai, China. \*Microbial Drug Resistance\*, 26\(12\), 1526-1539.](#)

1170 [Ghosh, D. \(2014\). Food safety regulations in Australia and New Zealand food standards.](#)  
1171 [\*Journal of the Science of Food and Agriculture\*, 94\(10\), 1970-1973.](#)

1172 [Jaffee, S., Henson, S., Unnevehr, L., Grace, D., & Cassou, E. \(2018\). \*The safe food\*](#)  
1173 [imperative: Accelerating progress in low-and middle-income countries: World Bank](#)  
1174 [Publications.](#)

1175 [Kaakoush, N. O., Castaño-Rodríguez, N., Mitchell, H. M., & Man, S. M. \(2015\). Global](#)  
1176 [epidemiology of \*Campylobacter\* infection. \*Clinical microbiology reviews\*, 28\(3\), 687-](#)  
1177 [720.](#)

1178 [Kantama, L., & Jayanetra, P. \(1996\). \*Salmonella enteritidis\* outbreak in Thailand: study by](#)  
1179 [random amplified polymorphic DNA \(RAPD\) analysis. \*The Southeast Asian journal of\*](#)  
1180 [tropical medicine and public health](#), 27(1), 119-125.

1181 [Keisam, S., Tuikhar, N., Ahmed, G., & Jeyaram, K. \(2019\). Toxigenic and pathogenic potential](#)  
1182 [of enteric bacterial pathogens prevalent in the traditional fermented foods marketed in](#)  
1183 [the Northeast region of India. \*International journal of food microbiology\*, 296, 21-30.](#)

1184 [Kim, K., Kim, S., & Park, C.-Y. \(2020\). Food Security in Asia and the Pacific amid the COVID-](#)  
1185 [19 Pandemic. 1-15.](#)

1186 [Kowalska, A., & Manning, L. \(2020\). Using the rapid alert system for food and feed: potential](#)  
1187 [benefits and problems on data interpretation. \*Critical reviews in food science and\*](#)  
1188 [nutrition](#), 1-14.

1189 [Kumar, R., & Lalitha, K. V. \(2013\). Prevalence and molecular characterization of \*Vibrio\*](#)  
1190 [cholerae O1, non-O1 and non-O139 in tropical seafood in Cochin, India. \*Foodborne\*](#)  
1191 [pathogens and disease](#), 10(3), 278-283.

1192 [Kumari, S., & Sarkar, P. K. \(2014\). Prevalence and characterization of \*Bacillus cereus\* group](#)  
1193 [from various marketed dairy products in India. \*Dairy science & technology\*, 94\(5\), 483-](#)  
1194 [497.](#)

1195 [Lee, Y., Choi, Y., Lee, S., Lee, H., Kim, S., Ha, J., Lee, J., Oh, H., Kim, Y., & Yoon, Y. \(2019\).](#)  
1196 [Occurrence of pathogenic \*Vibrio parahaemolyticus\* in seafood distribution channels](#)  
1197 [and their antibiotic resistance profiles in S. Korea. \*Letters in applied microbiology\*,](#)  
1198 [68\(2\), 128-133.](#)

1199 [Lemoine, T., Germanetto, P., & Giraud, P. \(1999\). Toxi-infection alimentaire à \*V.\*](#)  
1200 [parahaemolyticus. \*Bull Epidemiol Hebdo\*, 10, 37-38.](#)

1201 [Machado-Moreira, B., Richards, K., Brennan, F., Abram, F., & Burgess, C. M. \(2019\). Microbial](#)  
1202 [contamination of fresh produce: what, where, and how? \*Comprehensive reviews in\*](#)  
1203 [food science and food safety](#), 18(6), 1727-1750.

1204 [Makendi, C., Page, A. J., Wren, B. W., Le Thi Phuong, T., Clare, S., Hale, C., Goulding, D.,](#)  
1205 [Klemm, E. J., Pickard, D., & Okoro, C. \(2016\). A phylogenetic and phenotypic analysis](#)  
1206 [of \*Salmonella enterica\* serovar Weltevreden, an emerging agent of diarrheal disease](#)  
1207 [in tropical regions. \*PLoS neglected tropical diseases\*, 10\(2\), e0004446.](#)

1208 [Mangla, S. K., Bhattacharya, A., Yadav, A. K., Sharma, Y. K., Ishizaka, A., Luthra, S., &](#)  
1209 [Chakraborty, R. \(2021\). A framework to assess the challenges to food safety initiatives](#)  
1210 [in an emerging economy. \*Journal of Cleaner Production\*, 284, 124709.](#)

1211 [Martinez-Urtaza, J., Simental, L., Velasco, D., DePaola, A., Ishibashi, M., Nakaguchi, Y.,](#)  
1212 [Nishibuchi, M., Carrera-Flores, D., Rey-Alvarez, C., & Pousa, A. \(2005\). Pandemic](#)  
1213 [vibrio parahaemolyticus O3: K6, Europe. \*Emerging infectious diseases\*, 11\(8\), 1319.](#)

1214 [McLauchlin, J., Aird, H., Andrews, N., Chattaway, M., De Pinna, E., Elviss, N., Jørgensen, F.,](#)  
1215 [Larkin, L., & Willis, C. \(2019\). Public health risks associated with \*Salmonella\*](#)  
1216 [contamination of imported edible betel leaves: analysis of results from England, 2011-](#)  
1217 [2017. \*International journal of food microbiology\*, 298, 1-10.](#)

1218 [McLauchlin, J., Aird, H., Charlett, A., Chattaway, M., Elviss, N., Hartman, H., Jenkins, C.,](#)  
1219 [Jørgensen, F., Larkin, L., & Sadler-Reeves, L. \(2018\). Imported edible leaves collected](#)  
1220 [at retail sale in England during 2017 with an emphasis on betel and curry leaves:](#)  
1221 [microbiological quality with respect to \*Salmonella\*, Shiga-toxin-producing \*E. coli\*](#)

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Cambria Math

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Cambria Math

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Cambria Math

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Cambria Math

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

1222 [\(STEC\) and levels of \*Escherichia coli\*, \*Journal of applied microbiology\*, 125\(4\), 1175-](#)  
1223 [1185.](#)

1224 [Mensah, L. D., & Julien, D. \(2011\). Implementation of food safety management systems in the](#)  
1225 [UK. \*Food Control\*, 22\(8\), 1216-1225.](#)

1226 [Nakaguchi, Y. \(2013\). Contamination with \*Vibrio parahaemolyticus\* and its virulent strains in](#)  
1227 [seafood marketed in Thailand, Vietnam, Malaysia, and Indonesia from 2008 to 2011.](#)  
1228 [\*Tropical Medicine and Health\*, 2011-2006.](#)

1229 [Nasheri, N., Vester, A., & Petronella, N. \(2019\). Foodborne viral outbreaks associated with](#)  
1230 [frozen produce. \*Epidemiology & Infection\*, 147.](#)

1231 [Nath, T. K., & Inoue, M. \(2009\). Sustainability attributes of a small-scale betel leaf agroforestry](#)  
1232 [system: a case study in north-eastern hill forests of Bangladesh. \*Small-scale Forestry\*,](#)  
1233 [8\(3\), 289-304.](#)

1234 [Nguyen, T. K., Bui, H. T., Truong, T. A., Lam, D. N., Ikeuchi, S., Ly, L. K. T., Hara-Kudo, Y.,](#)  
1235 [Taniguchi, T., & Hayashidani, H. \(2021\). Retail fresh vegetables as a potential source](#)  
1236 [of \*Salmonella\* infection in the Mekong Delta, Vietnam. \*International journal of food\*](#)  
1237 [microbiology, 341, 109049.](#)

1238 [Office of National Statistics. \(2011\). UK Census 2011. Available online:](#)  
1239 [https://www.ons.gov.uk/census/2011census](#)

1240 [Ortega, D. L. a. T., D.L. \(2017\). Demand for food safety in emerging and developing countries:](#)  
1241 [A research agenda for Asia and Sub-Saharan Africa. \*Journal of Agribusiness in\*](#)  
1242 [Developing and Emerging Economies\(7\), 21-34.](#)

1243 [Park, S., Szonyi, B., Gautam, R., Nightingale, K., Anciso, J., & Ivanek, R. \(2012\). Risk factors](#)  
1244 [for microbial contamination in fruits and vegetables at the preharvest level: a](#)  
1245 [systematic review. \*Journal of food protection\*, 75\(11\), 2055-2081.](#)

1246 [Pham, H. V., & Dinh, T. L. \(2020\). The Vietnam's food control system: Achievements and](#)  
1247 [remaining issues. \*Food Control\*, 108, 106862.](#)

1248 [Reddy, A. A., Ricart, S., & Cadman, T. \(2020\). Driving factors of food safety standards in India:](#)  
1249 [learning from street-food vendors' behaviour and attitude. \*Food Security\*, 1-17.](#)

1250 [Rezaei, M., & Liu, B. \(2017\). Food loss and waste in the food supply chain. \*International Nut\*](#)  
1251 [and Dried Fruit Council: Reus, Spain, 26-27.](#)

1252 [Rivas, L., Strydom, H., Paine, S., Wang, J., & Wright, J. \(2021\). Yersiniosis in New Zealand.](#)  
1253 [\*Pathogens\*, 10\(2\), 191.](#)

1254 [Singh, B. R., Singh, M., Singh, P., Babu, N., Chandra, M., & Agarwal, R. K. \(2006\). Prevalence](#)  
1255 [of multidrug-resistant \*Salmonella\* on ready-to-eat betel leaves \(Paan\) and in water](#)  
1256 [used for soaking betel leaves in North Indian cities. \*Journal of food protection\*, 69\(2\),](#)  
1257 [288-292.](#)

1258 [Sodagari, H. R., Wang, P., Robertson, I., Habib, I., & Sahibzada, S. \(2020\). Non-Typhoidal](#)  
1259 [Salmonella at the Human-Food-of-Animal-Origin Interface in Australia. \*Animals\*, 10\(7\),](#)  
1260 [1192.](#)

1261 [Somorin, Y. M., Odeyemi, O. A., & Ateba, C. N. \(2021\). \*Salmonella\* is the most common](#)  
1262 [foodborne pathogen in African food exports to the European Union: Analysis of the](#)  
1263 [Rapid Alert System for Food and Feed \(1999–2019\). \*Food Control\*, 123, 107849.](#)

1264 [Sugiri, Y. D., Goelz, G., Meeyam, T., Baumann, M. P., Kleer, J., Chaisowwong, W., & Alter,](#)  
1265 [T. \(2014\). Prevalence and antimicrobial susceptibility of \*Listeria monocytogenes\* on](#)  
1266 [chicken carcasses in Bandung, Indonesia. \*Journal of food protection\*, 77\(8\), 1407-](#)  
1267 [1410.](#)

1268 [Tähkääpää, S., Majjala, R., Korkeala, H., & Nevas, M. \(2015\). Patterns of food frauds and](#)  
1269 [adulterations reported in the EU rapid alert system for food and feed and in Finland.](#)  
1270 [\*Food Control\*, 47, 175-184.](#)

1271 [Ullah, A. S., Tani, M., Tsuchiya, J., Rahman, A. M., & Rahman, Z. M. \(2020\). Impact of Betel](#)  
1272 [Leaf Cultivation on the Protected Forest Area of Teknaf Peninsula, Bangladesh. \*Small-\*](#)  
1273 [scale Forestry, 19\(3\), 335-355.](#)

1274 [Vilain, S., Luo, Y., Hildreth, M. B., & Brözel, V. S. \(2006\). Analysis of the life cycle of the soil](#)  
1275 [saprophyte \*Bacillus cereus\* in liquid soil extract and in soil. \*Applied and environmental\*](#)  
1276 [microbiology, 72\(7\), 4970-4977.](#)

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

1277 [Vital, P. G., Dimasuay, K. G. B., Widmer, K. W., & Rivera, W. L. \(2014\). Microbiological quality](#)  
1278 [of fresh produce from open air markets and supermarkets in the Philippines. \*The\*](#)  
1279 [Scientific World Journal, 2014.](#)

1280 [Wadamori, Y., Gooneratne, R., & Hussain, M. A. \(2017\). Outbreaks and factors influencing](#)  
1281 [microbiological contamination of fresh produce. \*Journal of the Science of Food and\*](#)  
1282 [Agriculture, 97\(5\), 1396-1403.](#)

1283 [Wang, S., Weller, D., Falardeau, J., Strawn, L. K., Mardones, F. O., Adell, A. D., & Switt, A. I.](#)  
1284 [M. \(2016\). Food safety trends: from globalization of whole genome sequencing to](#)  
1285 [application of new tools to prevent foodborne diseases. \*Trends in Food Science &\*](#)  
1286 [Technology, 57, 188-198.](#)

1287 [Wong, H.-C., Chen, M.-C., Liu, S.-H., & Liu, D.-P. \(1999\). Incidence of highly genetically](#)  
1288 [diversified \*Vibrio parahaemolyticus\* in seafood imported from Asian countries.](#)  
1289 [International journal of food microbiology, 52\(3\), 181-188.](#)

1290 [World Health Organization. \(2015\). WHO estimates of the global burden of foodborne](#)  
1291 [diseases: foodborne disease burden epidemiology reference group 2007-2015; World](#)  
1292 [Health Organization.](#)

1293 [World Health Organization. \(2019\). Shiga Toxin-producing Escherichia Coli \(STEC\) and Food:](#)  
1294 [Attribution Characterization and Monitoring \(Vol. 19\): World Health Organization.](#)

1295 [World Health Organization. \(2020a\). Food Safety. Retrieved from \[https://www.who.int/news-\]\(https://www.who.int/news-room/fact-sheets/detail/food-safety\)](#)  
1296 [room/fact-sheets/detail/food-safety.](#)

1297 [World Health Organization. \(2020b\). Framework for action on food safety in the WHO South-](#)  
1298 [East Asia Region. New Delhi: World Health Organization, Regional Office for South-](#)  
1299 [East Asia.](#)

1300 [World Health Organization. \(2020c\). \*Salmonella\* – non-typhoidal. Available from:](#)  
1301 [https://www.who.int/news-room/fact-sheets/detail/salmonella-\(non-typhoidal\)](#)  
1302 [Accessed 25.02.2021.](#)

1303 [Worldometer. \(2021\). Current World Population. Retrieved from](#)  
1304 [https://www.worldometers.info/world-population/#ref-1.](#)

1305 [Xu, X., Cheng, J., Wu, Q., Zhang, J., & Xie, T. \(2016\). Prevalence, characterization, and](#)  
1306 [antibiotic susceptibility of \*Vibrio parahaemolyticus\* isolated from retail aquatic products](#)  
1307 [in North China. \*BMC microbiology, 16\*\(1\), 1-9.](#)

1308 [Yu, P., Zaleski, A., Li, Q., He, Y., Mapili, K., Pruden, A., Alvarez, P. J., & Stadler, L. B. \(2018\).](#)  
1309 [Elevated levels of pathogenic indicator bacteria and antibiotic resistance genes after](#)  
1310 [Hurricane Harvey's flooding in Houston. \*Environmental science & technology letters,\*](#)  
1311 [5\(8\), 481-486.](#)

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Not Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Not Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

1313 **Supplementary Table 1**

1314 RASFF notifications of pathogenic microorganisms in food originating from the Asia-Pacific Region (n=1873) from 2020 - 2020 based on  
 1315 notifying country.

Pathogen genus	Notifying country											Total (%)
	United Kingdom	Italy	Norway	Netherlands	Germany	Finland	Greece	Spain	Denmark	Sweden	Others	
<i>Salmonella</i>	515	95	102	172	137	122	80	47	30	44	216	1560(73.55%)
<i>Vibrio</i>	4	87	124	4	10	7	2	8	24	12	27	309(14.57%)
<i>Bacillus</i>	36	19	1	3	13	9	0	0	0	0	11	92(4.34%)
<i>Escherichia</i>	27	2	8	1	1	1	0	0	2	1	4	47(2.22%)
<i>Norovirus</i>	0	4	1	1	3	1	0	20	3	1	8	42(1.98%)
<i>Not specified</i>	2	11	0	1	1	0	0	0	1	5	7	28(1.32%)
<i>Clostridium</i>	2	18	0	0	1	0	0	0	0	0	1	22(1.04%)
<i>Campylobacter</i>	0	0	0	0	0	0	0	0	4	1	1	6(0.28%)
<i>Staphylococcus</i>	0	0	2	0	0	0	0	0	1	0	0	3(0.14%)
<i>Enterococcus</i>	0	1	0	0	0	0	0	0	1	0	0	2(0.09%)
<i>Hepatitis</i>	0	1	0	0	0	0	0	0	0	0	1	2(0.09%)
<i>Pseudomonas</i>	0	0	0	0	2	0	0	0	0	0	0	2(0.09%)
<i>Cronobacter</i>	0	0	0	1	0	0	0	0	0	0	0	1(0.05%)
Fungi	0	0	0	0	0	0	0	0	0	0	1	1(0.05%)
<i>Plesiomonas</i>	1	0	0	0	0	0	0	0	0	0	0	1(0.05%)
<i>Shigella</i>	0	0	0	0	0	0	0	0	1	0	0	1(0.05%)
<i>Yersinia</i>	0	1	0	0	0	0	0	0	0	0	0	1(0.05%)
<i>Streptococci</i>	0	0	0	0	0	1	0	0	0	0	0	1(0.05%)
<b>Total</b>	<b>587</b>	<b>239</b>	<b>238</b>	<b>183</b>	<b>168</b>	<b>141</b>	<b>82</b>	<b>75</b>	<b>67</b>	<b>64</b>	<b>277</b>	<b>2121</b>

1316 **Other\_N: Other notifying countries:** Cyprus, Austria, Slovenia, Ireland, Lithuania, Iceland, Romania, Switzerland, Bulgaria, Croatia, Estonia,  
 1317 Hungary, Latvia, Luxembourg, Slovakia, Sri Lanka. **Other\_O: Other originating countries:** Australia, Cambodia, New Zealand, Pakistan,  
 1318 Philippines, Hong Kong, Taiwan, Singapore, South Korea, Afghanistan, Nepal. The total number of pathogens (n=2121) exceeds the total  
 1319 number of notifications (n=1873) because of the co-occurrence of pathogens.

1320 **Supplementary Table 2**

1321 Pathogens associated with notifications and their diversities in Crustaceans and related

1322 products from Asia and Pacific (2000 - 2020).

Country of origin	Notifications (n)	Pathogens	No of pathogens	Crustaceans and related products
Bangladesh	38	<i>Salmonella</i> Bareilly, <i>Salmonella</i> Brunei, <i>Salmonella</i> enterica, <i>Salmonella</i> spp, <i>Vibrio cholerae</i> , <i>Vibrio parahaemolyticus</i>	43	Shrimps
		<i>Salmonella</i> spp, <i>Vibrio cholerae</i> , <i>Vibrio parahaemolyticus</i>	5	Prawns
China	12	<i>Salmonella</i> enterica ser. Newport, <i>Vibrio mimicus</i> , <i>Vibrio parahaemolyticus</i> , <i>Vibrio cholerae</i>	9	Crayfish
		<i>Salmonella</i> spp	2	Crab
		<i>Vibrio parahaemolyticus</i>	1	Shrimps
India	49	aerobic mesophiles, <i>Enterococcus</i> spp, <i>Pseudomonas</i> spp, <i>Salmonella</i> enterica, <i>Salmonella</i> spp, <i>Salmonella</i> Weltevreden, <i>Vibrio alginolyticus</i> , <i>Vibrio cholerae</i> , <i>Vibrio fluvialis</i> , <i>Vibrio parahaemolyticus</i> , <i>Vibrio vulnificus</i>	60	Shrimps
		<i>Salmonella</i> enterica, <i>Salmonella</i> spp, <i>Vibrio cholerae</i> , <i>Vibrio parahaemolyticus</i>	7	Prawns
		<i>Vibrio cholerae</i>	1	Crab
Indonesia	26	<i>Salmonella</i> enterica subsp. houtenae (VI), <i>Salmonella</i> Lexington, <i>Salmonella</i> spp, <i>Salmonella</i> Wandsworth, <i>Vibrio alginolyticus</i> , <i>Vibrio cholerae</i> , <i>Vibrio parahaemolyticus</i> , <i>Vibrio</i> spp, <i>Vibrio vulnificus</i> ,	21	Shrimps
		<i>Plesiomonas shigelloides</i> , <i>Salmonella</i> spp, <i>Vibrio parahaemolyticus</i> , <i>Vibrio cholerae</i>	12	Prawns
Malaysia	65	<i>Salmonella</i> enterica, <i>Salmonella</i> Give, <i>Salmonella</i> spp, <i>Vibrio cholerae</i> , <i>Vibrio parahaemolyticus</i> , <i>Vibrio vulnificus</i> ,	48	Shrimps
		<i>Salmonella</i> enterica, <i>Salmonella</i> spp, <i>Vibrio cholerae</i> , <i>Vibrio parahaemolyticus</i>	17	Prawns
		<i>Vibrio parahaemolyticus</i>	1	shellfish
Sri Lanka, Pakistan	4	<i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	4	Crab
		<i>Vibrio cholerae</i> , aerobic mesophiles, <i>Salmonella</i> spp	3	Shrimps
Thailand	23	<i>Salmonella</i> enterica, <i>Vibrio cholerae</i> , <i>Vibrio parahaemolyticus</i> , <i>Vibrio vulnificus</i>	21	Shrimps
		<i>Clostridium perfringens</i> , <i>Clostridium</i> sulphite reducer	3	Shrimp paste
		<i>Vibrio cholerae</i>	1	Crab
		<i>Vibrio parahaemolyticus</i>	1	Prawns

Country of origin	Notifications (n)	Pathogens	No of pathogens	Crustaceans and related products
Vietnam	48	<i>Escherichia coli</i> , <i>Salmonella enterica</i> , <i>Salmonella enterica ser. Weltevreden</i> , <i>Salmonella Kentucky</i> , <i>Salmonella Newport</i> , <i>Salmonella Oranienburg</i> , <i>Salmonella spp.</i> , <i>Salmonella Virchow</i> , <i>Salmonella Welshimeri</i> , <i>Salmonella Weltevreden</i> , <i>Vibrio cholerae</i> , <i>Vibrio parahaemolyticus</i> , <i>Vibrio spp.</i> , <i>Vibrio vulnificus</i>	55	Shrimps
		<i>Salmonella enterica</i> , <i>Salmonella spp.</i> , <i>Vibrio cholerae</i> , <i>Vibrio parahaemolyticus</i>	6	Prawns
		<i>Clostridium sulphite reducer</i>	1	Shrimp paste
		<i>Salmonella enterica ser. Hvitvingfoss</i>	1	Crab
		<i>Vibrio parahaemolyticus</i>	1	Clams
		<i>Salmonella enterica</i>	1	Crustaceans

- Formatted: Font: Not Italic
- Formatted: Font: Not Italic
- Formatted: Font: Not Italic
- Formatted: Font: Not Italic
- Formatted: Font: Not Italic
- Formatted: Font: Not Italic
- Formatted: Font: Not Italic
- Formatted: Font: Not Italic
- Formatted: Font: Not Italic
- Formatted: Font: Not Italic

1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343

1344 **Supplementary Table 3**

1345 Pathogens associated with notifications and their diversities in Fruits and Vegetables from

1346 Asia and Pacific (2000 - 2020).

Country of origin	No of notifications	Pathogen	Count of Pathogen species	Food	
Afghanistan	1	<i>Salmonella enterica ser-ser. Agona</i>	1	mulberry	Formatted
Australia	1	<i>Salmonella enterica ser-ser. Chester</i>	1	mushroom	Formatted
Bangladesh	151	<i>Salmonella enterica, Salmonella Jerusalem, Salmonella spp, Salmonella Typhimurium, Salmonella Virchow</i>	149	betel leaves	Formatted
Cambodia	6	<i>Salmonella enterica</i>	3	paan leaves	Formatted: Font: Italic
		<i>Salmonella spp</i>	2	Basil	Formatted: Font: Italic
		<i>Salmonella enterica, Salmonella enterica ser-ser. Stanley</i>	2	betel leaves	Formatted
Cambodia	6	<i>Salmonella Hvittingfoss, Salmonella Brunei, Salmonella enteritidis, Salmonella Typhimurium, Salmonella Weltevreden</i>	5	vine leaves	Formatted
		<i>Salmonella spp</i>	1	mushroom	Formatted: Font: Italic
Chile	1	<i>Salmonella spp</i>	1	mushroom	Formatted: Font: Italic
China	39	<i>Salmonella enteritidis</i>	1	beans	Formatted
				sprouts	
		<i>Escherichia coli, Salmonella spp</i>	2	Cep powde	Formatted: Font: Italic
		<i>Salmonella Thompson</i>	1	Chanterelle powder	Formatted
		<i>Bacillus cereus</i>	1	Cheese	Formatted: Font: Italic
		<i>Salmonella Rissen</i>	1	chlorella	Formatted
		<i>Salmonella spp</i>	1	goji berries	Formatted: Font: Italic
		<i>Bacillus cereus, Clostridium perfringens, Clostridium sporulated sulphite reducer, Clostridium spp, Salmonella spp, aerobic mesophiles, moulds, Salmonella enterica</i>	18	mushroom	Formatted: Font: Italic
		<i>Norovirus, Norovirus-Norovirus GII</i>	5	raspberries	Formatted
		<i>Norovirus-Norovirus GI-GI &amp; GII</i>	2	seaweed salad	Formatted
		<i>Bacillus cereus</i>	1	soy cheese	Formatted: Font: Italic
		<i>Salmonella Tennessee</i>	1	soy protein	Formatted
		Hepatitis A, Norovirus, Norovirus, GI-GI & GII	5	strawberries	Formatted
		<i>Bacillus cereus</i>	2	tofu	Formatted: Font: Italic
Hong Kong	2	<i>Salmonella Senftenberg</i>	1	mushrooms	Formatted: Font: Italic
		<i>Bacillus cereus</i>	1	tofu	Formatted: Font: Italic
India	192	<i>Salmonella enterica, Salmonella spp</i>	184	betel leaves	Formatted: Font: Italic
		<i>Salmonella enterica</i>	2	curry leaves	Formatted: Font: Italic
		<i>Salmonella spp</i>	1	fennel seed	Formatted: Font: Italic
		<i>Salmonella spp</i>	1	moringa leaves	Formatted: Font: Italic
		<i>Salmonella agona, Salmonella Westminster</i>	2	onions	Formatted: Font: Italic
		<i>Salmonella infantis</i>	1	sorrel leaves	Formatted: Font: Italic

Country of origin	No of notifications	Pathogen	Count of Pathogen species	Food	
Laos	1	<i>Escherichia coli</i> , <i>Salmonella Brunei</i> , <i>Salmonella Weltevreden</i>	3	vine leaves	Formatted: Font: Italic
Malaysia	15	<i>Salmonella enterica ser-ser. Aberdeen</i>	1	beans	Formatted: Font: Italic
		<i>Escherichia coli</i> , <i>Salmonella Colindale</i> , <i>Salmonella group D</i> , <i>Salmonella spp</i> , <i>Salmonella Weltevreden</i> , <i>Vibrio cholerae</i>	14	betel leaves	Formatted: Font: Italic
Pakistan	1	<i>Salmonella Weltevreden</i> , <i>Vibrio cholerae</i>	2	Alfalfa	Formatted: Font: Italic
		<i>Escherichia coli</i> , <i>Salmonella enterica</i> , <i>Salmonella enterica ser-ser. Weltevreden</i> , <i>Salmonella spp</i>	5	betel leaves	Formatted: Font: Italic
Sri Lanka	7	<i>Salmonella Weltevreden</i> , <i>Vibrio cholerae</i>	2	centella	Formatted: Font: Italic
		<i>Salmonella poona</i> , <i>Salmonella Weltevreden</i> , <i>Vibrio cholerae</i>	3	Spinach	Formatted: Font: Italic
		<i>Bacillus cereus</i>	1	tofu	Formatted: Font: Italic
Taiwan	1	<i>Bacillus cereus</i>	1	tofu	Formatted: Font: Italic
Thailand	81	<i>Salmonella anatum</i> , <i>Escherichia coli</i> , <i>Salmonella Stanley</i>	3	magosa	Formatted: Font: Italic
		<i>Salmonella spp</i> , <i>Salmonella Virchow</i>	2	Acacia	Formatted: Font: Italic
		<i>Campylobacter spp</i> , <i>Salmonella Bareilly</i>	2	Asparagus	Formatted: Font: Italic
		<i>Salmonella spp</i> , <i>Salmonella Weltevreden</i> , <i>Vibrio cholerae</i>	3	Banana leaves	Formatted: Font: Italic
		<i>Escherichia coli</i> , <i>Salmonella enterica</i> , <i>Salmonella Jerusalem</i> <i>Salmonella spp</i>	16	betel leaves	Formatted: Font: Italic
		<i>Bacillus cereus</i> , <i>Salmonella Weltevreden</i> , <i>Vibrio cholerae</i>	3	black fungus	Formatted: Font: Italic
		<i>Salmonella Thompson</i> , <i>Salmonella spp</i>	1	broccoli	Formatted: Font: Italic
		<i>Salmonella spp</i>	3	Cabbage, Cha leaves	Formatted: Font: Italic
		<i>Campylobacter spp</i> , <i>Salmonella enterica</i> , <i>Salmonella spp</i> , <i>Salmonella typhimurium</i> , <i>Salmonella Weltevreden</i> , <i>Salmonella Zanzibar</i> , <i>Shigella sonnei</i> , <i>Vibrio cholerae</i>	8	Corn	Formatted: Font: Italic
		<i>Salmonella spp</i>	2	fresh lime and banana, Ipomea aquatica	Formatted: Font: Italic
		<i>Salmonella Stanley</i>	1	kaffir lime leaves	Formatted: Font: Italic
		<i>Salmonella Hvittingfoss</i> , <i>Salmonella infantis</i> , <i>Salmonella Zanzibar</i>	3	lemon grass	Formatted: Font: Italic
		<i>Salmonella Hadar</i> , <i>Salmonella Lexington</i>	2	lime leaves	Formatted: Font: Italic
		<i>Salmonella enterica</i> , <i>Salmonella enterica ser-ser. Hvittingfoss</i> , <i>Salmonella enterica ser-ser. Newport</i> , <i>Salmonella spp</i> , <i>Salmonella Stanley</i>	6	morning glory	Formatted: Font: Italic
		<i>Salmonella Hvittingfoss</i> , <i>Salmonella spp</i>	2	mushroom	Formatted: Font: Italic
		<i>Salmonella enterica ser-ser. Hvittingfoss</i>	1	neem leaves	Formatted: Font: Italic

Country of origin	No of notifications	Pathogen	Count of Pathogen species	Food			
		<i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	2	okra	Formatted		
		<i>Salmonella</i> spp	1	ong choi	Formatted: Font: Italic		
		<i>Salmonella</i> bBraenderup	1	Other vegetables	Formatted		
		<i>Salmonella enterica</i>	1	paan leave	Formatted: Font: Italic		
		<i>Salmonella</i> aAltona, <i>Salmonella</i> Stanley	2	pak peaw	Formatted		
		<i>Salmonella</i> Brunei	1	pak wan	Formatted		
		<i>Salmonella enterica</i> ser-ser. Chester, <i>Salmonella enterica</i> ser-ser. Rubislaw, <i>Salmonella enterica</i> subsp. Salamae	3	pandang leaf	Formatted		
		<i>Salmonella</i> Ndolo, <i>Salmonella</i> Aberdeen, <i>Salmonella</i> bBovismorbificans, <i>Salmonella</i> enterica, <i>Salmonella</i> group B, <i>Salmonella</i> group D, <i>Salmonella</i> Hvittingfoss, <i>Salmonella</i> Mbandaka, <i>Salmonella</i> Newport	12	Spinach	Formatted: Font: Italic		
		<i>Campylobacter</i> spp, <i>Salmonella enterica</i> , <i>Salmonella</i> Hvittingfoss, <i>Salmonella</i> spp	5	spring onions	Formatted		
		<i>Salmonella</i> aAnatum	1	tamarind leaves	Formatted		
		<i>Salmonella</i> aAugustenborg	1	watercress	Formatted		
		<i>Salmonella</i> Saint Paul, <i>Salmonella</i> Stanley	2	wildbetal leafbush	Formatted		
		Vietnam	38	<i>Bacillus cereus</i> , <i>Salmonella</i> aAugustenborg, <i>Salmonella enterica</i> ser-ser. Stanley, <i>Salmonella</i> spp, <i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	8	black fungus	Formatted
				<i>Salmonella</i> Brunei	1	celery	Formatted
				<i>Salmonella</i> eEnterica	1	coriander	Formatted
<i>Salmonella</i> group B	1			morels	Formatted		
<i>Salmonella</i> spp	1			morning glory	Formatted		
<i>Bacillus cereus</i> , <i>Salmonella</i> Thompson, <i>Clostridium perfringens</i> , <i>Salmonella enterica</i> , <i>Salmonella enterica</i> ser-ser. Derby, <i>Salmonella enterica</i> ser-ser. Panama, <i>Salmonella enterica</i> ser-ser. Stanley, <i>Salmonella enterica</i> ser-ser. Virchow, <i>Salmonella</i> group B, <i>Salmonella</i> Java, <i>Salmonella</i> Meleagridis, <i>Salmonella</i> Rissen, <i>Salmonella</i> spp, <i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	29			mushroom	Formatted		
<i>Salmonella</i> Stanley	1			okra	Formatted		
<i>Campylobacter</i> spp	1			peppers	Formatted		
<i>Salmonella</i> spp	1			pineapple	Formatted		
<i>Salmonella</i> spp	1			Spinach	Formatted		

Country of origin	No of notifications	Pathogen	Count of Pathogen species	Food
		<i>Salmonella</i> Lexington, <i>Salmonella</i> Weltevreden	2	coriander, houttuynia and celery

Formatted

1347 Supplementary Table 4

1348 Pathogens associated with notifications and their diversities in nuts and related products  
 1349 from Asia and Pacific (2000 - 2020).

Country of origin	Notifications (n)	Pathogens	No of pathogens	Nuts and related products
China	7	<i>Salmonella</i> Newport, <i>Salmonella</i> Stanley	2	peanuts
		<i>Salmonella</i> spp.	1	pine nuts
		<i>Bacillus cereus</i>	1	pumpkin seeds
		<i>Bacillus cereus</i>	1	soy cheese
		<i>Salmonella</i> Senftenberg, <i>Salmonella</i> spp., <i>Salmonella</i> Tennessee	3	sunflower seeds
India	253	<i>Salmonella</i> spp.	1	amaranth
		<i>Salmonella enterica</i> ser. Matadi	1	cashew
		<i>Salmonella</i> eElomrane, <i>Salmonella enterica</i> ser. Bareilly	2	nigella seeds
		<i>Escherichia coli</i> , <i>Salmonella</i> spp.	2	Coconut
		<i>Bacillus cereus</i> , <i>Escherichia coli</i> , <i>Salmonella</i> #Mbandaka, <i>Salmonella</i> Montevideo, <i>Salmonella</i> spp., <i>Salmonella</i> aAgona, <i>Salmonella</i> Amsterdam, <i>Salmonella</i> aAnatum, <i>Salmonella</i> Bareilly, <i>Salmonella</i> bBinza, <i>Salmonella</i> Bbraenderup, <i>Salmonella</i> Chittagong, <i>Salmonella</i> dDallgow, <i>Salmonella</i> dDrypool, <i>Salmonella enterica</i> , <i>Salmonella enterica</i> ser. Agona, <i>Salmonella enterica</i> ser. Amsterdam, <i>Salmonella enterica</i> ser. Bareilly, <i>Salmonella enterica</i> ser. Braenderup, <i>Salmonella enterica</i> ser. Derby, <i>Salmonella enterica</i> ser. Hvittingfoss, <i>Salmonella enterica</i> ser. Isangi, <i>Salmonella enterica</i> ser. Kentucky, <i>Salmonella enterica</i> ser. Kisil, <i>Salmonella enterica</i> ser. Livingstone, <i>Salmonella enterica</i> ser. Londo, <i>Salmonella enterica</i> ser. Montevideo, <i>Salmonella enterica</i> ser. Orion, <i>Salmonella enterica</i> ser. Ouakam, <i>Salmonella enterica</i> ser. Pensacola, <i>Salmonella enterica</i> ser. Schwarzengrund, <i>Salmonella enterica</i> ser. Senftenberg, <i>Salmonella enterica</i> ser. Typhimurium, <i>Salmonella enterica</i> ser. Umbadah, <i>Salmonella</i> Eenteritidis, <i>Salmonella</i> eGaminara, <i>Salmonella</i> group C1, <i>Salmonella</i> Hvittingfoss	262	Sesame seeds

Formatted

Formatted

Formatted: Font: Italic

Formatted: Font: Italic

Formatted

Formatted: Font: Italic

Formatted

Formatted

Formatted

Formatted

Formatted



1360 **Supplementary Table 5**

1361 Yearly RASFF notifications of pathogenic microorganisms in nuts, nut products and seeds originating from Asia-Pacific Region (n=1873).

Year	Nuts, nut products and seeds					Sesame seeds		
	India	Indonesia	Thailand	Others	Total	India	Thailand	Total
2000	0	0	0	1	1	0	0	0
2001	3	0	1	2	6	3	1	4
2002	0	0	0	2	2	0	0	0
2003	1	0	0	2	3	1	0	1
2004	6	1	0	0	7	6	0	6
2005	10	1	0	1	12	10	0	10
2006	1	1	0	0	2	1	0	1
2007	4	0	0	1	5	4	0	4
2008	11	0	0	2	13	11	0	11
2009	13	0	0	2	15	13	0	13
2010	8	0	0	1	9	8	0	8
2011	6	0	0	0	6	6	0	6
2012	8	2	0	2	12	8	0	8
2013	5	0	0	2	7	5	0	5
2014	24	2	0	1	27	24	0	24
2015	68	1	0	1	70	67	0	67
2016	19	0	0	0	19	19	0	19
2017	20	1	0	0	21	18	0	18
2018	24	0	0	3	27	23	0	23
2019	17	0	0	0	17	17	0	17
2020	18	0	0	0	18	16	0	16
Total	266	9	1	23	299	260	1	261

1362 Others: Malaysia, Pakistan, Phillipines, Singapore, Sri Lanka, China, Vietnam



Country of origin	Notifications (n)	Pathogens	No of pathogens	Herbs and spices	
		<i>Bacillus cereus</i> , <i>Clostridium perfringens</i> , <i>Salmonella</i> Ohio	4	ginger powder	Formatted: Font: Italic
		<i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	2	ginseng powder	Formatted: Font: Italic
		<i>Bacillus cereus</i> , <i>Salmonella</i> group C, <i>Salmonella</i> spp.	3	ground cur	Formatted: Font: Italic
		<i>Salmonella</i> Agona, <i>Salmonella</i> Bareilly, <i>Salmonella</i> Cubana	3	ground cur and ground coriander	Formatted: Font: Italic
		<i>Salmonella</i> spp	1	ground nutmeg	Formatted: Font: Italic
		<i>Bacillus cereus</i> , <i>Bacillus pumilus</i> , <i>Bacillus subtilis</i> , <i>Clostridium perfringens</i> , <i>Salmonella</i> Bareilly, <i>Salmonella enterica</i> <i>Salmonella</i> Give, <i>Salmonella</i> Newport, <i>Salmonella</i> Richmond, <i>Salmonella</i> spp	16	ground pepper	Formatted: Font: Italic
		<i>Salmonella</i> Richmond	1	Indian ginseng powder	Formatted: Font: Not Italic
		<i>Salmonella</i> spp	1	moringa leaves	Formatted: Font: Italic
		<i>Salmonella enterica</i>	3	onion powder	Formatted: Font: Italic
		<i>Salmonella</i> spp			Formatted: Font: Italic
		<i>Salmonella</i> Butantan	1	other herbs	Formatted: Font: Italic
		<i>Bacillus cereus</i> , <i>Clostridium perfringens</i> , <i>Salmonella</i> Amsterdam, <i>Salmonella</i> spp	4	other spices	Formatted: Font: Italic
		<i>Salmonella enterica</i>	1	sesame seeds	Formatted: Font: Italic
		<i>Salmonella</i> Senftenberg, <i>Salmonella</i> Amsterdam	2	turmeric powder	Formatted: Font: Italic
		<i>Salmonella enterica</i>	1	ground peppers	Formatted: Font: Italic
		<i>Salmonella enterica</i> , <i>Salmonella enterica</i> ser. Agona, <i>Salmonella</i> Senftenberg, <i>Salmonella</i> Livingstone, <i>Salmonella</i> spp, <i>Salmonella</i> Virchow	7	turmeric powder	Formatted: Font: Italic
					Formatted: Font: Italic
					Formatted: Font: Italic

Country of origin	Notifications (n)	Pathogens	No of pathogens	Herbs and spices	
Indonesia	4	<i>Bacillus cereus</i>	1	cinnamon powder	Formatted: Font: Italic
		<i>Salmonella</i> Aberdeen, <i>Salmonella</i> Albany, <i>Salmonella</i> Mbandaka	3	ground pepper	Formatted
					Formatted: Font: Italic
Laos	23	<i>Escherichia coli</i> , <i>Salmonella</i> Lexington, <i>Salmonella</i> spp.	4	basil	Formatted
		<i>Salmonella</i> spp.	1	chives	Formatted: Font: Italic
		<i>Salmonella</i> spp.	1	coriander	Formatted
		<i>Escherichia coli</i> , <i>Salmonella</i> Brunei, <i>Salmonella</i> spp., <i>Salmonella</i> Weltevreden	8	mint	Formatted
		<i>Salmonella</i> spp.	1	parsley	Formatted
		<i>Escherichia coli</i> , <i>Salmonella</i> Brunei, <i>Salmonella</i> Heidelberg, <i>Salmonella</i> spp., <i>Salmonella</i> Thompson, <i>Salmonella</i> typhimurium	12	perilla	Formatted
		<i>Escherichia coli</i> , <i>Salmonella</i> Aegona, <i>Salmonella</i> Bareilly, <i>Salmonella</i> Hvitvingfoss, <i>Salmonella</i> Javiana	8	piper lolot	Formatted
		<i>Escherichia coli</i> , <i>Salmonella</i> Hvitvingfoss, <i>Salmonella</i> Meleagridis, <i>Salmonella</i> Rissen, <i>Salmonella</i> spp.	6	praew leaf	Formatted
		<i>Salmonella enterica</i>	1	ground min	Formatted: Font: Italic
Malaysia	3	<i>Salmonella</i> Bareilly, <i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	3	Curry powd	Formatted
		<i>Salmonella</i> spp.	1	parsley and betel leaves	Formatted: Font: Italic
Nepal	1	<i>Salmonella enterica</i> ser. Java	1	betel leave	Formatted
Pakistan	3	<i>Salmonella</i> eEdinburgh	1	ground pepper	Formatted
		<i>Bacillus cereus</i> , <i>Salmonella</i> Hvitvingfoss	2	other spice	Formatted
Singapore	1	<i>Salmonella enterica</i>	1	ground pepper	Formatted: Font: Italic
Sri Lanka	8	<i>Bacillus cereus</i>	1	cinnamon powder	Formatted: Font: Italic
		<i>Salmonella</i> wWaycross, <i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	3	curry leaves	Formatted
		Moulds, <i>Salmonella</i> Aegona, <i>Salmonella</i> Java, <i>Salmonella</i> Pparatyphi B	4	ground pepper	Formatted
		<i>Escherichia coli</i> , <i>Salmonella</i> spp.	2	moringa leaves	Formatted
		<i>Salmonella</i> spp.	1	other herbs	Formatted

Country of origin	Notifications (n)	Pathogens	No of pathogens	Herbs and spices
Thailand	158	<i>Salmonella</i> spp.	1	betel leave
		<i>Salmonella</i> spp. <i>Salmonella</i> Stanley	2	pennywort
		<i>Salmonella</i> spp.	1	phak-phae
		<i>Salmonella</i> Hvittingfoss, <i>Escherichia coli</i> , <i>Salmonella enterica</i> , <i>Salmonella</i> Hadar, <i>Salmonella</i> spp., <i>Salmonella</i> Stanley, <i>Salmonella</i> Virchow, <i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	10	acacia
		<i>Salmonella</i> spp.	1	amaranth greens
		<i>Salmonella</i> spp.	1	Asiatic pennywort
		<i>Salmonella</i> Thompson, <i>Escherichia coli</i> , <i>Salmonella</i> spp., <i>Salmonella</i> Aagona, <i>Salmonella</i> Amsterdam, <i>Salmonella</i> arizonae, <i>Salmonella</i> augustenborg, <i>Salmonella</i> Bareilly, <i>Salmonella</i> Brunei, <i>Salmonella</i> Dublin, <i>Salmonella enterica</i> , <i>Salmonella</i> Goverdhan, <i>Salmonella</i> Java, <i>Salmonella enterica</i> subsp. Salamae, <i>Salmonella</i> group D, <i>Salmonella</i> Hvittingfoss, <i>Salmonella</i> Javiana, <i>Salmonella</i> Lexington, <i>Salmonella</i> Rubislaw, <i>Salmonella</i> Saint Paul, <i>Salmonella</i> spp., <i>Salmonella</i> Stanley, <i>Salmonella</i> Thompson, <i>Salmonella</i> Weltevreden	65	basil
		<i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	2	betel leave
		<i>Salmonella</i> spp.	1	Cha leaves
		<i>Salmonella</i> Augustenborg	1	chives
		<i>Salmonella</i> Aenatum, <i>Salmonella</i> Augustenborg, <i>Salmonella</i> Brunei, <i>Salmonella</i> Corvallis, <i>Salmonella</i> enterica, <i>Salmonella enterica</i> ser. Hvittingfoss, <i>Salmonella enterica</i> ser. Lexington, <i>Salmonella</i> Hvittingfoss, <i>Salmonella</i> Jerusalem, <i>Salmonella</i> Newport, <i>Salmonella</i> Rissen, <i>Salmonella</i> Rubislaw, <i>Salmonella</i> Singapore, <i>Salmonella</i> spp., <i>Salmonella</i> Virchow, <i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	24	coriander
		<i>Salmonella</i> Augustenborg	1	doksadao
		<i>Salmonella</i> Mbandaka	1	edible flower
		<i>Salmonella</i> Weltevreden	1	galingale

Country of origin	Notifications (n)	Pathogens	No of pathogens	Herbs and spices	
		<i>Salmonella</i> Hvittingfoss	2	ground pepper	Formatted
		<i>Salmonella</i> spp			Formatted: Font: Italic
		<i>Salmonella</i> spp	1	guichai leaves	Formatted
		<i>Salmonella</i> spp	2	horopa leaves	Formatted
		<i>Escherichia coli</i> , <i>Salmonella</i> bBraenderup	2	horseradish	Formatted
		<i>Salmonella</i> spp	1	kayang leaves	Formatted
		<i>Salmonella</i> spp	1	lalo leaves	Formatted
		<i>Salmonella</i> group C, <i>Salmonella</i> spp, <i>Salmonella</i> Wandsworth	3	lemon grass	Formatted
		<i>Salmonella</i> Lexington	1	lime leaves	Formatted
		<i>Salmonella</i> spp	1	magaso sadao	Formatted
		<i>Salmonella</i> spp	1	malabar nightshade	Formatted
		<i>Salmonella enterica</i>	1	mimosa	Formatted: Font: Italic
		<i>Escherichia coli</i> , <i>Salmonella</i> Javiana, <i>Salmonella</i> spp, <i>Salmonella</i> Stanley, <i>Salmonella</i> Zanzibar, <i>Salmonella</i> Weltevreden, <i>Salmonella</i> eChester, <i>Salmonella enterica</i> , <i>Salmonella</i> Heidelberg, <i>Salmonella</i> Hvittingfoss, <i>Salmonella</i> Javiana, <i>Salmonella</i> Ppoona, <i>Salmonella</i> spp, <i>Salmonella</i> Stanley, <i>Salmonella</i> Thompson, <i>Salmonella</i> Zanzibar	25	mint	Formatted
		<i>Salmonella</i> Give, <i>Salmonella</i> Kkedougou, <i>Salmonella</i> Typhimurium	3	mint and basil	Formatted
		<i>Salmonella enterica</i>	3	mint leaves	Formatted: Font: Italic
		<i>Salmonella</i> Rissen			Formatted: Font: Italic
		<i>Salmonella</i> Virchow			Formatted: Font: Italic
		<i>Salmonella</i> Bleadon, <i>Salmonella</i> Hvittingfoss, <i>Salmonella</i> Kentucky, <i>Salmonella</i> Mount Pleasant, <i>Salmonella</i> Rfamat-gan, <i>Salmonella</i> Rissen, <i>Salmonella</i> spp	8	morning gl	Formatted: Font: Italic
		<i>Salmonella</i> Hvittingfoss, <i>Salmonella</i> Javiana, <i>Salmonella</i> Stanley, <i>Salmonella</i> Weltevreden, <i>Salmonella</i> Zanzibar	5	other herbs	Formatted
		<i>Salmonella</i> spp	1	pakdum	Formatted
		<i>Salmonella enterica</i>	2	pandang leaves	Formatted: Font: Italic
		<i>Salmonella</i> spp			Formatted: Font: Italic
		<i>Salmonella</i> augustenburg, <i>Salmonella</i> Newport, <i>Salmonella</i> spp	3	parsley	Formatted
		<i>Salmonella</i> Virchow	1	perilla	Formatted
		<i>Salmonella</i> spp	3	praew leav	Formatted

Country of origin	Notifications (n)	Pathogens	No of pathogens	Herbs and spices	
		<i>Salmonella enterica</i>	1	rice paddy herb	Formatted: Font: Italic
		<i>Salmonella</i> Saint Paul, <i>Salmonella</i> spp., <i>Salmonella</i> Stanley	3	Spinach	Formatted
		<i>Salmonella</i> Hvittingfoss	1	wildbetal le...	Formatted: Font: Italic
		<i>Salmonella</i> Augustenborg, <i>Salmonella</i> Paratyphi B (variant Java monophasic variant 4,5,12)	2	Basil and ri paddy herb	Formatted
		<i>Salmonella</i> Augustenborg, <i>Salmonella</i> Lexington	2	basiland lin leaves	Formatted
Vietnam	49	<i>Salmonella</i> spp.	2	pennywort	Formatted: Font: Italic
		<i>Escherichia coli</i> , <i>Salmonella</i> Javiana, <i>Salmonella</i> spp., <i>Salmonella</i> Virchow	8	basil	Formatted
		<i>Salmonella enterica</i>	1	celery	Formatted: Font: Italic
		<i>Salmonella enterica</i> ser. Virchow	1	chilli	Formatted
		<i>Salmonella</i> spp.	1	coriander	Formatted
		<i>Salmonella</i> spp.	1	eryngo	Formatted
		<i>Salmonella</i> spp.	2	green herb	Formatted
		<i>Salmonella enterica</i> , <i>Salmonella</i> Lexington, <i>Salmonella</i> Orion, <i>Salmonella</i> spp., <i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	13	ground pepper	Formatted
		<i>Salmonella</i> Stanley	1	hing choi	Formatted
		<i>Salmonella</i> spp.	1	houttunya leaves	Formatted
		<i>Salmonella</i> spp.	1	houttunya leaves	Formatted
		<i>Salmonella</i> spp.	2	kinh gioi leaves	Formatted
		<i>Salmonella</i> spp.	4	la lot leaves	Formatted
		<i>Salmonella enterica</i> , <i>Salmonella</i> spp.	3	mint	Formatted
		<i>Salmonella</i> spp.	1	mong toi leaves	Formatted
		<i>Salmonella</i> spp.	1	morning gl	Formatted
		<i>Salmonella</i> Abony, <i>Salmonella</i> Dublin, <i>Salmonella</i> Orientalis, <i>Salmonella</i> Virchow, <i>Salmonella</i> Weltevreden, <i>Salmonella</i> spp.	7	other herbs	Formatted
		<i>Salmonella</i> spp.	3	parsley	Formatted
		<i>Salmonella</i> spp.	1	perilla	Formatted
		<i>Salmonella</i> spp.	1	Piper sarmentosum	Formatted

Country of origin	Notifications (n)	Pathogens	No of pathogens	Herbs and spices
		<i>Salmonella</i> spp.	1	rau ram leaves
		<i>Salmonella</i> Javiana	1	rice paddy herb

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Bold

1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407

Abatcha, M. G., Effarizah, M. E., & Rusul, G. (2018). Prevalence, antimicrobial resistance, resistance genes and class 1 integrons of *Salmonella* serovars in leafy vegetables, chicken carcasses and related processing environments in Malaysian fresh food markets. *Food Control*, *91*, 170-180.

Ananchaipattana, C., Hosotani, Y., Kawasaki, S., Pongswat, S., Latiful, B. M., Isobe, S., & Inatsu, Y. (2012). Bacterial contamination of soybean curd (tofu) sold in Thailand. *Food Science and Technology Research*, *19*(6), 843-848.

Bag, P. K., Nandi, S., Bhadra, R. K., Ramamurthy, T., Bhattacharya, S., Nishibuchi, M., Hamabata, T., Yamasaki, S., Takeda, Y., & Nair, G. B. (1999). Clonal diversity among recently emerged strains of *Vibrio parahaemolyticus* O3:K6 associated with pandemic spread. *Journal of clinical microbiology*, *37*(7), 2354-2357.

Banerjee, M., & Sarkar, P. K. (2003). Microbiological quality of some retail spices in India. *Food Research International*, *36*(5), 469-474.

Bangtrakulnonth, A., Pornreongwong, S., Pulsrikarn, C., Sawanpanyalert, P., Hendriksen, R. S., Wong, D. M. L. F., & Aarestrup, F. M. (2004). *Salmonella* serovars from humans and other sources in Thailand, 1993-2002. *Emerging infectious diseases*, *10*(1), 131.

Bernard, H., Faber, M., Wilking, H., Haller, S., Höhle, M., Schielke, A., Ducomble, T., Siffczyk, C., Merbecks, S., & Fricke, G. (2014). Large multistate outbreak of norovirus gastroenteritis associated with frozen strawberries, Germany, 2012. *Eurosurveillance*, *19*(8), 20719.

Bisht, A., Kamble, M. P., Choudhary, P., Chaturvedi, K., Kohli, G., Juneja, V. K., Sehgal, S., & Taneja, N. K. (2021). A surveillance of food borne disease outbreaks in India: 2009-2018. *Food Control*, *121*, 107630.

Borchers, A., Teuber, S. S., Keen, C. L., & Gershwin, M. E. (2010). Food safety. *Clinical reviews in allergy & immunology*, *39*(2), 95-141.

Briet, A., Helsen, N., Delannoy, S., Debuiche, S., Brisabois, A., Midelet, G., & Granier, S. (2018). NDM-1-producing *Vibrio parahaemolyticus* isolated from imported seafood. *J Antimicrob Chemother*, *doi*, 10.

Carstens, C. K., Salazar, J. K., & Darkoh, C. (2019). Multistate outbreaks of foodborne illness in the United States associated with fresh produce from 2010 to 2017. *Frontiers in microbiology*, *10*, 2667.

Chaoniruthisai, P., Punnakitikashem, P., & Rajchamaha, K. (2018). Challenges and difficulties in the implementation of a food safety management system in Thailand: A survey of BRC certified food productions. *Food Control*, *93*, 274-282.

Chen, A., Giese, M., & Chen, D. (2020). Flood impact on Mainland Southeast Asia between 1985 and 2018—The role of tropical cyclones. *Journal of Flood Risk Management*, *13*(2), e12598.

Chen, X., Zhu, Q., Yu, F., Zhang, W., Wang, R., Ye, X., Jin, L., Liu, Y., Li, S., & Chen, Y. (2018). Serology, virulence and molecular characteristics of *Vibrio parahaemolyticus* isolated from seafood in Zhejiang province. *PLoS one*, *13*(10), e0204892.

1408 Cook, N., Williams, L., & D'Agostino, M. (2019). Prevalence of Norovirus in produce sold at retail in the  
1409 United Kingdom. *Food microbiology*, 79, 85–89.

1410 DEFRA. (2020). Food Statistics in your pocket: Global and UK supply. Department for Environment  
1411 Food & Rural Affairs.

1412 Dolan, C., & Humphrey, J. (2000). Governance and trade in fresh vegetables: the impact of UK  
1413 supermarkets on the African horticulture industry. *Journal of development studies*, 37(2), 147–  
1414 176.

1415 Dos Santos, S. F., Cardoso, R. d. C. V., Borges, Í. M. P., e Almeida, A. C., Andrade, E. S., Ferreira, I. O., &  
1416 do Carmo Ramos, L. (2020). Post-harvest losses of fruits and vegetables in supply centers in  
1417 Salvador, Brazil: Analysis of determinants, volumes and reduction strategies. *Waste  
1418 Management*, 101, 161–170.

1419 ECDC/EFSA. (2021). Multi-country outbreak of Salmonella Enteritidis sequence type (ST)11 infections  
1420 linked to poultry products in the EU/EEA and the United Kingdom – 25 February 2021.  
1421 Stockholm: ECDC/EFSA; 2021. European Centre for Disease Prevention and Control (ECDC) and  
1422 European Food Safety Authority (EFSA).

1423 EFSA/ECDC. (2019). Scientific report on the European Union one health 2018. Zoonoses report.  
1424 European Centre for Disease Prevention and Control (ECDC) and European Food Safety  
1425 Authority (EFSA). *EFSA J*, 17, 5926.

1426 EFSA/ECDC. (2021). The European Union One Health 2019 Zoonoses Report. European Food Safety  
1427 Authority (EFSA) and European Centre for Disease Prevention and Control (ECDC). *EFSA  
1428 Journal* 19, 6406.

1429 EFSA/ECDC. (2018). Multi-country outbreak of Salmonella Agona infections possibly linked to ready-  
1430 to-eat food. European Food Safety Authority (EFSA) and European Centre for Disease  
1431 Prevention Control (ECDC). In: Wiley Online Library.

1432 Emberland, K., Ethelberg, S., Kuusi, M., Vold, L., Jensvoll, L., Lindstedt, B., Nygård, K., Kjelsø, C.,  
1433 Torpdahl, M., & Sørensen, G. (2007). Outbreak of Salmonella Weltevreden infections in  
1434 Norway, Denmark and Finland associated with alfalfa sprouts, July–October 2007. *Weekly  
1435 releases (1997–2007)*, 12(48), 3321.

1436 European Commission. (2014). Commission implementing decision of 13 February 2014 suspending  
1437 temporarily imports from Bangladesh of foodstuffs containing or consisting of betel leaves  
1438 ('Piper betle'). Available from: [https://eur-lex.europa.eu/legal-  
1439 content/EN/TXT/PDF/?uri=CELEX:32014D0088&from=DA](https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32014D0088&from=DA) Accessed 25.02.2021.

1440 European Commission. (2015). Commission implementing regulation (EU) 2015/943 of 18 June 2015.  
1441 *Official Journal of the European Union*(154), 8–9.

1442 European Commission (2016). RASFF — The Rapid Alert System for Food and Feed — 2015 annual  
1443 report. Available from: [https://op.europa.eu/en/publication-detail/-/publication/98e04cfc-  
1444 aa54-11e6-aab7-01aa75ed71a1/language-en/format-PDF/source-174743070](https://op.europa.eu/en/publication-detail/-/publication/98e04cfc-aa54-11e6-aab7-01aa75ed71a1/language-en/format-PDF/source-174743070) Accessed  
1445 19.02.2021.

1446 European Commission. (2020a). Commission implementing regulation (EU) 2020/625 of 6 May 2020.  
1447 *Official Journal of the European Union*(63), 13.

1448 European Commission. (2020b). Monitoring EU Agri-Food Trade: Developments in 2019. Available  
1449 from: [https://ec.europa.eu/info/news/2019-record-year-eu-agri-food-trade-2020-mar-  
1450 27\\_en#moreinfo](https://ec.europa.eu/info/news/2019-record-year-eu-agri-food-trade-2020-mar-27_en#moreinfo) Accessed 12.02.2021.

1451 European Food Safety Authority. (2021). The European Union One Health 2019 Zoonoses Report. *EFSA  
1452 Journal*, 19(2).

1453 Fakruddin, M., Sultana, R., Hossain, M. N., Rahaman, M. M., Islam, M. K., & Ahmed, M. M. (2017).  
1454 Occurrence of ingestion of Salmonella spp. in Betel leaf (Piper betle L.). *International Journal  
1455 of Food Contamination*, 4(1), 1–10.

1456 Ferrari, R. G., Rosario, D. K., Cunha Neto, A., Mano, S. B., Figueiredo, E. E., & Conte Junior, C. A. (2019).  
1457 Worldwide epidemiology of Salmonella serovars in animal-based foods: a meta-analysis.  
1458 *Applied and environmental microbiology*, 85(14).

1459 Food and Agricultural Organisation. (2018). Addressing the food safety challenges of the Asia-Pacific  
1460 region. 34th Session of the FAO Regional Conference for Asia and the Pacific. Available from:  
1461 <http://www.fao.org/3/MV819en/mv819en.pdf> Accessed 24.02.2021.

1462 Food and Agriculture Organization. (2019). Foodborne illnesses in Asia-Pacific 'needlessly' sicken 275  
1463 million annually and threaten trade. Retrieved from  
1464 [http://www.fao.org/asiapacific/news/detail-  
1465 events/en/c/1197008/#:~:text=In%20the%20Asia%2DPacific%20region,of%20Zero%20Hunger  
1466 %20by%202030.](http://www.fao.org/asiapacific/news/detail-events/en/c/1197008/#:~:text=In%20the%20Asia%2DPacific%20region,of%20Zero%20Hunger%20by%202030.)

1467 Fu, H., Yu, P., Liang, W., Kan, B., Peng, X., & Chen, L. (2020). Virulence, Resistance, and Genomic  
1468 Fingerprint Traits of *Vibrio cholerae* Isolated from 12 Species of Aquatic Products in Shanghai,  
1469 China. *Microbial Drug Resistance*, 26(12), 1526-1539.

1470 Ghosh, D. (2014). Food safety regulations in Australia and New Zealand food standards. *Journal of the  
1471 Science of Food and Agriculture*, 94(10), 1970-1973.

1472 Jaffee, S., Henson, S., Unnevehr, L., Grace, D., & Cassou, E. (2018). *The safe food imperative:  
1473 Accelerating progress in low and middle income countries*. World Bank Publications.

1474 Kaakoush, N. O., Castaño-Rodríguez, N., Mitchell, H. M., & Man, S. M. (2015). Global epidemiology of  
1475 *Campylobacter* infection. *Clinical microbiology reviews*, 28(3), 687-720.

1476 Kantama, L., & Jayanetra, P. (1996). Salmonella enteritidis outbreak in Thailand: study by random  
1477 amplified polymorphic DNA (RAPD) analysis. *The Southeast Asian journal of tropical medicine  
1478 and public health*, 27(1), 119-125.

1479 Keisam, S., Tuikhar, N., Ahmed, G., & Jeyaram, K. (2019). Toxigenic and pathogenic potential of enteric  
1480 bacterial pathogens prevalent in the traditional fermented foods marketed in the Northeast  
1481 region of India. *International journal of food microbiology*, 296, 21-30.

1482 Kim, K., Kim, S., & Park, C. Y. (2020). Food Security in Asia and the Pacific amid the COVID-19  
1483 Pandemic. 1-15.

1484 Kowalska, A., & Manning, L. (2020). Using the rapid alert system for food and feed: potential benefits  
1485 and problems on data interpretation. *Critical reviews in food science and nutrition*, 1-14.

1486 Kumar, R., & Lalitha, K. V. (2013). Prevalence and molecular characterization of *Vibrio cholerae* O1,  
1487 non-O1 and non-O139 in tropical seafood in Cochin, India. *Foodborne pathogens and disease*,  
1488 10(3), 278-283.

1489 Kumari, S., & Sarkar, P. K. (2014). Prevalence and characterization of *Bacillus cereus* group from  
1490 various marketed dairy products in India. *Dairy science & technology*, 94(5), 483-497.

1491 Lee, Y., Choi, Y., Lee, S., Lee, H., Kim, S., Ha, J., Lee, J., Oh, H., Kim, Y., & Yoon, Y. (2019). Occurrence of  
1492 pathogenic *Vibrio parahaemolyticus* in seafood distribution channels and their antibiotic  
1493 resistance profiles in S. Korea. *Letters in applied microbiology*, 68(2), 128-133.

1494 Lemoine, T., Germanetto, P., & Giraud, P. (1999). Toxi-infection alimentaire à *V. parahaemolyticus*.  
1495 *Bull. Epidémiol. Hebdo*, 10, 37-38.

1496 Machado-Moreira, B., Richards, K., Brennan, F., Abram, F., & Burgess, C. M. (2019). Microbial  
1497 contamination of fresh produce: what, where, and how? *Comprehensive reviews in food  
1498 science and food safety*, 18(6), 1727-1750.

1499 Makendi, C., Page, A. J., Wren, B. W., Le Thi Phuong, T., Clare, S., Hale, C., Goulding, D., Klemm, E. J.,  
1500 Pickard, D., & Okoro, C. (2016). A phylogenetic and phenotypic analysis of *Salmonella enterica*  
1501 serovar Weltevreden, an emerging agent of diarrheal disease in tropical regions. *PLoS  
1502 neglected tropical diseases*, 10(2), e0004446.

1503 Mangla, S. K., Bhattacharya, A., Yadav, A. K., Sharma, Y. K., Ishizaka, A., Luthra, S., & Chakraborty, R.  
1504 (2021). A framework to assess the challenges to food safety initiatives in an emerging  
1505 economy. *Journal of Cleaner Production*, 284, 124709.

1506 Martinez-Urtaza, J., Simental, L., Velasco, D., DePaola, A., Ishibashi, M., Nakaguchi, Y., Nishibuchi, M.,  
1507 Carrera-Flores, D., Rey-Alvarez, C., & Pousa, A. (2005). Pandemic *vibrio parahaemolyticus* O3:  
1508 K6, Europe. *Emerging infectious diseases*, 11(8), 1319.

1509 McLauchlin, J., Aird, H., Andrews, N., Chattaway, M., De Pinna, E., Elviss, N., Jørgensen, F., Larkin, L., &  
1510 Willis, C. (2019). Public health risks associated with Salmonella contamination of imported  
1511 edible betel leaves: analysis of results from England, 2011–2017. *International journal of food*  
1512 *microbiology*, *298*, 1–10.

1513 McLauchlin, J., Aird, H., Charlett, A., Chattaway, M., Elviss, N., Hartman, H., Jenkins, C., Jørgensen, F.,  
1514 Larkin, L., & Sadler-Reeves, L. (2018). Imported edible leaves collected at retail sale in England  
1515 during 2017 with an emphasis on betel and curry leaves: microbiological quality with respect  
1516 to Salmonella, Shiga-toxin-producing *E. coli* (STEC) and levels of *Escherichia coli*. *Journal of*  
1517 *applied microbiology*, *125*(4), 1175–1185.

1518 Mensah, L. D., & Julien, D. (2011). Implementation of food safety management systems in the UK.  
1519 *Food Control*, *22*(8), 1216–1225.

1520 Nakaguchi, Y. (2013). Contamination with *Vibrio parahaemolyticus* and its virulent strains in seafood  
1521 marketed in Thailand, Vietnam, Malaysia, and Indonesia from 2008 to 2011. *Tropical Medicine*  
1522 *and Health*, 2011–2006.

1523 Nasheri, N., Vester, A., & Petronella, N. (2019). Foodborne viral outbreaks associated with frozen  
1524 produce. *Epidemiology & Infection*, *147*.

1525 Nath, T. K., & Inoue, M. (2009). Sustainability attributes of a small-scale betel leaf agroforestry system:  
1526 a case study in north-eastern hill forests of Bangladesh. *Small-scale Forestry*, *8*(3), 289–304.

1527 Nguyen, T. K., Bui, H. T., Truong, T. A., Lam, D. N., Ikeuchi, S., Ly, L. K. T., Hara-Kudo, Y., Taniguchi, T., &  
1528 Hayashidani, H. (2021). Retail fresh vegetables as a potential source of Salmonella infection in  
1529 the Mekong Delta, Vietnam. *International journal of food microbiology*, *341*, 109049.

1530 Office of National Statistics. (2011). UK Census 2011. Available online:  
1531 <https://www.ons.gov.uk/census/2011census>

1532 Ortega, D. L. a. T., D. L. (2017). Demand for food safety in emerging and developing countries: A  
1533 research agenda for Asia and Sub-Saharan Africa. *Journal of Agribusiness in Developing and*  
1534 *Emerging Economies*(7), 21–34.

1535 Park, S., Szonyi, B., Gautam, R., Nightingale, K., Anciso, J., & Ivanek, R. (2012). Risk factors for microbial  
1536 contamination in fruits and vegetables at the preharvest level: a systematic review. *Journal of*  
1537 *food protection*, *75*(11), 2055–2081.

1538 Pham, H. V., & Dinh, T. L. (2020). The Vietnam's food control system: Achievements and remaining  
1539 issues. *Food Control*, *108*, 106862.

1540 Reddy, A. A., Ricart, S., & Cadman, T. (2020). Driving factors of food safety standards in India: learning  
1541 from street food vendors' behaviour and attitude. *Food Security*, 1–17.

1542 Rezaei, M., & Liu, B. (2017). Food loss and waste in the food supply chain. *International Nut and Dried*  
1543 *Fruit Council: Reus, Spain*, 26–27.

1544 Rivas, L., Strydom, H., Paine, S., Wang, J., & Wright, J. (2021). Yersiniosis in New Zealand. *Pathogens*,  
1545 *10*(2), 191.

1546 Singh, B. R., Singh, M., Singh, P., Babu, N., Chandra, M., & Agarwal, R. K. (2006). Prevalence of  
1547 multidrug-resistant Salmonella on ready-to-eat betel leaves (Paan) and in water used for  
1548 soaking betel leaves in North Indian cities. *Journal of food protection*, *69*(2), 288–292.

1549 Sodagari, H. R., Wang, P., Robertson, I., Habib, I., & Sahibzada, S. (2020). Non-Typhoidal Salmonella at  
1550 the Human-Food-of-Animal-Origin Interface in Australia. *Animals*, *10*(7), 1192.

1551 Somorin, Y. M., Odeyemi, O. A., & Ateba, C. N. (2021). Salmonella is the most common foodborne  
1552 pathogen in African food exports to the European Union: Analysis of the Rapid Alert System  
1553 for Food and Feed (1999–2019). *Food Control*, *123*, 107849.

1554 Sugiri, Y. D., Goelz, G., Meeyam, T., Baumann, M. P., Kleer, J., Chaisowwong, W., & Alter, T. (2014).  
1555 Prevalence and antimicrobial susceptibility of *Listeria monocytogenes* on chicken carcasses in  
1556 Bandung, Indonesia. *Journal of food protection*, *77*(8), 1407–1410.

1557 Tähkääpää, S., Maijala, R., Korkeala, H., & Nevas, M. (2015). Patterns of food frauds and adulterations  
1558 reported in the EU rapid alert system for food and feed and in Finland. *Food Control*, *47*, 175–  
1559 184.

1560 Ullah, A. S., Tani, M., Tsuchiya, J., Rahman, A. M., & Rahman, Z. M. (2020). Impact of Betel Leaf  
1561 Cultivation on the Protected Forest Area of Teknaf Peninsula, Bangladesh. *Small-scale*  
1562 *Forestry*, *19*(3), 335-355.

1563 Vilain, S., Luo, Y., Hildreth, M. B., & Brözel, V. S. (2006). Analysis of the life cycle of the soil saprophyte  
1564 *Bacillus cereus* in liquid soil extract and in soil. *Applied and environmental microbiology*, *72*(7),  
1565 4970-4977.

1566 Vital, P. G., Dimasuay, K. G. B., Widmer, K. W., & Rivera, W. L. (2014). Microbiological quality of fresh  
1567 produce from open air markets and supermarkets in the Philippines. *The Scientific World*  
1568 *Journal*, 2014.

1569 Wadamori, Y., Gooneratne, R., & Hussain, M. A. (2017). Outbreaks and factors influencing  
1570 microbiological contamination of fresh produce. *Journal of the Science of Food and*  
1571 *Agriculture*, *97*(5), 1396-1403.

1572 Wang, S., Weller, D., Falardeau, J., Strawn, L. K., Mardones, F. O., Adell, A. D., & Switt, A. I. M. (2016).  
1573 Food safety trends: from globalization of whole genome sequencing to application of new  
1574 tools to prevent foodborne diseases. *Trends in Food Science & Technology*, *57*, 188-198.

1575 Wong, H.-C., Chen, M.-C., Liu, S.-H., & Liu, D.-P. (1999). Incidence of highly genetically diversified *Vibrio*  
1576 *parahaemolyticus* in seafood imported from Asian countries. *International journal of food*  
1577 *microbiology*, *52*(3), 181-188.

1578 World Health Organization. (2015). *WHO estimates of the global burden of foodborne diseases: foodborne disease burden epidemiology reference group 2007-2015*. World Health Organization.

1580 World Health Organization. (2019). *Shiga Toxin producing Escherichia Coli (STEC) and Food: Attribution Characterization and Monitoring (Vol. 19)*. World Health Organization.

1581 World Health Organization. (2020a). Food Safety. Retrieved from <https://www.who.int/news-room/fact-sheets/detail/food-safety>.

1582 World Health Organization. (2020b). Framework for action on food safety in the WHO South-East Asia Region. New Delhi: World Health Organization, Regional Office for South-East Asia.

1583 World Health Organization. (2020c). Salmonella – non-typhoidal. Available from: [https://www.who.int/news-room/fact-sheets/detail/salmonella-\(non-typhoidal\)](https://www.who.int/news-room/fact-sheets/detail/salmonella-(non-typhoidal)) Accessed 25.02.2021.

1584 Worldometer. (2021). Current World Population. Retrieved from <https://www.worldometers.info/world-population/#ref-1>.

1585 Xu, X., Cheng, J., Wu, Q., Zhang, J., & Xie, T. (2016). Prevalence, characterization, and antibiotic susceptibility of *Vibrio parahaemolyticus* isolated from retail aquatic products in North China. *BMC microbiology*, *16*(1), 1-9.

1586 Yu, P., Zaleski, A., Li, Q., He, Y., Mapili, K., Pruden, A., Alvarez, P. J., & Stadler, L. B. (2018). Elevated levels of pathogenic indicator bacteria and antibiotic resistance genes after Hurricane Harvey's flooding in Houston. *Environmental science & technology letters*, *5*(8), 481-486.

**Declaration of competing interest**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**CRedit authorship contribution statement**

**Ayokunle C. Dada:** Conceptualization, Methodology, Formal analysis, Writing - original draft, Review and Editing, Writing - review & editing. **Yinka M. Somorin:** Writing - original draft, Review and Editing, Writing - review & editing. **Collins N. Ateba:** Writing - original draft, Review and Editing, Writing - review & editing. **Helen Onyeaka:** Review and Editing, Writing - review & editing. **Amarachukwu Anyogu:** Review and Editing, Writing - review & editing. **Nor Azman Kasan:** Writing - original draft, Review and Editing, Writing - review & editing. **Olumide A. Odeyemi:** Conceptualization, Methodology, Formal analysis, Writing - original draft, Review and Editing, Writing - review & editing.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65