



## **UWL REPOSITORY**

**repository.uwl.ac.uk**

Global disease burden linked to diet high in red meat and colorectal cancer from 1990 to 2019 and its prediction up to 2030

Yang, X, Wu, D, Liu, Y, He, Z, Manyande, Anne ORCID logo ORCID: <https://orcid.org/0000-0002-8257-0722>, Fu, H and Xiang, H (2024) Global disease burden linked to diet high in red meat and colorectal cancer from 1990 to 2019 and its prediction up to 2030. *Frontiers in Nutrition*, 11. ISSN 2296-861X

<http://dx.doi.org/10.3389/fnut.2024.1366553>

This is the Accepted Version of the final output.

**UWL repository link:** <https://repository.uwl.ac.uk/id/eprint/11245/>

**Alternative formats:** If you require this document in an alternative format, please contact: [open.research@uwl.ac.uk](mailto:open.research@uwl.ac.uk)

**Copyright:** Creative Commons: Attribution 4.0

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy:** If you believe that this document breaches copyright, please contact us at [open.research@uwl.ac.uk](mailto:open.research@uwl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

**Rights Retention Statement:**

## **Global disease burden linked to diet high in red meat and colorectal cancer from 1990 to 2019 and its prediction up to 2030**

Xuesong Yang<sup>1,#</sup>, Duozi Wu<sup>2,#</sup>, Yanbo Liu<sup>1</sup>, Zhigang He<sup>1</sup>, Anne Manyande<sup>3</sup>, Hongjun Fu<sup>4,\*</sup>,  
Hongbing Xiang<sup>1,5,\*</sup>

<sup>1</sup> Department of Anesthesiology and Pain Medicine, Hubei Key Laboratory of Geriatric Anesthesia and Perioperative Brain Health, Wuhan Clinical Research Center for Geriatric Anesthesia, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China

<sup>2</sup> Department of Anesthesiology, Hainan general Hospital, Haikou, Hainan, China

<sup>3</sup> School of Human and Social Sciences, University of West London, London, United Kingdom

<sup>4</sup> Department of Orthopedics, Xiangyang Hospital of Traditional Chinese Medicine, Xiangyang, Hubei, China

<sup>5</sup> Key Laboratory of Anesthesiology and Resuscitation (Huazhong University of Science and Technology), Ministry of Education, China

# These authors contributed equally

\*Address correspondence to: Hongbing Xiang, E-mail: [xhbtj2004@163.com](mailto:xhbtj2004@163.com); Hongjun Fu, E-mail:

[15971036634@163.com](mailto:15971036634@163.com)

## **Abstract**

Numerous studies have already identified an association between excessive consumption of red meat and colorectal cancer (CRC). However, there has been a lack of detailed understanding regarding the disease burden linked to diet high in red meat and CRC. Our objective was to evaluate global, regional, and national mortality rates and disability-adjusted Life years (DALYs) related to this diet. We also considered factors such as sex, age, the socio-demographic index (SDI), and evaluated the cross-national inequalities. Furthermore, we utilized DALYs data from 204 countries and regions to gauge cross-country inequalities of CRC by calculating the slope index of inequality and concentration index as standard indicators of absolute and relative inequalities. Our data was derived from the Global Burden of Disease (GBD) Study 2019. The results show that globally, the ASMR and ASDR related to CRC due to diet high in red meat have decreased, with EAPCs of -0.32% (95% CI -0.37 to -0.28) and -0.18% (95% CI -0.25 to -0.11). Notably, the burden was higher among males and the elderly. The slope index of inequality rose from 22.0 (95% CI 18.1 to 25.9) in 1990 to 32.9 (95% CI 28.3 to 37.5) in 2019 and the concentration index fell from 59.5 (95% CI 46.4 to 72.6) in 1990 to 48.9 (95% CI 34.6 to 63.1) in 2019. We aim to offer evidence-based guidance for developing effective strategies that can mitigate the elevated CRC burden in certain countries.

**Keywords:** colorectal cancer, red meat, Global Burden of Disease, mortality, disability-adjusted life years, health inequality, epidemiology

## **1. Introduction**

Colorectal cancer (CRC), also known as colorectal adenocarcinoma, primarily arises from the colon's glands and epithelial cells[1]. Worldwide, CRC is ranked third in cancer prevalence among males and second among females[2]. It is worth noting that more than half of the cases occur in more developed and industrialized countries[2, 3]. In these countries, CRC is the second most common tumor, with a lifetime incidence of 5%[4-6]. In regions with varying levels of economic development, there are significant differences in the 5-year survival rate of CRC. For instance, the overall 5-year survival rate in the United States is over 60%, whereas in developing countries, it is less than 40%[4].

The International Agency for Research on Cancer (IARC) defines red meat as unprocessed mammalian muscle meat[7, 8]. Dietary red meat, including beef, veal, pork, lamb, and mutton, provides us with several important essential nutrients[9, 10]. These essential nutrients comprise protein, essential amino acids, vitamins (including vitamin B12), minerals (including heme iron and zinc), and other micronutrients[11]. With the recent economic growth, the global demand for red meat has surged, in both developed and developing countries[12]. However, abundant evidence indicates that the diet high in red meat is associated with a range of health issues[13-15], including two major chronic diseases: cardiovascular disease and CRC[16-18]. A 1975 survey revealed a robust correlation ( $r=0.9$ ) between per capita meat consumption and the incidence of CRC among women from 23 different countries[19]. Other research also reported that an increase in red meat consumption by 100 grams per day is associated with an 11%-51% greater risk of various cancer incidence and appears to be unrelated to any health benefits[20]. IARC has classified red meat consumption as a probable human carcinogen based on evidence related to CRC, pancreatic cancer, and prostate cancer (Group 2A)[8]. Recent guidelines recommend that the public should moderate their daily intake of red meat[21].

Previous Global Burden of Disease (GBD) studies predominantly emphasized the disease burden from high dietary red meat, rather than specifically addressing the CRC burden attributable to such consumption[22, 23]. Currently, no detailed report exists based on the GBD dataset focusing on the CRC burden from high global red meat consumption. Hence, we utilized data from GBD 2019 to determine the temporal trends in incidence and disability-adjusted life years (DALYs) at the global, regional, and national levels, stratified by sex, age, and SDI. We systematically summarized the global burden and health development status, including the unequal distribution of disease burden among countries, in order to provide evidence for policymakers. Given the public accessibility of the data, ethical approval and informed consent were not required for our study.

## **2. Materials and method**

### **2.1 Data source**

The data from the Global Burden of Disease (GBD) for the year 2019 were used to estimate the incidence and burden of CRC. Data on CRC were sourced from various outlets, including hospital records, emergency department records, insurance claims, surveys, and the Global Vital Registration system.

The GBD data processing and model construction, which have been previously discussed in articles, were employed for this research[24, 25].

We extracted data from the Global Health Data Exchange website (<http://ghdx.healthdata.org/gbd-results-tool>) for the years 1990 to 2019, stratified by sex, age, and region. This data included CRC-

related deaths, DALYs (Disability-Adjusted Life Years), age-standardized mortality rates (ASMR), and age-standardized DALY rates (ASDR).

Additionally, the relationship between SDI and the burden of disease were investigated. SDI is a composite index that considers a country's per capita income, average years of education, and fertility rate. It categorized 204 countries and regions into five groups: low SDI (<0.45), medium SDI ( $\geq 0.45$  and <0.61), medium SDI ( $\geq 0.61$  and <0.69), medium SDI ( $\geq 0.69$  and <0.80), and high SDI ( $\geq 0.80$ )[26, 27].

High red meat consumption in the diet was defined as a daily intake of red meat (beef, pork, lamb, and goat, excluding poultry, fish, eggs, and all processed meats) exceeding 23g (ranging from 18g to 27g) as an optional level[28]. Detailed information about inclusion and exclusion criteria can be found in the preceding sections[25]. This research aimed to assess the incidence and burden of CRC while exploring its relationship with social demographic indices and diet high in red meat.

## **2.2. Statistical analyses**

We analyzed the burden of CRC at the global national level. To account for demographic differences, we used age-standardized incidence rates (ASIR), mortality rates, and Disability-Adjusted Life Years (DALY) rates to better reflect actual incidence and mortality rates. All rates are expressed as per 100,000 people.

Additionally, estimated annual percentage changes (EAPC) were calculated using a linear regression equation:  $y = \alpha + \beta x + \varepsilon$  (where  $y$  represents  $\ln(\text{ASMR or ASDR})$ , and  $x$  represents the calendar year) to assess trends in ASMR and ASDR. The exact calculation is  $\text{EAPC} = 100 (\exp(\beta) - 1)$ [29]. EAPC and its 95% confidence interval (CI) greater than 0 indicate an increasing trend in ASMR or ASDR over the years, while values lower than 0 indicate a decreasing trend. If EAPC is close to 0, it suggests stability[30].

Furthermore, to explore the factors affecting the EAPC of the CRC burden related to a diet high in red meat, we also assessed the correlation between the age-standardized indicator of diet high in red meat in 1990 and SDI in 2019 with EAPC at the national level using the Spearman rank test.

We used the concentration index (CI) and the slope index to quantify the health inequalities. The slope index of inequality and concentration index, are two standard indicators of absolute inequality and relative inequality, respectively[31]. The slope index of inequality is calculated by regressing the national DALYs ratio for all age groups on a relative positional scale associated with SDI, defined as the midpoint of the population cumulative range ranked by SDI[32]. Heteroscedasticity is explained by a weighted regression model. The concentration index is calculated by numerically integrating the area under the Lorenz concentration curve, which is fitted using the cumulative scores of DALYs and the cumulative relative distribution of the population based on SDI[33]. All statistical analyses were performed with R software (version 4.2.3).

## **3. Result**


### **3.1. Global spatial and temporal colorectal cancer burden attributable to diet high in red meat**

Globally, in 1990 and 2019, approximately 0.026 million and 0.052 million deaths from CRC were linked to a diet high in red meat. The male-to-female ratio for these deaths was around 1.3 and 1.5, respectively, for CRC. During this period, there was a rapid increase in deaths and DALYs associated with diet high in red meat. However, the ASMR and ASDR showed little change, with EAPC in ASMR for females being less than 0, and greater than 0 for males (Table 1, 2; Figure 1).

*Table 1. Deaths of colon and rectum cancer attributable to diet high in red meat in 1990 and 2019 for both sexes and all locations, with EAPC from 1990 and 2019.*

location	Deaths cases in 1990	ASMR per 100 000 in 1990	Deaths cases in 2019	ASMR per 100 000 in 2019	EAPC (1990– 2019)
Global	26087 (6690 to 50231)	0.71 (0.18 to 1.38)	52811 (13598 to 100688)	0.66 (0.17 to 1.26)	-0.32% (-0.37 to -0.28)
Both					
Female	12951 (3302 to 24856)	0.64 (0.16 to 1.23)	22760 (5663 to 44004)	0.52 (0.13 to 1.01)	-0.83% (-0.88 to -0.78)
Male	13136 (3427 to 25156)	0.81 (0.21 to 1.56)	30051 (7627 to 57328)	0.83 (0.21 to 1.59)	0.07% (0.01 to 0.12)
Region	14063 (3944 to 25419)	1.34 (0.38 to 2.42)	18849 (4893 to 35349)	0.97 (0.26 to 1.79)	-1.29% (-1.37 to -1.21)
High SDI					
High-middle SDI	8864 (2276 to 17225)	0.87 (0.22 to 1.7)	18793 (5033 to 35132)	0.93 (0.25 to 1.74)	0.09% (0.01 to 0.17)
Middle SDI	2278 (297 to 5662)	0.23 (0.03 to 0.59)	11723 (2560 to 23587)	0.49 (0.1 to 0.99)	2.95% (2.77 to 3.12)
Low-middle SDI	649 (122 to 1497)	0.12 (0.02 to 0.27)	2826 (617 to 5904)	0.22 (0.05 to 0.45)	2.38% (2.28 to 2.48)
Low SDI	221 (32 to 565)	0.1 (0.01 to 0.26)	598 (98 to 1464)	0.12 (0.02 to 0.3)	0.81% (0.72 to 0.9)
Central Sub-Saharan Africa	27 (5 to 67)	0.13 (0.02 to 0.32)	61 (10 to 150)	0.12 (0.02 to 0.3)	-0.18% (-0.51 to 0.15)
East Asia	2851 (408 to 6970)	0.34 (0.05 to 0.84)	16498 (4056 to 31569)	0.82 (0.2 to 1.58)	3.6% (3.35 to 3.86)
Eastern Europe	2995 (772 to 5768)	1.07 (0.28 to 2.07)	2708 (468 to 5947)	0.79 (0.14 to 1.72)	-1.91% (-2.21 to -1.62)
Eastern Sub-Saharan Africa	88 (10 to 231)	0.12 (0.01 to 0.32)	241 (27 to 626)	0.16 (0.02 to 0.4)	0.9% (0.79 to 1.02)
Andean Latin America	36 (4 to 92)	0.18 (0.02 to 0.47)	154 (17 to 392)	0.28 (0.03 to 0.71)	1.81% (1.68 to 1.95)
High-income Asia Pacific	910 (87 to 2421)	0.47 (0.05 to 1.25)	2196 (275 to 5546)	0.47 (0.06 to 1.14)	-0.25% (-0.35 to -0.15)
High-income North America	4798	1.36	6386	1.02	-1.21%

location	Deaths cases in 1990	ASMR per 100 000 in 1990	Deaths cases in 2019	ASMR per 100 000 in 2019	EAPC (1990– 2019)
	(1332 to 8719)	(0.38 to 2.44)	(1712 to 11567)	(0.28 to 1.82)	(-1.33 to -1.09)
Caribbean	83 (8 to 220)	0.33 (0.03 to 0.87)	209 (20 to 560)	0.4 (0.04 to 1.08)	0.9% (0.79 to 1.01)
Australasia	630 (276 to 959)	2.72 (1.19 to 4.15)	888 (367 to 1387)	1.75 (0.74 to 2.71)	-1.8% (-1.97 to -1.63)
Central Europe	1538 (322 to 3190)	1.06 (0.22 to 2.23)	2862 (699 to 5678)	1.33 (0.33 to 2.61)	1.04% (0.92 to 1.17)
Central Latin America	189 (24 to 461)	0.24 (0.03 to 0.58)	807 (130 to 1946)	0.34 (0.05 to 0.84)	1.38% (1.33 to 1.42)
Central Asia	271 (70 to 524)	0.57 (0.14 to 1.11)	389 (98 to 756)	0.56 (0.14 to 1.11)	0.39% (-0.13 to 0.91)
North Africa and Middle East	301 (29 to 820)	0.18 (0.02 to 0.5)	882 (86 to 2352)	0.22 (0.02 to 0.57)	0.76% (0.51 to 1.02)
Oceania	6 (1 to 16)	0.22 (0.03 to 0.59)	15 (2 to 41)	0.24 (0.03 to 0.64)	0.11% (-0.02 to 0.24)
South Asia	267 (70 to 583)	0.05 (0.01 to 0.12)	947 (228 to 2161)	0.07 (0.02 to 0.16)	0.84% (0.71 to 0.97)
Southeast Asia	406 (51 to 1036)	0.17 (0.02 to 0.42)	1822 (220 to 4550)	0.31 (0.04 to 0.77)	2.15% (2.08 to 2.21)
Southern Latin America	896 (376 to 1387)	2 (0.82 to 3.09)	1768 (705 to 2777)	2.1 (0.84 to 3.3)	0.15% (0.06 to 0.23)
Southern Sub-Saharan Africa	83 (11 to 211)	0.32 (0.04 to 0.84)	209 (30 to 498)	0.4 (0.06 to 0.95)	1.01% (0.84 to 1.18)
Tropical Latin America	388 (78 to 834)	0.45 (0.09 to 0.98)	2403 (890 to 3925)	1 (0.36 to 1.64)	3.02% (2.41 to 3.64)
Western Europe	9232 (2731 to 16476)	1.59 (0.48 to 2.82)	11104 (2953 to 20565)	1.16 (0.32 to 2.11)	-1.33% (-1.44 to -1.22)
Western Sub-Saharan Africa	92 (10 to 244)	0.11 (0.01 to 0.3)	262 (29 to 684)	0.16 (0.02 to 0.41)	1.35% (1.23 to 1.48)



**Figure 1.** The burden of colorectal cancer deaths (A) and DALYs (B) due to diet high in red meat from 1990 to 2019 by sex and SDI region. The bar is the number of colorectal cancer deaths and DALYs attributable to diet high in red meat. The line with 95% UI represents ASMR and ASDR attributable to diet high in red meat. ASMR, age-standardized mortality rate; DALYs, disability-adjusted life-years; ASDR, age-standardized DALY rate; UI, uncertainty interval; SDI, sociodemographic index.

**Table 2. DALYs of colon and rectum cancer attributable to diet high in red meat in 1990 and 2019 for both sexes and all locations, with EAPC from 1990 and 2019.**

location	DALYs in 1990	ASDR per 100 000 in 1990	DALYs in 2019	ASDR per 100 000 in 2019	EAPC (1990– 2019)
Global	627834	15.63	1234678	14.95	-0.18%
Both	(165197 to 1192278)	(4.08 to 29.8)	(332704 to 2306844)	(4.02 to 27.99)	(-0.25 to -0.11)
Female	296273 (76863 to 562565)	13.86 (3.6 to 26.4)	496397 (132360 to 944785)	11.43 (3.05 to 21.75)	-0.77% (-0.84 to -0.71)
Male	331561	17.73	738281	18.83	0.24%



location	DALYs in 1990	ASDR per 100 000 in 1990	DALYs in 2019	ASDR per 100 000 in 2019	EAPC (1990– 2019)
	(88528 to 627390)	(4.67 to 33.68)	(198240 to 1373853)	(5.03 to 35.12)	(0.16 to 0.32)
Region	314924	30.93	389921	22.76	-1.2%
High SDI	(92479 to 557564)	(9.16 to 54.6)	(108477 to 703495)	(6.57 to 40.43)	(-1.27 to -1.12)
High-middle SDI	221808	20.29	440051	21.86	0.09%
	(58212 to 422210)	(5.29 to 38.8)	(124026 to 802903)	(6.2 to 39.94)	(-0.01 to 0.19)
Middle SDI	65863	5.77	311976	11.96	3%
	(8875 to 161394)	(0.75 to 14.25)	(73750 to 608407)	(2.77 to 23.44)	(2.82 to 3.19)
Low-middle SDI	18653	2.83	75490	5.22	2.39%
	(3612 to 42686)	(0.54 to 6.52)	(17505 to 155025)	(1.19 to 10.78)	(2.29 to 2.49)
Low SDI	6319	2.43	16780	2.92	0.73%
	(893 to 16233)	(0.35 to 6.21)	(2643 to 41401)	(0.47 to 7.18)	(0.64 to 0.82)
Central Sub-Saharan Africa	796	3.15	1771	2.92	-0.17%
	(141 to 1950)	(0.57 to 7.77)	(274 to 4466)	(0.46 to 7.19)	(-0.48 to 0.14)
East Asia	83024	8.61	436252	20.8	3.72%
	(12430 to 200519)	(1.25 to 20.89)	(115075 to 813024)	(5.44 to 38.62)	(3.44 to 3.99)
Eastern Europe	77518	27.49	62825	18.92	-2.2%
	(20849 to 145612)	(7.36 to 51.58)	(11277 to 137394)	(3.42 to 41.18)	(-2.53 to -1.88)
Eastern Sub-Saharan Africa	2553	3.04	6912	3.71	0.79%
	(286 to 6777)	(0.35 to 8)	(738 to 18081)	(0.41 to 9.69)	(0.68 to 0.91)
Andean Latin America	912	4.18	3690	6.43	1.8%
	(96 to 2357)	(0.44 to 10.82)	(443 to 9188)	(0.76 to 16.05)	(1.66 to 1.93)
High-income Asia Pacific	22154	10.88	41772	10.94	-0.2%
	(2126 to 59218)	(1.05 to 29.1)	(5719 to 101560)	(1.65 to 25.79)	(-0.32 to -0.08)
High-income North America	110891	32.99	144668	25.35	-1.08%
	(32929 to 193179)	(9.96 to 57.08)	(42179 to 251558)	(7.51 to 43.92)	(-1.19 to -0.96)
Caribbean	1956	7.39	4608	8.91	0.87%
	(185 to 5166)	(0.7 to 19.52)	(437 to 12277)	(0.85 to 23.75)	(0.75 to 0.98)
Australasia	14698	63.82	18262	39.61	-1.91%
	(6625 to 22077)	(28.97 to 95.77)	(7951 to 28011)	(17.84 to 60.29)	(-2.08 to -1.74)
Central Europe	37988	25.72	63016	31.44	0.99%
	(8506 to 76660)	(5.77 to 51.88)	(16256 to 122005)	(8.24 to 60.21)	(0.84 to 1.14)


location	DALYs in 1990	ASDR per 100 000 in 1990	DALYs in 2019	ASDR per 100 000 in 2019	EAPC (1990– 2019)
Central Latin America	5030 (675 to 11890)	5.5 (0.72 to 13.21)	20610 (3627 to 48284)	8.47 (1.47 to 19.89)	1.59% (1.54 to 1.65)
Central Asia	8019 (2143 to 15197)	15.9 (4.21 to 30.17)	10914 (2929 to 21059)	13.59 (3.49 to 26.08)	-0.19% (-0.7 to 0.33)
North Africa and Middle East	8526 (802 to 23203)	4.5 (0.43 to 12.31)	23444 (2225 to 62629)	4.98 (0.48 to 13.24)	0.51% (0.25 to 0.77)
Oceania	187 (23 to 484)	5.47 (0.66 to 14.24)	454 (49 to 1206)	5.65 (0.59 to 15.03)	-0.02% (-0.14 to 0.1)
South Asia	7509 (2045 to 16221)	1.23 (0.32 to 2.67)	24347 (5853 to 55572)	1.66 (0.4 to 3.79)	0.86% (0.73 to 0.99)
Southeast Asia	11633 (1431 to 29882)	4.08 (0.51 to 10.45)	49492 (5926 to 123345)	7.56 (0.91 to 18.86)	2.1% (2.04 to 2.17)
Southern Latin America	20605 (9149 to 31548)	44.3 (19.61 to 67.88)	38064 (16077 to 58942)	46.7 (19.81 to 72.14)	0.17% (0.1 to 0.24)
Southern Sub-Saharan Africa	2216 (302 to 5423)	7.39 (0.98 to 18.32)	5485 (878 to 12736)	9.1 (1.4 to 21.36)	1.09% (0.91 to 1.27)
Tropical Latin America	10572 (2212 to 22079)	10.65 (2.16 to 22.52)	61167 (23985 to 97095)	24.61 (9.58 to 39.24)	3.15% (2.49 to 3.81)
Western Europe	198617 (60793 to 339982)	36.14 (11.18 to 61.52)	210073 (59872 to 372018)	25.63 (7.6 to 44.82)	-1.42% (-1.51 to -1.34)
Western Sub-Saharan Africa	2429 (262 to 6498)	2.61 (0.29 to 6.94)	6853 (714 to 18052)	3.39 (0.37 to 8.86)	1.16% (1.04 to 1.28)

On a global scale, regions with high SDI had the most deaths related to diet high in red meat (0.018 million) in 2019, while high-middle SDI regions had the highest DALYs (0.44 million), together accounting for over 35% of cases worldwide. However, the highest ASMR and ASDR were observed in high SDI regions. Over the years, ASMR and ASDR increased in low, low-middle, middle, and high-middle SDI regions, with middle and low-middle SDI regions experiencing a faster increase compared to low and high-middle SDI regions. In contrast, high SDI regions had a significant decrease in ASMR [EAPC -1.29% (95% CI: -1.37 to -1.21)] and ASDR [EAPC -1.2% (95% CI: -1.27 to -1.12)] (Table 1, 2).

At the regional level according to the Global Burden of Disease (GBD) classification, East Asia carried the heaviest burden in 2019, with 0.016 million deaths and 0.43 million DALYs worldwide. However, Australasia had the highest ASMR (1.75 per 100,000) and ASDR (39.61 per 100,000).

The most significant increase in ASMR and ASDR from 1990 to 2019 was observed in Tropical Latin America and East Asia, with Estimated Annual Percent Changes (EAPCs) exceeding 3 in both regions. On the other hand, Eastern Europe experienced the most substantial decrease in ASMR [EAPC -1.91% (95% CI: -2.21 to -1.62)] and ASDR [EAPC -2.2% (95% CI: -2.53 to -1.88)] (Table 1, 2).

At country level, China had the highest number of CRC deaths and DALYs attributed to a high red meat diet in 2019 (Table S1, S2). Argentina and Greenland had the highest ASMR and ASDR in 2019 (Figure 2A and B; Table S1, S2). However, the most substantial increase in ASMR and ASDR occurred in Equatorial Guinea, with EAPCs of 4.71% (95% CI: 4.27 to 5.14) and 4.31% (95% CI: 3.9 to 4.72), respectively (Figure S1; Table S1, S2).




**Figure 2.** These maps show the ASMR (A) and ASDR (B) of colorectal cancer due to diet high in red meat per 100,000 people in 2019 in 204 countries and territories, for both sexes. ASMR, age-standardized mortality rate; ASDR, age-standardized DALY rate.

### 3.2. Global colorectal cancer burden attributable to diet high in red meat by age and sex

In 2019, the number of deaths from CRC attributed to diet high in red meat exhibited a synchronous pattern in males and females, first rising and then declining with age. The peak age of incidence was in the 70-74 years group (Figure S2). More deaths were observed in the 65-74 years age range, with a higher number of age-specific deaths in males compared to females (Figure S2). Consequently, the age-specific mortality rate in males was greater than that in females, and showed a rapid increase for both sexes (Figure 3A).


The age-specific number of colorectal cancer DALYs followed a similar pattern to that of deaths, but the peak point appeared in the 65-69 years age group, with more DALYs occurring in the 60-69 years age range (Figure S2). The age-specific number of CRC DALYs was significantly higher in males than in females (Figure S2). Correspondingly, the age-specific DALY rate in males was larger

than that in females. The trend of the age-specific DALY rate resembled that of the mortality rate but declined in the 80-84 years age group (Figure 3B).



**Figure 3.** Age-specific rates of mortality (A) and DALYs (B) of colorectal cancer due to diet high in red meat by sex, 2019. DALYs, disability-adjusted life-years.


Globally, the age-specific mortality rate has increased in the 25-49 years and over 95 years age groups for both sexes from 1990 to 2019, with the most rapid increase occurring in the 25-29 years age group (Figure 4A). In males, the age-specific mortality rate rose in the 25-59 years and over 85 years age groups from 1990 to 2019, but fell in the 65-84 years age group, with the most significant increases and decreases occurring in the 65-69 years and 70-74 years age groups, respectively (Figure 4A). Conversely, the age-specific mortality rates in females declined in most age groups (Figure 4A). The age-specific mortality rates increased in low, low-middle, and middle SDI regions from 1990 to 2019. In high SDI regions, the age-specific mortality rates dropped in the 35-89 years age range, followed by an increase in older age groups. In high-middle SDI regions, the age-specific mortality rates decreased in the 60-69 years age group and increased in relatively younger and older age groups (Figure 4B). The EAPCs in age-specific DALY rates exhibited a similar pattern to that in age-specific mortality rates (Figure S3).



**Figure 4.** The age distribution of colorectal cancer due to a diet high in red meat trend in mortality rate from 1990 to 2019 by sex (A) and location (B). EAPC, estimated annual percentage change; SDI, sociodemographic index.


### 3.3. The association between SDI, global inequalities, and the burden of colorectal cancer burden attributable to diet high in red meat

In general, the ASMR displayed an inverted V-shaped correlation with the SDI, reaching its peak around an SDI value of approximately 0.75 (Figure 5A). The EAPC in ASMR exhibited a strong negative correlation with ASMR values in 1990 ( $\rho = -0.551$ ,  $P < 0.001$ ), particularly when ASMR was low (Figure 5B). Furthermore, the EAPC in ASMR showed a negative association with SDI values in 2019 ( $\rho = -0.216$ ,  $P < 0.001$ ), particularly for SDI values exceeding 0.75 (Figure 5C). Similar patterns were also evident when considering ASDR and their correlation with SDI, as well as the relationship between EAPC in ASDR and ASMR in 1990 and SDI in 2019 (Figure S2).



**Figure 5.** The correlation between colorectal cancer due to diet high in red meat in ASMR and SDI (A), between EAPC in ASMR and ASMR in 1990 (B), and between EAPC in ASMR and SDI in 2019 (C). EAPC, estimated annual percentage change; ASMR, age-standardized mortality rate; SDI, sociodemographic index.

In 1990 and 2019, 204 countries and territories experienced significant income-related inequalities in DALYs caused by CRC, the slope index of inequality in the DALYs rate between the highest and the lowest SDI country increased from 22.0 (95% CI 18.1 to 25.9) in 1990 to 32.9 (95% CI 28.3 to 37.5) in 2019 (Figure 6A; Table 3). The concentration index was 59.5 (95% CI 46.4 to 72.6) in 1990 and 48.9 (95% CI 34.6 to 63.1) in 2019 indicating that the burden was disproportionately concentrated in more-affluent countries (Figure 6B; Table 3).



**Figure 6.** Income-related health inequality regression (A) and concentration curves (B) for DALYs of colorectal cancer due to diet high in red meat across 204 counties and territories, 1990 and 2019. DALYs, disability-adjusted life-years.

**Table 3. Summary measures for cross-country inequalities related to SDI in DALYs of colorectal cancer burden attributable to diet high in red meat.**

Diseases	Health inequality metrics	Year	Value	95% CI
colon and rectum cancer	Slope index of inequality	1990	22.0	28.3 to 37.5
		2019	32.9	18.1 to 25.9
	Concentration index	1990	59.5	46.4 to 72.6
		2019	48.9	34.6 to 63.1

#### 4. Discussion

Our research analyzed the global burden of CRC caused by high red meat consumption in diets from 1990 to 2019, stratified by factors such as year, age, sex, region, and sociodemographic index. Our findings revealed a consistent global rise in mortality and DALYs rate linked to CRC due to elevated red meat consumption. This indicates that the burden it imposes remains a significant global public health concern.

In the 2019 report, there were over 50,000 reported cases of death, with the elderly being a high-risk group. Although the age-standardized death rate related to high red meat consumption in diets and CRC has declined from 1990 to 2019, the absolute number of fatalities has more than doubled. Population growth and age demographic shifts might account for this rise. Furthermore, the burden is closely related to socioeconomic development and how it is distributed unevenly. As high and high-middle SDI regions shoulder a greater burden, their ASDR and ASMR consistently decrease

annually. Specifically, Western Europe has seen significant reductions in ASMR and ASDR. Conversely, there has been a concerning increase in low SDI regions, such as tropical Latin America. No doubt, economic growth has brought more affordable red meat and shifted the disease burden pattern in low and middle-income countries from infectious diseases, maternal and neonatal diseases, and nutritional diseases to non-communicable chronic diseases[24, 34], with CRC importantly of noteworthy. Additionally, the escalating male-to-female ratio in CRC-related deaths indicates that males are being affected to a greater extent. The reason for this may be that males are more likely to consume unprocessed red meat, leading to higher estimated intake levels[35]. Another hypothesis is that dietary-related effects may vary by sex due to hormonal differences between males and females, as well as the tendency for females to develop proximal tumors and males to develop distal and rectal tumors[36].

Meat processing, preservation, and high-temperature cooking can yield carcinogens such as HCAs, PAHs, and NOCs, which are implicated in the onset of several cancers[37-40]. Furthermore, red meat is a major source of heme iron. Many epidemiological and evidence-based studies have suggested a possible association between the intake of heme iron and the risk of various cancers, including colorectal cancer[41, 42]. Compared to low-income countries, high-income countries have a red meat consumption level that is approximately five times higher[43]. Despite the health inequality analysis indicating that the burden is disproportionately concentrated in wealthier countries, high-income countries have exhibited reduced red meat consumption over recent decades. This decline is primarily due to the development of relevant dietary guidelines and an increased level of compliance with these guidelines. Additionally, people are becoming more aware of the negative health effects of red meat consumption[44-46]. Another study has demonstrated the merit of public health initiatives that limit red meat intake in developed countries[47]. Therefore, high-income countries should emphasize the benefits of a healthy diet and use evidence-based intervention measures to educate and empower individuals to understand the risks associated with excessive red meat consumption[23, 48]. At the same time, although the burden in developing countries is relatively smaller, deaths related to CRC are rapidly increasing. From a nutritional perspective, animal-source foods (ASF) including red meat remain important in developing countries[43]. In these nations, it might not be necessary to completely avoid the consumption of red meat[12, 49, 50]. Most African countries are low-income or middle-income nations and their people do not consume red meat, even at potentially beneficial levels and this can contribute to severe nutritional deficiencies associated with a low life expectancy[51, 52]. For such countries, vigilance should not be relaxed, as paradoxically, excessive red meat consumption can eventually burden these nations[22]. Policymakers must remain cognizant of this risk.

Furthermore, though excessive red meat consumption is unhealthy, our study did not directly link it to CRC causality. We could only delineate the present burden and trends. Our study utilized data from GBD 2019, which primarily consists of national and regional data. The accuracy of our research hinges on the GBD study's data quality and volume. We lack individual-level data to support our findings. In addition, the highest consumers of red meat (the top 20% of red meat consumers) also have the highest body mass index, are more likely to smoke, engage in less physical activity, generally have lower levels of education and health literacy, consume smaller quantities of fruits and vegetables, and have higher daily calorie intake. Any or a combination of these residual confounding factors could potentially influence the association between red meat intake and colorectal cancer[16].



## **5. Conclusions**

Although the ASIR and ASDR of CRC associated with diet high in red meat have decreased globally from 1990 to 2019, the absolute number of cases is still on the rise. The number of deaths has more than doubled since 1990, with a greater impact on males and older individuals. CRC linked to diet high in red meat exhibited significant income-related inequality, disproportionately burdening more affluent countries. However, in developing countries, including some countries in tropical Latin America and Africa, ASIR and ASDR related to cancer were higher. We hope that our research can provide insights for policymakers to tailor dietary guidelines, disease control, and prevention strategies based on the specific circumstances of different countries, especially for least developed countries with relatively lower economic levels.

## **Author Contributions**

All authors contributed to the manuscript presented methodology, conceptualization, data analysis, and manuscript writing.

## **Funding**

This project was supported by the Hainan Province Clinical Medical Center and the Key Research and Development Program of Hainan Province (ZDYF2021SHFZ087).

## **Institutional Review Board Statement**

The study complied with the Helsinki Declaration and has been approved by the Medical Ethics Committee of Tongji Hospital.

## **Informed Consent Statement**

The institutions responsible for conducting the surveys were obligated to obtain informed consent from the participants.

## **Data Availability Statement**

The data is accessible through a public, open-access repository. All data is publicly available, and it can also be obtained upon request.

## **Conflicts of Interest**

The authors declare no conflict of interest.

## **Abbreviations**

CI: confidence interval; DALYs, disability-adjusted life years; EAPC: estimated annual percent change; GBD: disability-adjusted life years; SDI: sociodemographic index; UI: uncertainty interval; ASMR= age standardized mortality rate; ASDR= age-standardized disability adjusted life year rate; WHO: World Health Organization; CRC: colorectal cancer; HCA: heterocyclic amines; PAHs: polycyclic aromatic hydrocarbons; NOCs: N-nitroso compounds.

## **References**

- [1] Yang VW. The molecular genetics of colorectal cancer. *Curr Gastroenterol Rep* 1999; 1:

449-454.

- [2] Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D and Bray F. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 2015; 136: E359-386.
- [3] Huxley RR, Ansary-Moghaddam A, Clifton P, Czernichow S, Parr CL and Woodward M. The impact of dietary and lifestyle risk factors on risk of colorectal cancer: a quantitative overview of the epidemiological evidence. *Int J Cancer* 2009; 125: 171-180.
- [4] Dekker E, Tanis PJ, Vleugels JLA, Kasi PM and Wallace MB. Colorectal cancer. *Lancet* 2019; 394: 1467-1480.
- [5] Russo MW, Wei JT, Thiny MT, Gangarosa LM, Brown A, Ringel Y, Shaheen NJ and Sandler RS. Digestive and liver diseases statistics, 2004. *Gastroenterology* 2004; 126: 1448-1453.
- [6] Jemal A, Tiwari RC, Murray T, Ghafoor A, Samuels A, Ward E, Feuer EJ and Thun MJ. Cancer statistics, 2004. *CA Cancer J Clin* 2004; 54: 8-29.
- [7] Farvid MS, Sidahmed E, Spence ND, Mante Angua K, Rosner BA and Barnett JB. Consumption of red meat and processed meat and cancer incidence: a systematic review and meta-analysis of prospective studies. *Eur J Epidemiol* 2021; 36: 937-951.
- [8] Bouvard V, Loomis D, Guyton KZ, Grosse Y, Ghissassi FE, Benbrahim-Tallaa L, Guha N, Mattock H and Straif K. Carcinogenicity of consumption of red and processed meat. *Lancet Oncol* 2015; 16: 1599-1600.
- [9] Speedy AW. Global production and consumption of animal source foods. *J Nutr* 2003; 133: 4048s-4053s.
- [10] Wang X, Lin X, Ouyang YY, Liu J, Zhao G, Pan A and Hu FB. Red and processed meat consumption and mortality: dose-response meta-analysis of prospective cohort studies. *Public Health Nutr* 2016; 19: 893-905.
- [11] Lafarga T and Hayes M. Bioactive peptides from meat muscle and by-products: generation, functionality and application as functional ingredients. *Meat Sci* 2014; 98: 227-239.
- [12] Walker P, Rhubart-Berg P, McKenzie S, Kelling K and Lawrence RS. Public health implications of meat production and consumption. *Public Health Nutr* 2005; 8: 348-356.
- [13] Wolk A. Potential health hazards of eating red meat. *J Intern Med* 2017; 281: 106-122.
- [14] Ekmekcioglu C, Wallner P, Kundi M, Weisz U, Haas W and Hutter HP. Red meat, diseases, and healthy alternatives: A critical review. *Crit Rev Food Sci Nutr* 2018; 58: 247-261.
- [15] Qian F, Riddle MC, Wylie-Rosett J and Hu FB. Red and Processed Meats and Health Risks: How Strong Is the Evidence? *Diabetes Care* 2020; 43: 265-271.
- [16] Cross AJ, Leitzmann MF, Gail MH, Hollenbeck AR, Schatzkin A and Sinha R. A prospective study of red and processed meat intake in relation to cancer risk. *PLoS Med* 2007; 4: e325.
- [17] Giovannucci E, Rimm EB, Stampfer MJ, Colditz GA, Ascherio A and Willett WC. Intake of fat, meat, and fiber in relation to risk of colon cancer in men. *Cancer Res* 1994; 54: 2390-2397.
- [18] Kontogianni MD, Panagiotakos DB, Pitsavos C, Chrysohoou C and Stefanadis C. Relationship between meat intake and the development of acute coronary syndromes: the CARDIO2000 case-control study. *Eur J Clin Nutr* 2008; 62: 171-177.
- [19] Armstrong B and Doll R. Environmental factors and cancer incidence and mortality in different countries, with special reference to dietary practices. *Int J Cancer* 1975; 15: 617-

631.

- [20] Huang Y, Cao D, Chen Z, Chen B, Li J, Guo J, Dong Q, Liu L and Wei Q. Red and processed meat consumption and cancer outcomes: Umbrella review. *Food Chem* 2021; 356: 129697.
- [21]
- [22] Liu D, Shi Q, Cheng G, Huang Q and Li S. Worldwide burden attributable to diet high in red meat from 1990 to 2019. *Arch Med Sci* 2023; 19: 1-15.
- [23] Wu Z, Wang W, Zhang K, Fan M and Lin R. The Global Burden of Disease Attributable to Diet High in Red Meat in 204 Countries and Territories, 1999-2019: An updated Analysis of the Global Burden of Disease Study. *Mol Nutr Food Res* 2023; e2300144.
- [24] Global burden of 369 diseases and injuries in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* 2020; 396: 1204-1222.
- [25] Global burden of 87 risk factors in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* 2020; 396: 1223-1249.
- [26] Global age-sex-specific fertility, mortality, healthy life expectancy (HALE), and population estimates in 204 countries and territories, 1950-2019: a comprehensive demographic analysis for the Global Burden of Disease Study 2019. *Lancet* 2020; 396: 1160-1203.
- [27] Kocarnik JM, Compton K, Dean FE, Fu W, Gaw BL, Harvey JD, Henrikson HJ, Lu D, Pennini A, Xu R, Ababneh E, Abbasi-Kangevari M, Abbastabar H, Abd-Elsalam SM, Abdoli A, Abedi A, Abidi H, Abolhassani H, Adedeji IA, Adnani QES, Advani SM, Afzal MS, Aghaali M, Ahinkorah BO, Ahmad S, Ahmad T, Ahmadi A, Ahmadi S, Ahmed Rashid T, Ahmed Salih Y, Akalu GT, Aklilu A, Akram T, Akunna CJ, Al Hamad H, Alahdab F, Al-Aly Z, Ali S, Alimohamadi Y, Alipour V, Aljunid SM, Alkhayyat M, Almasi-Hashiani A, Almasri NA, Al-Maweri SAA, Almustanyir S, Alonso N, Alvis-Guzman N, Amu H, Anbesu EW, Ancuceanu R, Ansari F, Ansari-Moghaddam A, Antwi MH, Anvari D, Anyasodor AE, Aqeel M, Arabloo J, Arab-Zozani M, Aremu O, Ariffin H, Aripov T, Arshad M, Artaman A, Arulappan J, Asemi Z, Asghari Jafarabadi M, Ashraf T, Atorkey P, Aujayeb A, Ausloos M, Awedew AF, Ayala Quintanilla BP, Ayenew T, Azab MA, Azadnajafabad S, Azari Jafari A, Azarian G, Azzam AY, Badiye AD, Bahadory S, Baig AA, Baker JL, Balakrishnan S, Banach M, Bärnighausen TW, Barone-Adesi F, Barra F, Barrow A, Behzadifar M, Belgaumi UI, Bezabhe WMM, Bezabih YM, Bhagat DS, Bhagavathula AS, Bhardwaj N, Bhardwaj P, Bhaskar S, Bhattacharyya K, Bhojaraja VS, Bibi S, Bijani A, Biondi A, Bisignano C, Bjørge T, Bleyer A, Blyuss O, Bolarinwa OA, Bolla SR, Braithwaite D, Brar A, Brenner H, Bustamante-Teixeira MT, Butt NS, Butt ZA, Caetano Dos Santos FL, Cao Y, Carreras G, Catalá-López F, Cembranel F, Cerin E, Cernigliaro A, Chakinala RC, Chattu SK, Chattu VK, Chaturvedi P, Chimed-Ochir O, Cho DY, Christopher DJ, Chu DT, Chung MT, Conde J, Cortés S, Cortesi PA, Costa VM, Cunha AR, Dadras O, Dagnew AB, Dahlawi SMA, Dai X, Dandona L, Dandona R, Darwesh AM, das Neves J, De la Hoz FP, Demis AB, Denova-Gutiérrez E, Dhamnetiya D, Dhimal ML, Dhimal M, Dianatinasab M, Diaz D, Djalalinia S, Do HP, Doaei S, Dorostkar F, Dos Santos Figueiredo FW, Driscoll TR, Ebrahimi H, Eftekharzadeh S, El Tantawi M, El-Abid H, Elbarazi I, Elhabashy HR, Elhadi M, El-Jaafary SI, Eshrati B, Eskandarieh S, Esmaeilzadeh F, Etemadi A, Ezzikouri S, Faisaluddin M, Faraon EJA, Fares J, Farzadfar F, Feroze AH, Ferrero S, Ferro Desideri L, Filip I, Fischer F, Fisher JL, Foroutan M, Fukumoto T, Gaal PA, Gad MM, Gadanya MA, Gallus S, Gaspar Fonseca M, Getachew Obsa A, Ghafourifard M, Ghashghaee A, Ghith

N, Gholamalizadeh M, Gilani SA, Ginindza TG, Gizaw ATT, Glasbey JC, Golechha M, Goleij P, Gomez RS, Gopalani SV, Gorini G, Goudarzi H, Grosso G, Gubari MIM, Guerra MR, Guha A, Gunasekera DS, Gupta B, Gupta VB, Gupta VK, Gutiérrez RA, Hafezi-Nejad N, Haider MR, Haj-Mirzaian A, Halwani R, Hamadeh RR, Hameed S, Hamidi S, Hanif A, Haque S, Harlianto NI, Haro JM, Hasaballah AI, Hassanipour S, Hay RJ, Hay SI, Hayat K, Heidari G, Heidari M, Herrera-Serna BY, Herteliu C, Hezam K, Holla R, Hossain MM, Hossain MBH, Hosseini MS, Hosseini M, Hosseinzadeh M, Hostiuc M, Hostiuc S, Househ M, Hsairi M, Huang J, Hugo FN, Hussain R, Hussein NR, Hwang BF, Iavicoli I, Ibitoye SE, Ida F, Ikuta KS, Ilesanmi OS, Ilic IM, Ilic MD, Irham LM, Islam JY, Islam RM, Islam SMS, Ismail NE, Isola G, Iwagami M, Jacob L, Jain V, Jakovljevic MB, Javaheri T, Jayaram S, Jazayeri SB, Jha RP, Jonas JB, Joo T, Joseph N, Joukar F, Jürisson M, Kabir A, Kahrizi D, Kalankesh LR, Kalhor R, Kaliyadan F, Kalkonde Y, Kamath A, Kameran AI-Salihi N, Kandel H, Kapoor N, Karch A, Kasa AS, Katikireddi SV, Kauppila JH, Kavetsky T, Kebede SA, Keshavarz P, Keykhaei M, Khader YS, Khalilov R, Khan G, Khan M, Khan MN, Khan MAB, Khang YH, Khater AM, Khayamzadeh M, Kim GR, Kim YJ, Kisa A, Kisa S, Kissimova-Skarbek K, Kopec JA, Koteeswaran R, Koul PA, Koulmane Laxminarayana SL, Koyanagi A, Kucuk Bicer B, Kugbey N, Kumar GA, Kumar N, Kumar N, Kurmi OP, Kutluk T, La Vecchia C, Lami FH, Landires I, Lauriola P, Lee SW, Lee SWH, Lee WC, Lee YH, Leigh J, Leong E, Li J, Li MC, Liu X, Loureiro JA, Lunevicius R, Magdy Abd El Razek M, Majeed A, Makki A, Male S, Malik AA, Mansournia MA, Martini S, Masoumi SZ, Mathur P, McKee M, Mehrotra R, Mendoza W, Menezes RG, Mengesha EW, Mesregah MK, Mestrovic T, Miao Jonasson J, Miazgowski B, Miazgowski T, Michalek IM, Miller TR, Mirzaei H, Mirzaei HR, Misra S, Mithra P, Moghadaszadeh M, Mohammad KA, Mohammad Y, Mohammadi M, Mohammadi SM, Mohammadian-Hafshejani A, Mohammed S, Moka N, Mokdad AH, Molokhia M, Monasta L, Moni MA, Moosavi MA, Moradi Y, Moraga P, Morgado-da-Costa J, Morrison SD, Mosapour A, Mubarik S, Mwanri L, Nagarajan AJ, Nagaraju SP, Nagata C, Naimzada MD, Nangia V, Naqvi AA, Narasimha Swamy S, Ndejjo R, Nduaguba SO, Negoï I, Negru SM, Neupane Kandel S, Nguyen CT, Nguyen HLT, Niazi RK, Nnaji CA, Noor NM, Nuñez-Samudio V, Nzopotam CI, Oancea B, Ochir C, Odukoya OO, Ogbo FA, Olagunju AT, Olakunde BO, Omar E, Omar Bali A, Omonisi AEE, Ong S, Onwujekwe OE, Orru H, Ortega-Altamirano DV, Otstavnov N, Otstavnov SS, Owolabi MO, P AM, Padubidri JR, Pakshir K, Pana A, Panagiotakos D, Panda-Jonas S, Pardhan S, Park EC, Park EK, Pashazadeh Kan F, Patel HK, Patel JR, Pati S, Pattanshetty SM, Paudel U, Pereira DM, Pereira RB, Perianayagam A, Pillay JD, Pirouzpanah S, Pishgar F, Podder I, Postma MJ, Pourjafar H, Prashant A, Preotescu L, Rabiee M, Rabiee N, Radfar A, Radhakrishnan RA, Radhakrishnan V, Rafiee A, Rahim F, Rahimzadeh S, Rahman M, Rahman MA, Rahmani AM, Rajai N, Rajesh A, Rakovac I, Ram P, Ramezanzadeh K, Ranabhat K, Ranasinghe P, Rao CR, Rao SJ, Rawassizadeh R, Razeghinia MS, Renzaho AMN, Rezaei N, Rezaei N, Rezapour A, Roberts TJ, Rodriguez JAB, Rohloff P, Romoli M, Ronfani L, Roshandel G, Rweggerera GM, S M, Sabour S, Saddik B, Saeed U, Sahebkar A, Sahoo H, Salehi S, Salem MR, Salimzadeh H, Samaei M, Samy AM, Sanabria J, Sankararaman S, Santric-Milicevic MM, Sardiwalla Y, Sarveazad A, Sathian B, Sawhney M, Saylan M, Schneider IJC, Sekerija M, Seylani A, Shafaat O, Shaghaghzi Z, Shaikh MA, Shamsoddin E, Shannawaz M, Sharma R, Sheikh A, Sheikhbahaei S, Shetty A, Shetty JK, Shetty PH, Shibuya K, Shirkoohi R, Shivakumar KM, Shivarov V, Siabani S, Siddappa

Malleshappa SK, Silva DAS, Singh JA, Sintayehu Y, Skryabin VY, Skryabina AA, Soeberg MJ, Sofi-Mahmudi A, Sotoudeh H, Steiropoulos P, Straif K, Subedi R, Sufiyan MB, Sultan I, Sultana S, Sur D, Szerencsés V, Szócska M, Tabarés-Seisdedos R, Tabuchi T, Tadbiri H, Taherkhani A, Takahashi K, Talaat IM, Tan KK, Tat VY, Tedla BAA, Tefera YG, Tehrani-Banihashemi A, Temsah MH, Tesfay FH, Tessema GA, Thapar R, Thavamani A, Thoguluva Chandrasekar V, Thomas N, Tohidinik HR, Touvier M, Tovani-Palome MR, Traini E, Tran BX, Tran KB, Tran MTN, Tripathy JP, Tusa BS, Ullah I, Ullah S, Umapathi KK, Unnikrishnan B, Upadhyay E, Vacante M, Vaezi M, Valadan Tahbaz S, Velazquez DZ, Veroux M, Violante FS, Vlassov V, Vo B, Volovici V, Vu GT, Waheed Y, Wamai RG, Ward P, Wen YF, Westerman R, Winkler AS, Yadav L, Yahyazadeh Jabbari SH, Yang L, Yaya S, Yazie TSY, Yeshaw Y, Yonemoto N, Younis MZ, Yousefi Z, Yu C, Yuze D, Yunusa I, Zadnik V, Zare F, Zastrozhin MS, Zastrozhina A, Zhang J, Zhong C, Zhou L, Zhu C, Ziapour A, Zimmermann IR, Fitzmaurice C, Murray CJL and Force LM. Cancer Incidence, Mortality, Years of Life Lost, Years Lived With Disability, and Disability-Adjusted Life Years for 29 Cancer Groups From 2010 to 2019: A Systematic Analysis for the Global Burden of Disease Study 2019. *JAMA Oncol* 2022; 8: 420-444.

- [28] Health effects of dietary risks in 195 countries, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 2019; 393: 1958-1972.
- [29] Gao S, Yang WS, Bray F, Va P, Zhang W, Gao J and Xiang YB. Declining rates of hepatocellular carcinoma in urban Shanghai: incidence trends in 1976-2005. *Eur J Epidemiol* 2012; 27: 39-46.
- [30] Zhou L, Deng Y, Li N, Zheng Y, Tian T, Zhai Z, Yang S, Hao Q, Wu Y, Song D, Zhang D, Lyu J and Dai Z. Global, regional, and national burden of Hodgkin lymphoma from 1990 to 2017: estimates from the 2017 Global Burden of Disease study. *J Hematol Oncol* 2019; 12: 107.
- [31] Organization WH. Handbook on health inequality monitoring: With a special focus on low-and middle-income countries. 2013;
- [32] Peng J, Xu H and Tang X. Global inequalities in the burden of digestive diseases from 1990 to 2019: findings from the Global Burden of Disease Study 2019. *Gastroenterology* 2023;
- [33] Ordunez P, Martinez R, Soliz P, Giraldo G, Mujica OJ and Nordet P. Rheumatic heart disease burden, trends, and inequalities in the Americas, 1990-2017: a population-based study. *Lancet Glob Health* 2019; 7: e1388-e1397.
- [34] Lippi G and Mattiuzzi C. The global burden of pancreatic cancer. *Arch Med Sci* 2020; 16: 820-824.
- [35] Frank SM, Jaacks LM, Batis C, Vanderlee L and Taillie LS. Patterns of Red and Processed Meat Consumption across North America: A Nationally Representative Cross-Sectional Comparison of Dietary Recalls from Canada, Mexico, and the United States. *Int J Environ Res Public Health* 2021; 18:
- [36] Jacobs ET, Thompson PA and Martínez ME. Diet, gender, and colorectal neoplasia. *J Clin Gastroenterol* 2007; 41: 731-746.
- [37] Cross AJ, Freedman ND, Ren J, Ward MH, Hollenbeck AR, Schatzkin A, Sinha R and Abnet CC. Meat consumption and risk of esophageal and gastric cancer in a large prospective study. *Am J Gastroenterol* 2011; 106: 432-442.
- [38] Ferguson LR. Natural and human-made mutagens and carcinogens in the human diet.

- Toxicology 2002; 181-182: 79-82.
- [39] Skog KI, Johansson MA and Jägerstad MI. Carcinogenic heterocyclic amines in model systems and cooked foods: a review on formation, occurrence and intake. *Food Chem Toxicol* 1998; 36: 879-896.
- [40] Tricker AR and Preussmann R. Carcinogenic N-nitrosamines in the diet: occurrence, formation, mechanisms and carcinogenic potential. *Mutat Res* 1991; 259: 277-289.
- [41] Carr PR, Walter V, Brenner H and Hoffmeister M. Meat subtypes and their association with colorectal cancer: Systematic review and meta-analysis. *Int J Cancer* 2016; 138: 293-302.
- [42] Cross AJ, Ferrucci LM, Risch A, Graubard BI, Ward MH, Park Y, Hollenbeck AR, Schatzkin A and Sinha R. A large prospective study of meat consumption and colorectal cancer risk: an investigation of potential mechanisms underlying this association. *Cancer Res* 2010; 70: 2406-2414.
- [43] Ranabhat CL, Park MB and Kim CB. Influence of Alcohol and Red Meat Consumption on Life Expectancy: Results of 164 Countries from 1992 to 2013. *Nutrients* 2020; 12:
- [44] Zeng L, Ruan M, Liu J, Wilde P, Naumova EN, Mozaffarian D and Zhang FF. Trends in Processed Meat, Unprocessed Red Meat, Poultry, and Fish Consumption in the United States, 1999-2016. *J Acad Nutr Diet* 2019; 119: 1085-1098.e1012.
- [45] Stewart C, Piernas C, Cook B and Jebb SA. Trends in UK meat consumption: analysis of data from years 1-11 (2008-09 to 2018-19) of the National Diet and Nutrition Survey rolling programme. *Lancet Planet Health* 2021; 5: e699-e708.
- [46] Clonan A, Roberts KE and Holdsworth M. Socioeconomic and demographic drivers of red and processed meat consumption: implications for health and environmental sustainability. *Proc Nutr Soc* 2016; 75: 367-373.
- [47] Park MB. Effect of red meat, vegetable, tobacco, and alcohol consumption on national cancer mortality index: Data from 1989 to 2013 in 37 developed countries. *Front Nutr* 2022; 9: 929553.
- [48] Micha R, Wallace SK and Mozaffarian D. Red and processed meat consumption and risk of incident coronary heart disease, stroke, and diabetes mellitus: a systematic review and meta-analysis. *Circulation* 2010; 121: 2271-2283.
- [49] Larsen CS. Animal source foods and human health during evolution. *J Nutr* 2003; 133: 3893s-3897s.
- [50] Popkin BM. Reducing meat consumption has multiple benefits for the world's health. *Arch Intern Med* 2009; 169: 543-545.
- [51] Fanzo J, Hawkes C, Udomkesmalee E, Afshin A, Allemandi L, Assery O, Baker P, Battersby J, Bhutta Z and Chen K. 2018 Global Nutrition Report: Shining a light to spur action on nutrition. 2018;
- [52] Ranabhat CL, Jakovljevic M, Dhimal M and Kim CB. Structural Factors Responsible for Universal Health Coverage in Low- and Middle-Income Countries: Results From 118 Countries. *Front Public Health* 2019; 7: 414.