

Qualitative Health Research

Women's Experiences of Coping With Pregnancy Termination for Fetal Abnormality

Journal:	<i>Qualitative Health Research</i>
Manuscript ID:	QHR-2012-0296.R2
Manuscript Type:	Research Article
Keywords:	abortion, bereavement / grief, coping and adaptation, interpretative phenomenological analysis (IPA), lived experience, perinatal health, pregnancy, psychology, research, qualitative

SCHOLARONE™
Manuscripts

Review

1
2
3 Women's Experiences of Coping With Pregnancy Termination for Fetal Abnormality
4

5 Caroline Lafarge,¹ Kathryn Mitchell,¹ and Pauline Fox¹
6

7
8 ¹University of West London, London, UK.
9

10 **Corresponding Author:**

11
12 Caroline Lafarge, University of West London, Paragon, Boston Manor Road, Brentford
13

14
15 TW8 9GA, UK
16

17
18 Email: caroline.lafarge@uwl.ac.uk
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Abstract

Pregnancy termination for fetal abnormality (TFA) can have significant psychological consequences. Most research focus on measuring the psychological outcome of TFA, and little is known about the coping strategies involved. In this article, we report women's coping strategies used during and after the procedure. Our account is based on experiences of 27 women, which we analyzed using interpretative phenomenological analysis. Coping comprised four structures, consistent across time points: support, acceptance, avoidance, and meaning attribution. Women mostly used "adaptive" coping strategies but reported inadequacies in aftercare, which challenged their resources. The implications of this study focus on the provision of a sensitive, nondirective care rooted in the acknowledgment of the unique nature of TFA. Enabling women to reciprocate to emotional support, promoting adaptive coping strategies, highlighting the potential value of spending time with the baby, and providing long-term support (including during subsequent pregnancies) might promote psychological adjustment to TFA.

Keywords

Abortion; bereavement / grief; coping and adaptation; interpretative phenomenological analysis (IPA); lived experience; perinatal health; pregnancy; psychology; research, qualitative

1
2
3 Progresses in antenatal screening and delayed maternal age have resulted in the increased
4
5 detection of fetal abnormalities and terminations for fetal abnormality (TFA; Korenromp et al.,
6
7 2007). For example, in England and Wales in 2010, TFA represented 1% of all terminations
8
9 (Department of Health, 2011). Despite being small, this number is likely to rise as more couples
10
11 delay starting a family, thus increasing the risk of fetal abnormalities occurring (Royal College
12
13 of Obstetricians and Gynaecologists, 2009). TFA is at the center of many ethical and clinical
14
15 debates. First, it is closely associated with the abortion debate and the question of the impact of
16
17 abortion on women's mental health. Evidence on the subject is inconsistent with some reviews
18
19 indicating a negative impact of abortion (Coleman, 2011), whereas others suggest otherwise
20
21 (Major et al., 2009; National Collaborating Centre for Mental Health, 2011). TFA is also linked
22
23 to the eugenics debate, as demonstrated by arguments about how late in the pregnancy and for
24
25 which medical conditions could TFA be performed (House of Commons, 2007a,b; the Heartbeat
26
27 bill, nd). Ethical questioning also occurs at an individual level; first, for the women who might
28
29 be challenged by the morality of their decision (McCoyd, 2007). It also extends to the health
30
31 professionals who might object to these procedures (Marshall & Raynor, 2002; Strickland,
32
33 2011). These debates have legal, political, and clinical implications, which affect the care
34
35 provided to women and their psychosocial well-being.
36
37
38
39
40
41
42

43
44 Much research has centered on the decision making following antenatal screening,
45
46 particularly when a diagnosis of fetal abnormality is made. Evidence suggests that this is a
47
48 complex process (McCoyd, 2008; Sandelowski & Barroso, 2005). Many researchers have
49
50 attempted to understand the psychological impact of TFA on women's mental health, and the
51
52 literature suggests that TFA can have long lasting psychological consequences for those
53
54 involved. Occurrences of complicated grief, depression, and posttraumatic stress following TFA
55
56 have been well documented (Davies, Gledhill, McFadyen, Whitlow, & Economides, 2005; Green
57
58
59
60

1
2
3 & Statham, 2007; Kersting et al., 2005; Korenromp et al., 2007). Complications are particularly
4
5 manifest among mothers, but accounts of psychological distress have also been reported among
6
7 fathers (Robson, 2002). Grief reactions following TFA have been shown to resemble those
8
9 experienced in other types of perinatal loss (Keefe-Cooperman, 2004; Salvesen, Oyen, Schmidt,
10
11 Malt, & Eik-Nes, 1997). Still, ending a pregnancy for fetal abnormality bears an additional moral
12
13 component in that the parents choose to terminate a pregnancy. Research indicates that several
14
15 years post-TFA, women continue to display symptoms of grief (Green & Statham, 2007). Thus,
16
17 like other types of bereavement, the process of coping with TFA is ongoing.
18
19

20
21
22 To date, little is known about the coping processes involved in dealing with TFA, which
23
24 we aim to address in this article. Coping is a relevant field of investigation within the context of
25
26 TFA. Research indicates that people who face similar stressful events adjust in different ways,
27
28 and coping processes influence psychological outcomes (Carver & Connor-Smith, 2010). Many
29
30 research studies have emphasized the role of coping in adjusting to stressful situations among
31
32 various populations (Cartwright, Endean, & Porter, 2009; Schnider, Elhai, & Gray, 2007).
33
34 Lazarus and Folkman (1984) defined coping as psychological processes that occur when facing
35
36 stressful situations, comprising appraisal of the threat (perception and evaluation) and coping *per*
37
38 *se* (resources involved in managing the problem). Coping consists of appraising the threat to
39
40 reduce its magnitude and its subsequent impact on the individual (Carver & Connor-Smith,
41
42 2010).
43
44
45
46
47

48
49 There are many ways of categorizing coping processes from approach/avoidance coping
50
51 (Roth & Cohen, 1986), problem/emotional focused coping (Lazarus & Folkman, 1984), through
52
53 to the multidimensional approach to coping that includes processes, such as behavioral
54
55 disengagement, religion, and acceptance (Carver, 1997). More recently, researchers have
56
57 identified five core categories of coping: problem solving, support seeking, distraction,
58
59
60

1
2
3 avoidance, and positive restructuring (Skinner, Edge, Altman, & Sherwood, 2003). Additional
4
5 categories have also emerged such as accommodative, meaning-focused, and proactive coping
6
7 (Carver & Connor-Smith, 2010). Given the influence of coping on psychological adaptation, it is
8
9 important to understand the coping strategies women use when they undergo TFA, during the
10
11 termination procedure itself and afterwards. In this article, we report a qualitative study of 27
12
13 women who underwent TFA. To our knowledge, this is the first time that an investigation
14
15 focused exclusively on coping strategies employed both during and after the procedure.
16
17
18
19

20 **Methods**

21
22 We conducted a cross-sectional retrospective study to explore women's coping strategies during
23
24 and after TFA. The qualitative interview schedule consisted of open-ended questions about the
25
26 process of coping with the termination, with specific attention paid to coping strategies. We left
27
28 the questions open and provided few prompts to ensure that the feedback collected was, as much
29
30 as possible, participant-led (Biggerstaff & Thompson, 2008). Topics such as what the
31
32 termination meant for the participants and how they felt about the future were also investigated.
33
34 We collected the following information about the terminated pregnancy: number of pregnancies,
35
36 existence of other children at the time of TFA, gestational age, reproductive history post-TFA,
37
38 method of abnormality detection (routine scan or directed search), abnormality prognosis
39
40 (incompatible or compatible with life), method of termination (medical, surgical, or induced
41
42 labor), and how participants felt about their decision to end the pregnancy. Demographic data
43
44 were also gathered.
45
46
47
48
49

50
51 Participants were members of a support group based in the United Kingdom, which
52
53 provides support to parents when an abnormality is detected in their unborn baby. We recruited
54
55 the participants through the group's e-mail network and forum. We provided them with
56
57 information about the study (e.g., aims, confidentiality, and right to withdraw) and the
58
59
60

1
2
3 opportunity to ask questions before we formally obtained their informed consent. To be eligible,
4
5 women had to be at least 18 years old and, initially, they had to have undergone TFA at least six
6
7 months prior to participating in the survey. We later broadened the latter criterion to include any
8
9 women who had undergone TFA irrespective of when the procedure took place, in light of
10
11 comments from potential participants who were willing but unable to take part because of the six
12
13 months exclusion criterion. Altogether, 27 women participated in this study. We decided to stop
14
15 collecting data after 27 accounts, given that saturation of themes was reached at 25. Participants
16
17 were mainly White, and two-thirds had completed higher education. Their ages ranged between
18
19 24 and 44 years. Pregnancies were terminated between 12- and 30-weeks of gestation. For 13
20
21 participants, this was their first pregnancy, whereas two had undergone three TFAs.
22
23
24
25
26

27 The data collection took place online between April 2011 and February 2012.
28
29 Participants had the flexibility to leave and re-enter the survey, enabling them to complete it in
30
31 their own time and pace. Collecting data electronically presented a number of advantages, not
32
33 least enhancing confidentiality with no intervention from the researcher in the data collection.
34
35 We also believe that answering questions online might have enabled women to open up more
36
37 easily than in a face-to-face environment and to think carefully about their answers, thus
38
39 providing valuable insight. At the end of the survey, women had the opportunity to comment on
40
41 the study and leave their contact details if they wished to receive a summary of the findings. We
42
43 conducted a pilot study with three participants, which resulted in no changes being made.
44
45
46
47 However, one respondent suggested that a preview of the questions would be helpful to start the
48
49 reflection process prior to filling the survey, which we subsequently implemented.
50
51
52

53 We used interpretative phenomenological analysis (IPA) to analyze the data. IPA is
54
55 particularly suited to this type of analysis because it uncovers people's individual experience and
56
57 the meaning they attribute to it. Thus, it is deeply rooted in individual narratives and represents a
58
59
60

1
2
3 bottom-up rather than a top-down analytical approach. In analyzing the data using IPA, the
4
5 researcher also seeks to provide an interpretation of these narratives (Larkin & Thompson,
6
7 2012). IPA is most commonly used to analyze data collected through semi-structured interviews;
8
9 however, other types of data have also been successfully used in conjunction with IPA. These
10
11 include diaries, personal accounts, letters, questionnaires with open-ended questions aiming to
12
13 capture additional comments, internet forum posts (Mulveen & Hepworth, 2006; Smith, Flowers,
14
15 & Larkin, 2009), and focus groups (Palmer, Larkin, de Visser, & Fadden, 2010). A key
16
17 requirement of IPA is that the data should be rich and “participants should have been granted an
18
19 opportunity to tell their stories, to speak freely and reflectively, and to develop their ideas and
20
21 express their concerns at some length” (Smith, Flowers, & Larkin, 2009, p.56). Our data fulfilled
22
23 these criteria.
24
25
26
27

28
29 Conducting IPA analysis consists of identifying commonalities and differences about the
30
31 meaning of a given experience. We formatted and line-numbered the data and changed names to
32
33 protect participants’ identities. The first author read and re-read the texts and made some notes
34
35 about the concepts emerging, the text itself, its emotional content, the language used, and so
36
37 forth. From these observations emerged a number of themes, which were classified into
38
39 categories or superordinate themes. Themes were compared across individual accounts, and a
40
41 framework of superordinate themes and subthemes was created, which most accurately
42
43 epitomized women’s experiences. In accordance with IPA requirements, the aim of our analysis
44
45 was to create a “whole” that would go beyond the sum of its parts to provide an interpretation for
46
47 the meaning of the experience (Smith, Flowers, & Larkin, 2009). Throughout the analysis
48
49 process, peer-debriefing was used to enhance transparency and trustworthiness. Randomly
50
51 selected sections of text were also co-coded by the third author to enhance rigor.
52
53
54
55
56
57
58
59
60

1
2
3 Ethical approval was obtained from the University of West London's Psychology
4 Department. Given the sensitive nature of the subject, several ethical considerations had to be
5 addressed. First, we had to consider the potential for distress to be experienced by participants as
6 a result of participating in the study. Accordingly, participants were advised to contact the
7 support group helpline or network of volunteers should they feel the need to speak to someone
8 after completing the survey. However, evidence has suggested that participation in group
9 discussions is beneficial to women who have experienced TFA (Statham, Solomou, & Green,
10 2001). Similarly, self-disclosure of thoughts and feelings has been shown to enhance
11 psychological adjustment (Pennebaker, 2004), and many participants commented that
12 participating in our study constituted an opportunity to share their story and help other women in
13 the future. None of the respondents reported experiencing distress as a result of participating in
14 our study. Thus, the benefits of their participation might have outweighed any potential
15 negatives.

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34 The second ethical consideration related to the first author's membership to the support
35 group's network of volunteers. This had potential implications in terms of confidentiality and
36 duality of roles. To safeguard confidentiality, the data were password-protected and participants'
37 names were changed. With regard to a potential conflict of interest, we genuinely felt that being
38 a group volunteer benefited the study in many ways. Still, in an effort to promote high ethical
39 standards, the first author's details were removed from the list of volunteers sent to women
40 contacting the helpline as a result of participating in the study.

41 42 43 44 45 46 47 48 49 50 **Results**

51
52 We set out to investigate coping strategies at two different points of the TFA process, during the
53 procedure and afterwards, to identify differences and commonalities in coping. These periods are
54 different in nature. Whereas the procedure represents a short-term medical intervention, the
55
56
57
58
59
60

1
2
3 aftermath represents the beginning of a long-term process of grieving. Thus, we wondered
4
5 whether different coping strategies could be used at each point. In our study, we identified more
6
7 commonalities than differences and uncovered the role of key coping strategies that are support,
8
9 acceptance, avoidance, and meaning attribution throughout the experience of TFA.
10
11

12 13 *Coping With the Procedure*

14
15 Five coping strategies emerged as most relevant to women when dealing with the termination
16
17 procedure. These included receiving/giving support, acknowledging the baby, problem solving,
18
19 dissociating oneself from the procedure, and attributing meaning to the birth experience.
20
21

22
23 *Receiving/giving support.* Support was one of the main coping strategies used during
24
25 the procedure. Women unanimously identified their partners as their main source of support.
26
27 Nevertheless, support from health professionals was also considered essential to women's
28
29 experience of the procedure. It involved listening to and addressing women's fears about the
30
31 procedure, as well as providing physical and emotional comfort. Ellie described, "The consultant
32
33 also held my hand tight . . . this warmth from the staff I will always remember." The provision of
34
35 professional support led to better coping, but the absence of it resulted in negative experiences.
36
37 Justine was left on her own for most of the procedure, including the baby's delivery:
38
39

40
41
42 It was so undignified and to see that little body on my own with no support was terrible. I
43
44 called for the midwife who laughed and made a joke about how quick I had been then she
45
46 left the room, allowing me then to deliver the after birth myself, again alone.
47
48

49
50 Support was also derived from the presence, actual or virtual, of other family members.
51
52 Ellie mentioned her mother waiting in the corridor; Wendy had her mother and Rose, her
53
54 mother-in-law, in the delivery room. Christine carried a picture of her three year old son in the
55
56
57
58
59
60

1
2
3 delivery room, which provided focus and strength. Religious support was important to some
4
5 participants. Given the perceived moral dimension of TFA, some women such as Laura
6
7 expressed the fear of being judged: “We went to the Hospital Chapel and met with a Chaplain.
8
9 She was absolutely lovely. I was initially worried that she may judge us, but she didn’t. She was
10
11 sympathetic and very understanding, which helped.”
12
13

14
15 For some women, supporting their partner through the procedure and focusing on their
16
17 partner’s fear and pain helped alleviate their own. For others, supporting their baby through the
18
19 procedure was essential. This was particularly the case when the termination involved a feticide,
20
21 a procedure to induce the baby’s death through intra-cardiac or intra-amniotic injections.
22
23 Recalling the moment the feticide was performed, Rose described how she tried to shield her
24
25 baby from the stress of the procedure:
26
27

28
29 I tried to just breathe through it using deep calm breaths so as not to move too much. I
30
31 didn’t want to cause the baby any distress and I wanted him to die peacefully. It was very
32
33 important that I obeyed the doctor’s instruction to make it as quick and straightforward
34
35 for them.
36
37
38
39
40

41 *Acknowledging the baby.* Acknowledging the baby was seen as a way to accept, and
42
43 thus, cope with the loss. This took many forms. Most participants spent time with their baby after
44
45 delivery, which most found comforting. It provided women with the opportunity to acknowledge
46
47 and bond with their baby. Women were also able to check any resemblance with family
48
49 members, thus recognizing the baby as their own, and see their child’s anomaly for themselves.
50
51 Seeing her baby helped Laura come to terms with her decision: “He did not have any legs and
52
53 this made it easier to know that he definitely did have abnormalities that were apparent and that
54
55 we had made the right decision.” For many women, spending time with their baby was the first
56
57
58
59
60

1
2
3 step in accepting what had happened. For a few, however, it proved upsetting, confronting them
4
5 with the reality of the baby's death. Others felt disturbed at how "weird" their baby looked. A
6
7 handful of participants opted against seeing their baby, mostly because they feared being scarred
8
9 for life.
10

11
12 Acknowledgement also involved conferring an identity on the baby. This included
13
14 naming the baby, taking photos, and showing them to others. The importance of a clear identity
15
16 for the baby is illustrated by Penny's story. Penny was distraught to learn that the gender given to
17
18 her baby at the autopsy differed from the birth report. It left her confused about which baby she
19
20 had lost:
21
22

23
24 We were devastated to realize that we'd had the funeral in the wrong name, the wrong
25
26 name was on the coffin and we'd told our friends and family the wrong name which left
27
28 us having to deal with some very difficult questions. This left us absolutely furious, even
29
30 to the extent that it felt like the initial grieving process had been taken away from us or
31
32 even at times that we'd lost two babies.
33
34
35
36
37

38
39 Underlying this anger was the feeling of having betrayed and denied the baby a proper
40
41 goodbye, another important factor in coping with the loss. Women described leaving the hospital
42
43 without their baby as heart-breaking and the process of saying goodbye as surreal. Parting with
44
45 the baby typically involved three actions: arranging the funeral (which most participants hoped
46
47 would bring some sense of closure), having the baby blessed, and/or placing soft toys or photos
48
49 of themselves in the coffin with the baby. Anna recalled, "I held the toy throughout the labor and
50
51 then left it with him. It was cremated with him. This helped and still brings me comfort to know
52
53 he had a 'friend' to look after him on his journey."
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
Problem solving. In some instances, the procedure was thought of as a practical problem to be solved. This implied preparing for the procedure and focusing on the task ahead. Preparation for the procedure consisted of seeking factual information about the baby's conditions and the termination. This information was often sourced from the Internet and, in some cases, from support group leaflets. Women also relied on experiential knowledge. Isobel had already undergone two TFAs, and thus, felt prepared for the third procedure: "We knew the drill as it were." Others had contact with a group volunteer before the termination, and therefore were aware of what to expect during the procedure. Some of the women who had not been through labor before sought information about the birth process from friends and relatives. Information seeking sometimes engendered a sense of empowerment as exemplified by Laura: "I am the type of person who likes to have prior knowledge about things as it helps me deal with them when they happen; it prepares me." Concentrating on the task of delivering the baby was another way of focusing the mind, and thus coping with the procedure. For Yvonne, "[T]aking one thing at a time, or think[ing] of it in four hour blocks" helped her through the termination.

36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Dissociating oneself from the procedure. Most women described TFA, and, in particular, the procedure to end the pregnancy, as the most difficult experience they ever had to go through. Thus, many tried to distance themselves from what they considered an ordeal. This involved self-distraction, such as watching television, reading magazines and newspaper, or going for walks. Attempts to block the pain were also common, with some participants reporting having an "out of body experience" or going into "autopilot." The use of analgesia during labor was also widespread because it was seen as a way to disconnect with the procedure and avoid unnecessary suffering. Keira used analgesia despite being physically comfortable: "Directly afterwards I was offered some pethidine for the pain, and although I wasn't in pain I accepted it.

1
2
3 It numbed my brain and helped me sleep when I got home.” Underpinning the use of analgesia
4
5 laid expectations and meaning attributed to the birth experience itself.
6
7

8 *Attributing meaning to the birth experience.* Among women who gave birth to their
9
10 baby, the birth experience was an influential factor in the way they coped with the whole
11
12 procedure. The birth often conveyed specific expectations and meanings. In some cases,
13
14 attributing meaning to the birth constituted a resource women could draw on to help them cope
15
16 with their baby’s delivery. Whereas some women tried to shield themselves from the physical
17
18 pain of labor as much as possible, others considered the pain to be constitutive of the birth
19
20 process and embody a particular meaning. Beatrice believed the pain to be “cathartic.” Anna
21
22 regarded the use of analgesia as a failure on her part and wanted to experience the pain to punish
23
24 herself for the decision she had made: “Initially I wanted to feel all the pain of my labor to make
25
26 myself suffer, but then I couldn’t cope so the midwife gave me diamorphine.” For Theresa,
27
28 giving birth without analgesia was her way to normalize the birth and bond with her child. She
29
30 had used very little pain relief for the birth of her previous children and wanted to do the same
31
32 for this baby:
33
34
35
36
37

38 I also wanted to approach my daughter’s labor as similarly as possible to the labors of my
39
40 previous children. I felt like I owed it to her. That at least I could bond with her to that
41
42 extent, give her the same labor as I’d given my other two children . . . For me it was very
43
44 important. There maybe wasn’t a baby to “protect” as such, but there was still a baby to
45
46 bond and connect with for as long as possible. I needed to be with her throughout and
47
48 feel her.
49
50
51
52
53
54

55 Finally, some women mentioned their wish to give birth with dignity, which implied the
56
57 provision of sensitive care. Although this was in most cases provided, there were instances where
58
59
60

1
2
3 it was not, which could have far-reaching consequences on women's psychological adjustment to
4
5 TFA.
6

7 8 *Post Termination Coping*

9
10 Women mostly relied on four coping strategies to deal with the aftermath of the termination:
11
12 remembering the baby, receiving/providing emotional support, avoidance, and looking to the
13
14 future.
15

16
17 *Remembering the baby.* Remembering the baby was a key element in post-TFA
18
19 coping. This involved talking about the baby to partners, friends, and family, as well as sharing
20
21 stories with other group members, which most found soothing. Having a remembrance service
22
23 also enabled women to acknowledge the baby and provided some closure. A few women chose
24
25 to read poems at the service, others played meaningful music; however, for some, the service
26
27 was a painful experience, emphasizing the unnatural order of the loss in which a parent loses a
28
29 child. Holding a service was a way to give the baby social recognition. This was particularly
30
31 important given that many women felt that their loss was not fully sanctioned by society. Many,
32
33 like Justine, reported feeling isolated and misunderstood in their grief: "Nobody can understand
34
35 the torture unless they have experienced it themselves as I don't think it can be likened to
36
37 anything else."
38
39
40
41
42

43
44 The use of rituals and token acts was widespread. Visiting a place of remembrance was
45
46 common and most women kept a memory box. In some cases, the memory box was provided by
47
48 the hospital, but more often it was sourced by the parents. Rose recalled her partner building the
49
50 box over a period of time, almost mirroring his mourning process: "My partner built a little box
51
52 to put little mementos into. It took him many months to complete it as I think it was his time to
53
54 think about the son he had lost." The memory box usually contained copies of the scans of the
55
56 baby, photos of the baby, letters of condolence, poems, baby's hand and foot prints, toys, clothes,
57
58
59
60

1
2
3 and other symbolic objects. Women reported initially looking at the box on a daily basis, but this
4
5 need tended to lessen in time. Gemma recalled the location of the box being a source of conflict
6
7
8 with her partner, a possible reflection of grieving differences between them:
9

10 More recently he has been happier to talk about it; though he would prefer that her
11
12 memory box was tucked away (it is on top of the unit in the living room and not
13
14 obvious). But for a long time, he got cross.
15
16
17

18
19 Some women carried a photo album in their bag. Others displayed the baby's photos next to their
20
21 other children's. Other token acts included getting a tattoo, or planting a tree, a permanent and
22
23 ever-growing feature in the women's lives.
24
25

26
27 *Receiving/providing emotional support.* Support after the termination was essential to
28
29 the way women coped. This support took many forms. Support from health professionals was
30
31 particularly important but in many cases, this support was limited post termination. Often, it was
32
33 the interaction women had with health professionals during the procedure, which had a lasting
34
35 influence on the way they coped with the event. Instances where clinical care went beyond the
36
37 traditional doctor-patient relationship were received with much gratitude and had a soothing
38
39 effect long after the termination. As Bonnie recalled,
40
41

42
43 I think of his words. He very kindly and reassuringly said that he would make the same
44
45 decision in our position—something that he confessed he shouldn't say but something we
46
47 found so helpful since he was the person who had the skill to "fix the problem."
48
49
50

51
52 Support from health professionals post termination extended to other forms of care and
53
54 communication. Rose's district nurse had not read the paperwork prior to visiting her. Rose was
55
56 particularly upset when the nurse came to obtain a blood sample from the baby: "I had to tell her
57
58
59
60

1
2
3 the baby had died. She hadn't read the notes properly! I was furious and very distressed." The
4
5 language used to refer to the baby, particularly in postmortem reports, also influenced women's
6
7 experience, as illustrated by Olivia:
8
9

10 All of her paperwork stated she was a "nonviable fetus." I felt she deserved to be
11
12 recognized . . . because to us, she very much existed. P.S. - something that really doesn't
13
14 help, is paperwork saying "non-viable fetus." Very upsetting to read.
15
16
17

18
19 Professional support was also dispensed in the form of counseling, which was mainly used by
20
21 women more advanced in the recovery process. Even though the majority found counseling
22
23 helpful, some felt that they had started the process too early and felt vulnerable as a result. The
24
25 timing of the intervention might therefore be an important factor.
26
27

28
29 Emotional support was also sourced from friends and relatives, but its quality varied
30
31 greatly. Support was key in helping women cope with their loss, but most women indicated that
32
33 it faded over time. A lack of support, particularly in the early stage, engendered much suffering
34
35 as exemplified by Penny:
36
37

38 I was horrified to spend a weekend with some of [my] closest friends, the week of the
39
40 funeral and not once did they ask about the funeral or how we were . . . remember feeling
41
42 like a leper after that weekend and that people avoided talking to me.
43
44
45

46
47 In rare instances, women chose to withdraw from their social network, particularly when friends
48
49 went on to have successful pregnancies.
50
51

52 Support groups were another source of emotional support, which is unsurprising given
53
54 that the sample was drawn from a support group. Support groups form a distinctive category in
55
56 that they involve a reciprocal support relationship based on a common experience. Most women
57
58
59
60

1
2
3 considered the group e-mail network/forum to be one of the most helpful elements in the coping
4
5 process. Being able to reciprocate, getting support as well as offering support, was of great
6
7 benefit to the women. Gemma recalled,
8
9

10 [It] has been the most amazing thing though as everyone really understands more than
11
12 anyone else ever could. I still read the posts every day and have done some fund-raising
13
14 for them. I would never, ever have done so well without their support.
15
16
17

18
19
20 Reasons for using the group e-mail network/forum varied, but the underlying drive was to derive
21
22 a sense of solidarity. For some women, like Christine, the forum became the only opportunity to
23
24 share their story: “I feel completely alone in my grief as no one seems to understand just how
25
26 profound it is. This includes my partner.” However, other women reported having ambivalent
27
28 feelings about using the forum. Ulrika saw the forum as a lifeline during dark days, but actively
29
30 avoided it on good days to prevent being “brought back to the darkness.” Bonnie also had mixed
31
32 feelings about the e-mail network because some of the comments led her to believe that her way
33
34 of grieving was inadequate:
35
36
37

38 Sometimes I found reading other people’s accounts on the forum unhelpful as I felt guilty
39
40 for not feeling as emotional or terrible as they did, but in time I was able to feel that this
41
42 was positive, that I was coping and mentally strong.
43
44
45
46
47

48 Going to parents’ meetings was another coping resource, and it provided women with a “sense
49
50 check” regarding where they were in the healing process. A few participants also used a support
51
52 group specialized in stillbirth and neonatal deaths. Some found it helpful, but others reported
53
54 feeling uncomfortable given that this particular group only dealt with naturally occurring or
55
56 accidental deaths rather than terminations.
57
58
59
60

1
2
3 *Avoidance.* Avoidance constituted another coping strategy, although a less commonly
4 used one. Avoidance usually involved self-distraction, such as going away after the termination
5 or on the baby's due date, decorating, or moving house. Keeping busy was seen as an antidote to
6 feeling the pain as illustrated by Holly:
7
8
9

10
11 When it was most painful I kept myself busy so that I [made] an effort to put it to the
12 back of my mind and made sure I was tired each evening so that I could get some sleep at
13 night.
14
15
16
17
18
19

20
21 Similarly, some women were eager to return to work soon after the termination. Although
22 women reported that this initially helped, they also recognized that this strategy might have
23 pushed emotions aside and hampered the grieving process. Keira recalled, "Keeping busy and
24 going back to work might have helped initially, but I think they stored up the emotions for later."
25 For some participants like Megan, concentrating on other children helped alleviate the pain, by
26 channeling their energy and deriving meaning: "I kept myself busy with my other kids. I went
27 into overdrive I think; almost like I had to prove my worth as a mum." Avoidance also meant
28 actively trying to block the pain, which only a few participants mentioned. This implied avoiding
29 thinking about the baby and trying to rationalize the loss by framing it in a practical rather than
30 an emotional way. This strategy was used by women like Natalie for whom this was not the first
31 TFA: "I have never allowed myself to think as if I was having a baby. I've always thought it too
32 good to be true so I have just not imagined a baby at the end. I think this has really helped." A
33 handful of participants also recalled drinking heavily initially in an attempt to block the pain.
34 Avoidance also extended to avoiding pregnant women or women with young babies, which a
35 few participants admitted to.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 *Looking to the future.* Most women reported that looking to the future helped them
4
5 cope with their loss. One of the first steps in this process was to regain a sense of normality. This
6
7 was achieved by getting back into a routine, going back to work, and trying to function as
8
9 normally as possible. As Beatrice recalled, “I went back to work four weeks after the
10
11 termination. I thought getting back into a routine would help the grieving process.” Looking to
12
13 the future was also linked to getting closure. All women reported the process of healing as
14
15 uneven, similar to a roller coaster, and lengthy. This process was usually punctuated by
16
17 important milestones such as the funeral, postmortem results, due date, and the first birthday.
18
19 Ellie described, “As closure was
20
21 provided through the funeral, postmortem, due date passing, the first birthday, now it is time to
22
23 move on with our lives.”
24
25
26
27
28

29 The baby’s funeral was usually the first milestone in the process, and generally provided
30
31 some closure with the physical side of the loss: the loss of the “actual” baby. The postmortem
32
33 constituted a second milestone for those who requested it. Participants found the postmortem
34
35 results helpful in reaching some closure on the decision they had made, as illustrated by Frances:
36
37 “The postmortem report also helped as it confirmed the diagnosis and put closure to that
38
39 irrational fear that we might have made the wrong decision.” The baby’s due date was perhaps
40
41 the most important milestone. It was often an anxious time because it directly confronted women
42
43 with the reality of the loss, as described by Ulrika: “I will be mourning the loss of Oscar all over
44
45 again.” Many reported being unable to “let go” before the due date, and it appeared to provide
46
47 some closure on the pregnancy, the “virtual” baby, and the hopes the pregnancy had brought.
48
49 The baby’s first birthday was the fourth significant milestone. It signaled the beginning of a new
50
51 type of mourning process, which up to that point consisted of getting from one milestone to the
52
53 other, each milestone bringing closure to some aspects of the loss. After the first birthday, the
54
55
56
57
58
59
60

1
2
3 focus moved from grieving for the lost pregnancy, dream, and baby to a long-term grieving for
4 the child. For Gemma, birthdays were important and a way to honor her daughter's memory: "I
5 celebrate Louise's birthdays with cake, lantern, a card, and I often receive cards from lovely
6 people who remember." Focusing on another pregnancy was another widely used coping
7 strategy. It involved closing on the TFA and projecting oneself in the future. This focus was, in
8 some instances, all-consuming and often bittersweet. A new pregnancy was usually a source of
9 worry and failure to get pregnant often led to feelings of despair, as illustrated by Keira:

10
11
12
13
14
15
16
17
18
19
20 The only time when I felt like I was unable to cope again was when it came to be nearly a
21 year and half after the termination and I was still not pregnant. I felt that only having
22 another baby would heal the wound which I still have, and a seeming inability to
23 conceive again was almost like a punishment.
24
25
26
27
28
29
30
31
32

33 Women who became pregnant were eager to stress that their new baby was not a replacement for
34 the baby they had lost.
35
36
37

38 Finally, looking for positives in their experiences also contributed to women's coping.
39 Although women described the decision to end their pregnancy as the hardest they ever made,
40 most were at peace with their decision. Some drew comfort from the fact that their baby's
41 condition was detected early enough to enable them to make a decision. For some, terminating
42 the pregnancy was the only parental decision they could make, as Laura recalled:
43
44
45
46
47
48
49

50 I understood that as a mother, there are times when you have to make decisions for your
51 children, on their behalf if it relates to safety, etc. Although this decision was a bit
52 extreme, I felt that as a mother, I was making a decision to take away my child's pain and
53 it was for reasons out of love.
54
55
56
57
58
59
60

1
2
3
4
5
6 Looking for positives also involved putting the experience to good use. Some participants
7
8 sponsored a child, some ran marathons to raise money for charities specialized in their baby's
9
10 condition, while some found solace in helping other women. For some participants, the
11
12 termination also instigated personal positive growth, with some reporting feeling stronger and
13
14 more confident as a result. For Theresa, the termination triggered a sense of urgency to live life
15
16 to the full, fulfill her potential, and follow her dreams:
17

18
19 This feeling of doing things in her memory and of using her memory to instigate positive
20
21 things has also been hugely important in terms of ways of coping . . . I've been filled with
22
23 an urge to do things that I hadn't been confident to do otherwise beforehand. It has made
24
25 me realize the fragility of life, and it has made me want to treasure each day more, risk
26
27 things more in order to live the life I really want to live.
28
29
30
31
32
33

34 For others, the termination triggered a desire to address unresolved issues or explore new career
35
36 paths, thus having a positive impact.
37

38 **Discussion**

39
40 To our knowledge, this study is the first to focus exclusively on women's coping strategies used
41
42 throughout the process of TFA, both during and after the procedure. Thus, our study provides a
43
44 valuable insight into the lived experiences of women who are coping with this traumatic life
45
46 event. We hope that this insight will inform clinical practice and be valuable to women who
47
48 undergo this procedure. In this article, we suggest that coping with the procedure of TFA
49
50 involved five processes: seeking and providing support, acknowledging the baby, problem
51
52 solving, dissociating oneself from the procedure, and attributing meaning to the birth experience.
53
54
55
56
57
58
59
60

1
2
3 Regarding post termination coping, we identified four processes: seeking and providing
4 emotional support, remembering the baby, avoidance, and looking to the future.
5
6

7
8 Throughout the TFA process, four major structures of coping emerged: support,
9 acceptance, avoidance, and meaning attribution. The consistency of these structures across time
10 indicates the importance of dispositional characteristics. Problem solving was specific to coping
11 with the procedure, whereas elements of looking to the future (e.g., returning to normality and
12 focusing on a new pregnancy) only related to post termination coping. Our results are in line
13 with some of the current literature on coping. For example, in their review of 100 coping
14 assessments, Skinner and colleagues (2003) suggest that there are five core categories of coping:
15 problem solving, support seeking, avoidance, distraction, and positive cognitive restructuring.
16
17

18
19 Support was the first essential coping structure present throughout the TFA process. The
20 concept of support involved two separate activities: receiving and providing it. Receiving
21 support was crucial to women. Research has shown that receiving social support contributes to
22 general well-being (Lakey & Orehek, 2011). It is particularly important in the context of TFA
23 because a lack of support has been associated with mood disturbances in women (Green &
24 Statham, 2007). Similar to research studies about perinatal loss (Lasker & Toedter, 1991) and
25 TFA (Statham et al., 2001), partner's support was critical to the women in our study. Many
26 participants felt that their loss was not fully recognized by society and, in numerous cases,
27 women only fully shared their experience with their partners.
28
29

30
31 Support from health professionals was also critical to women's coping. This included
32 emotional support from health professionals, as documented elsewhere (Geerinck-Vercammen &
33 Kanhai, 2003), and care provision both during and after the procedure. Most participants in our
34 article found the care in hospital satisfactory, which contributed to alleviating their distress.
35
36 However, the aftercare was often seen as deficient. Many women reported feeling unsupported
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 post termination, a finding in line with most TFA literature (Green & Statham, 2007; Statham et
4
5 al., 2001).

6
7
8 Many women used various support groups as a source of comfort, which is unsurprising
9
10 given that all participants were to some degree active on the forum or e-mail network of the
11
12 group we recruited from. Nevertheless, the extensive use of support groups also underlined the
13
14 importance of the reciprocal nature of support. The positive impact on mental well-being of
15
16 being in a position to provide support has been reported in studies across various fields of
17
18 psychology (Knoll, Kienle, Bauer, Pfuller, & Luszczynska, 2007; Warner, Schuz, Wurm,
19
20 Ziegelman, & Tesch-Romer, 2010). Comments left by participants at the end of our survey
21
22 corroborate this view. Thus, supporting others seemed to enable women to transcend their pain,
23
24 put it to good use, and frame it in a positive way.
25
26
27

28
29 Acceptance was the second key coping structure common to both periods. It involved
30
31 acknowledging and remembering the baby, which was attained in different ways and
32
33 predominantly experienced as helpful. Trying not to think of the baby, blocking the pain, or
34
35 busying oneself often resulted in short-term gain but long-term loss. Women engaged in rituals
36
37 and token acts. Women who opted to see their baby after delivery found this comforting, but a
38
39 minority found it disturbing.
40
41
42

43
44 Whether or not to spend time with the baby in the context of perinatal loss has generated
45
46 clinical discussion. For example, until recently in the United Kingdom, parents were routinely
47
48 encouraged to spend time with their baby on the assumption that doing so facilitated the grieving
49
50 process (Bennett, Litz, Lee, & Maguen, 2005; Geerinck-Vercammen & Kanhai, 2003). However,
51
52 a study by Hughes, Turton, Hopper, and Evans (2002) on psychological adjustment to stillbirth
53
54 found that women who choose to see their baby adjust to their loss less easily than those who
55
56 choose not to. Similarly, Green and Statham (2007) suggested that women who do not seek care
57
58
59
60

1
2
3 or engage in any form of ritual following TFA fare better psychologically than those who do. In
4
5 light of this contrasted evidence, the UK National Institute for Health and Clinical Excellence
6
7 (2007) issued guidelines about care in perinatal loss recommending not to routinely encourage
8
9 mothers to spend time with their baby, but instead to leave the decision to parents. By
10
11 highlighting the range of emotions associated with spending time with the baby, our findings
12
13 support this clinical recommendation.
14
15
16

17
18 The third coping structure common to the procedure and its aftermath was avoidance. It
19
20 included themes of self-distraction, dissociation, and avoidance, with the latter more prominent
21
22 in post-termination coping. It was the least homogeneous structure, with some elements
23
24 experienced as helpful and others unhelpful. Traditionally, self-distraction and dissociation have
25
26 been linked to avoidant coping styles. Thus, they are considered maladaptive strategies,
27
28 primarily because they are difficult to sustain (Folkman & Lazarus, 1988; Schnider et al., 2007).
29
30 Nevertheless, in our study, both elements served an adaptive function, particularly during the
31
32 procedure, by protecting women from further suffering. In contrast, most coping components of
33
34 the avoidance theme in post termination coping involved various degrees of active denial and
35
36 disengagement (actively trying not to think about the baby, drinking, getting busy to the point of
37
38 exhaustion), which could be considered as maladaptive. These strategies are of limited use in the
39
40 long term because they do not address the event and its effect in an individual's life (Carver &
41
42 Connor-Smith, 2010).
43
44
45
46
47

48
49 The fourth common coping structure was meaning attribution. This included attributing
50
51 meaning to the birth, deriving meaning, and experiencing growth as a result of TFA. This coping
52
53 structure is echoed in the concept of resilience. Research in that field has shown that people can
54
55 experience stress-related growth following a traumatic event (Bonanno, 2008). Research studies
56
57 have established that the action of attributing meaning might lead to positive changes in people's
58
59
60

1
2
3 lives (Helgeson, Reynolds, & Tomich, 2006). Therefore, stress-related growth can be seen as a
4
5 central part of the coping process rather than as a positive, unexpected outcome of a difficult
6
7 experience. Our results show that, as a result of TFA, a number of women have discovered
8
9 strengths they were unaware of having. Others have unearthed a drive to take control of their
10
11 lives and fulfill their potentials.
12
13

14
15 Problem solving was a coping strategy used solely during the procedure. The benefit of
16
17 preoperative information on levels of anxiety prior to surgery (McDonald, Hetrick, & Green,
18
19 2004) and on recovery has been documented (Shulldham, 1999). Focusing on the task rather than
20
21 thinking about its meaning was also helpful. Both strategies were used as buffers against the
22
23 stress and pain of the procedure. The strategy of looking to the future, not surprisingly,
24
25 predominantly belonged to post-termination coping, although meaning attribution was common
26
27 during both periods. It involved a return to normality and focusing on another pregnancy, and
28
29 thus, reflected a drive to restore equilibrium and move forward.
30
31
32
33

34
35 The focus on a new pregnancy engendered mixed feelings. Many women experienced
36
37 anxiety during their new pregnancy, a finding echoed in the literature (Rillstone & Hutchinson,
38
39 2001). This also raises a question about the recommended time between the termination and a
40
41 subsequent pregnancy. A study by Hughes, Turton, and Evans (1999) in psychological
42
43 adjustment to stillbirth suggests that women should preferably wait 12 months following their
44
45 loss to conceive again. This has implications not only for the mother but also for the infant
46
47 because anxiety and distress during pregnancy can negatively affect the well-being of the baby
48
49 (Brouwers, van Baar, & Pop, 2001). To our knowledge such information applied in the context
50
51 of TFA is lacking, and hence warrants further research.
52
53
54

55
56 Our research shows that women used coping strategies that are largely considered to be
57
58 adaptive during and after the termination. According to the coping literature (Carver, 1997;
59
60

1
2
3 Carver & Connor-Smith, 2010; Folkman & Lazarus, 1988) seeking support, whether
4
5 instrumental (e.g., aspects of care being provided by health professional) or emotional (e.g.,
6
7 derived from partner and/or friends) is an adaptive coping strategy. Similarly,
8
9 acknowledging/remembering the baby, which is linked to the concept of acceptance, and
10
11 meaning attribution, which is related to positive reframing and growth, are also considered
12
13 adaptive. When traditionally labeled maladaptive strategies were used (self-distraction,
14
15 dissociation, and avoidance), they either served an adaptive purpose of short-term protection
16
17 and/or were used by only a few participants.
18
19
20
21

22 The debate about which coping strategy is adaptive and which is maladaptive is ongoing.
23
24 Whether a strategy is adaptive or not is subjective. Carver (1997) himself advised against
25
26 classifying coping strategies into discrete categories, such as problem-focused or emotional
27
28 active/avoidant coping. His advice instead was to examine each aspect of coping independently.
29
30 Coping with a traumatic event is a personal process, determined by individual (e.g., personality,
31
32 personal history, attitudes, and beliefs) and situational factors (e.g., type of stressor, timing of
33
34 event). Consequently, what is adaptive for some might not be so for others.
35
36
37
38

39 Our findings have several practical implications, the main one being care provision. In
40
41 view of the importance of support, it is key for health professionals to understand the nature of
42
43 the loss to provide suitable care. Although different in nature, the grieving process in TFA has
44
45 been compared to grieving after a stillbirth (Salvesen et al., 1997; Statham et al., 2001). If the
46
47 unexpected nature of a stillbirth affects the grieving process, other elements might complicate the
48
49 grieving process for a baby one has chosen not to have. White-Van Mourik and colleagues
50
51 (1992) argued that most negative feelings inherent to TFA can be articulated around the loss of
52
53 self-esteem: loss of biological self-esteem because of the pregnancy ending in a failure; loss of
54
55 moral self-esteem caused by feelings of responsibility and sometimes guilt that accompany the
56
57
58
59
60

1
2
3 decision to end the pregnancy; finally, loss of social self-esteem derived by a perceived inability
4 to bring a disabled child into the world and subsequent mixed feelings about what is “gained”
5 from the loss. These issues alongside elements of guilt and self-blame, which have also surfaced
6 in the data, must be taken into account. Research has also shown that women who undergo TFA
7 often feel alienated and stigmatized in their loss (Bryar, 1997; McCoyd, 2010; Rillstone &
8 Hutchinson, 2001), predominantly because TFA is still conceptualized within the abortion
9 paradigm (McCoyd, 2010). In addition, because providing support to others, preparation, and
10 experiential knowledge were experienced as helpful, early referral to support groups might be
11 beneficial.
12
13
14
15
16
17
18
19
20
21
22
23

24 Our findings also point to the potential benefit of promoting coping strategies to women
25 who undergo TFA, such as acceptance, positive reframing, and benefit finding. It might also be
26 useful to emphasize psychological concepts such as self-esteem and resilience. Although
27 controversial, the potential value of seeing the baby and creating memories might also be
28 emphasized alongside suggestions of ways to achieve this. Conversely, women who express a
29 desire to distance themselves from the event should be allowed to do so. Care needs to be open,
30 sensitive, nondirective, and ultimately tailored to each woman’s particular needs. Our study also
31 revealed some inadequacies in aftercare, which might result in poor psychological adjustment.
32 These could have significant consequences on women’s quality of life and their well-being in
33 subsequent pregnancies. Previous prenatal loss affects the mental health of women who conceive
34 again, with typically higher levels of anxiety and depression being observed among those who
35 have experienced a loss compared to those who have not (Blackmore et al., 2011). Similarly,
36 women who conceive after TFA have reported a new pregnancy to trigger a “re-emergence of
37 anguish” (Rillstone & Hutchinson, 2001). Consequently, it might be beneficial to monitor
38 women who have undergone TFA during subsequent pregnancies.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 One the many strengths of this study is the novel and prolific nature of the feedback
4 gathered. The study also provided a welcomed opportunity for women to share their experiences,
5
6 with many who indicated that their participation in the study had a therapeutic effect. By making
7
8 a summary of the results available to all participants, we also endeavored to disseminate the
9
10 results directly to the group most in need of this information and of the changes this information
11
12 may bring. Our study was purposively exploratory, and thus, has limitations that warrant further
13
14 research. First, the sample was limited to women who were, to various degrees, active members
15
16 of a support group. Research has shown that the use of online groups can lead to personal
17
18 empowerment, through the provision of a forum for self-expression and social support. This
19
20 empowerment might act as a buffer against distress (Barak, Boniel-Nissim, & Suler, 2008) and
21
22 influence coping processes. Consequently, further research based on women who are not using a
23
24 support group is indicated.
25
26
27
28
29
30

31 In addition, as with many studies on TFA conducted in high-income countries, the
32
33 sample in our study predominantly comprised White, well-educated participants. It is possible
34
35 that women who are less articulate might find it more difficult to voice their feelings, which
36
37 might influence the way they cope. Thus, widening the demographic profile of participants
38
39 would be beneficial. This observation also raises the question of health literacy, given that TFA
40
41 is a situation that involves a large exchange of complex information and often requires the input
42
43 of many different health professionals (e.g., geneticists, neurologists, obstetricians).
44
45
46
47

48 Second, it is possible that some accounts reflect a social desirability bias (van de Mortel,
49
50 2008). Thus, some women might have under-reported the use of more maladaptive coping
51
52 strategies. Similarly, some women might have felt under pressure to report their experience in a
53
54 way they believed fit with society's stereotypes of motherhood. Comments from a few
55
56 participants suggest that some women might feel inadequate if they are not overwhelmed by
57
58
59
60

1
2
3 sadness and pain. Hochschild (1983) suggested that society is governed by “feeling rules,” which
4
5 are sets of beliefs about how people think they ought to feel. These rules could also apply to TFA
6
7 (McCoyd, 2009) and might influence the way coping processes are reported. Third, the
8
9 retrospective nature of the study raises the possibility of a degree of post hoc rationalization. This
10
11 is particularly relevant in the context of TFA because of the ethical issues it raises. Women
12
13 might have post-rationalized their coping processes in light of the decision they made. This is
14
15 suggested by some research studies in the field of abortion, antenatal screening, and prenatal sex
16
17 identification, which have emphasized the concept of cognitive dissonance (Burke, 1992;
18
19 Sandelowski & Borroso, 2005). Cognitive dissonance occurs when there is an uncomfortable
20
21 conflict between ideas or feelings that are held simultaneously (Festinger, Riecken, & Schachter,
22
23 1956). The underlying assumption is that people are naturally driven to reduce this dissonance as
24
25 much as possible by either changing their attitudes or rationalizing and justifying their actions.
26
27
28
29
30

31
32 The concept of cognitive dissonance is pertinent to TFA (Statham, H. personal
33
34 communication, July 4, 2011) given that the pregnancy is, in most cases, wanted, and thus, the
35
36 decision to end it is particularly complex (McCoyd, 2008). In some instances the decision
37
38 strongly conflicts with personal religious beliefs, which accentuates the dissonance. It is
39
40 therefore possible that when women recall their termination, they try, to some extent, to justify
41
42 their decision by rationalizing it, and in turn, rationalizing the coping strategies involved. Ideally,
43
44 further research would gather prospective data, from the time a fetal abnormality is detected to
45
46 well after the termination. This study contributes to the knowledge in the area of pregnancy
47
48 termination for fetal abnormality. In doing so, it has the potential to inform clinical practice, as
49
50 well as to benefit women who are facing TFA. It also clearly represents a starting point rather
51
52 than a conclusion, and thus might initiate exploration into new areas of research.
53
54
55
56

57 **Authors' Note**

1
2
3 Portions of this article were presented at the British Psychological Society *Division of Health*
4
5 *Psychology Conference*, September 15, 2011, in Southampton, UK.
6
7

8 **Acknowledgements**

9
10 We thank Antenatal Results and Choices (ARC) for their contribution to the study.
11

12 **Declaration of Conflicting Interests**

13
14 The authors declared the following potential conflict of interest with respect to the research,
15
16 authorship, and/or publication of this article: Caroline Lafarge is a member of the volunteers'
17
18 network of Antenatal Results and Choices (ARC). The other authors (Kathryn Mitchell and
19
20 Pauline Fox) have no interest to declare.
21
22
23

24 **Funding**

25
26 The authors received no financial support for the research, authorship, and/or publication of this
27
28 article.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- Barak, A., Boniel-Nissim, M., & Suler, J. (2008). Fostering empowerment in online support groups. *Computers in Human Behavior, 24*, 1867–1883. doi:10.1016/j.chb.2008.02.004
- Bennett, S. M., Litz, B. T., Lee, B. S., & Maguen, S. (2005). The scope and impact of perinatal loss: Current status and future directions. *Professional Psychology: Research and Practice, 36*(2), 180–187. doi:10.1037/0735-7028.36.2.180
- Biggerstaff, D. L., & Thompson, A. R. (2008). Interpretative Phenomenological Analysis (IPA): A qualitative methodology of choice in healthcare research. *Qualitative Research in Psychology, 5*(3), 214–224. doi:10.1080/14780880802314304
- Blackmore, R. E., Côté-Arsenault, D., Tang, W., Glover, V., Evans, J., Golding, J., & O'Connor, T. G. (2011). Previous prenatal loss as a predictor of perinatal depression and anxiety. *British Journal of Psychiatry, 198*, 373–378. doi: 10.1192/bjp.bp.110.083105
- Bonanno, G. A. (2008). Loss, trauma, and human resilience: Have we underestimated the human capacity to thrive after extremely aversive events? *Psychological Trauma: Theory, Research, Practice and Policy, 5*(1), 101–113. doi: 10.1037/1942-9681.5.1.101
- Brouwers, E. P. M., van Baar, A. L., & Pop, V. J. M. (2001). Maternal anxiety during pregnancy and subsequent infant development. *Infant Behavior and Development, 24*(1), 95–106. Retrieved from [http://dx.doi.org/10.1016/S0163-6383\(01\)00062-5](http://dx.doi.org/10.1016/S0163-6383(01)00062-5)
- Bryar, S. H. (1997). One day you're pregnant and one day you're not: Pregnancy interruption for fetal anomalies. *Journal of Obstetric, Gynecologic & Neonatal Nursing, 26*(5), 559–566. doi: 10.1111/j.1552-6909.1997.tb02159.x
- Burke, M. B. (1992). Genetic counsellor attitudes towards fetal sex identification and selective abortion. *Social Science & Medicine, 34*(11), 1263–1269.

- 1
2
3 Cartwright, T., Endean, N., & Porter, A. (2009). Illness perceptions, coping and quality of life in
4 patients with alopecia. *British Journal of Dermatology*, *160*(5), 1034–1039. doi:
5
6 10.1111/j.1365-2133.2008.09014.x
7
8
9
10
11 Carver, C. S. (1997). You want to measure coping but your protocol's too long: Consider the
12 Brief COPE. *International Journal of Behavioral Medicine*, *4*(1), 92–100. doi:
13
14 10.1207/s15327558ijbm0401_6
15
16
17
18 Carver, C. S., & Connor-Smith, J. (2010). Personality and coping. *Annual Review of Psychology*,
19 *61*, 679–704. doi: 10.1146/annurev.psych.093008.100352
20
21
22
23 Coleman, P. K. (2011). Abortion and mental health: Quantitative synthesis and analysis of
24 research published 1995–2009. *British Journal of Psychiatry*, *199*, 180–186. doi:
25
26 10.1192/bjp.bp.110.077230
27
28
29
30 Davies, V., Gledhill, J., McFadyen, A., Whitlow, B., & Economides, D. (2005). Psychological
31 outcome in women undergoing termination of pregnancy for ultrasound-detected fetal
32 anomaly in the first and second trimesters: A pilot study. *Ultrasound in Obstetrics and*
33 *Gynecology*, *25*(4), 389–392. doi: 10.1002/uog.1854
34
35
36
37
38
39
40 Department of Health. (2011). *Abortions Statistics, England and Wales in 2010*. Retrieved from
41 [http://www.dh.gov.uk/prod_consum_dh/groups/dh_digitalassets/documents/
42 digitalasset/dh_127202.pdf](http://www.dh.gov.uk/prod_consum_dh/groups/dh_digitalassets/documents/digitalasset/dh_127202.pdf).
43
44
45
46
47 Festinger, L., Riecken, H. W., & Schachter, S. (1956). *When prophecy fails*. Minneapolis:
48 University of Minnesota Press.
49
50
51
52 Folkman, S., & Lazarus, R. S. (1988). Coping as a mediator of emotion. *Journal of Personality
53 and Social Psychology*, *54*(3), 466–475. doi: 10.1037/0022-3514.54.3.466
54
55
56
57
58
59
60

- 1
2
3 Green, J. M., & Statham, H. E. (2007). Psychosocial aspects of prenatal diagnosis: the challenges
4
5 for doctors and patients. In J. Cockburn & M. Pawson (Eds.), *Psychological challenges*
6
7 *in obstetrics and gynecology: The clinical management* (pp. 107–120). London:
8
9 Springer-Verlag.
10
11
12 Geerinck-Vercammen, C. R., & Kanhai, H. H. H. (2003). Coping with termination of pregnancy
13
14 for fetal abnormality in a supportive environment. *Prenatal Diagnosis*, 23(7), 543–548.
15
16 doi: 10.1002/pd.636
17
18
19 Heartbeat bill. (n.d.). *The Heartbeat bill*. Retrieved from <http://heartbeatbill.com>
20
21
22 Helgeson, V. S., Reynolds, K. A., & Tomich, P. L. (2006). A meta-analytic review of benefit
23
24 finding and growth. *Journal of Consulting and Clinical Psychology*, 74(5), 797–816. doi:
25
26 10.1037/0022-006X.74.5.797
27
28
29 Hochschild, A. R. (1983). *The managed heart: Commercialization of human feelings*. Berkeley
30
31 and Los Angeles: University of California Press.
32
33
34 House of Commons. (2007a). *Science and Technology Committee Publications. Twelfth report*.
35
36 Retrieved from
37
38 [http://www.publications.parliament.uk/pa/cm200607/cmselect/cmsctech/1045/104506.ht](http://www.publications.parliament.uk/pa/cm200607/cmselect/cmsctech/1045/104506.htm)
39
40 [m](http://www.publications.parliament.uk/pa/cm200607/cmselect/cmsctech/1045/104506.htm)
41
42
43 House of Commons. (2007b). *Science and Technology. Written Evidence*. Retrieved from
44
45 [http://www.publications.parliament.uk/pa/cm200607/cmselect/cmsctech/1045/1045we30.](http://www.publications.parliament.uk/pa/cm200607/cmselect/cmsctech/1045/1045we30.htm)
46
47 [htm](http://www.publications.parliament.uk/pa/cm200607/cmselect/cmsctech/1045/1045we30.htm)
48
49
50 Hughes, P. M., Turton, P., & Evans, C. D. H. (1999). Stillbirth as risk factor for depression and
51
52 anxiety in the subsequent pregnancy: Cohort study. *British Medical Journal*, 318, 1721–
53
54 1724. doi: 10.1136/bmj.318.7200.1721
55
56
57
58
59
60

- 1
2
3 Hughes, P. M., Turton, P., Hopper, E., & Evans, C. D. H. (2002). Assessment of guidelines for
4 good practice in psychosocial care of mothers after stillbirth: A cohort study. *Lancet*,
5 360(9327), 114–118. Retrieved from [http://dx.doi.org/10.1016/S0140-6736\(02\)09410-2](http://dx.doi.org/10.1016/S0140-6736(02)09410-2)
6
7
8
9
10 Keefe-Cooperman, K. A. (2004-2005). A comparison of grief as related to miscarriage and
11 termination for fetal abnormality. *OMEGA-Journal of Death and Dying*, 50(4), 281–300.
12 Retrieved from
13
14 <http://baywood.metapress.com/openurl.asp?genre=article&id=doi:10.2190/QFDW->
15 LGEY-CYLM-N4LW
16
17
18
19
20
21
22 Kersting, A., Dorsch, M., Kreulich, C., Reutemann, M., Ohrmann, P., Baez, E., & Arolt, V.
23 (2005). Trauma and grief 2–7 years after termination of pregnancy because of fetal
24 anomalies—a pilot study. *Journal of Psychosomatic Obstetrics and Gynecology*, 26(1), 9–
25 14. doi:10.1080/01443610400022967
26
27
28
29
30
31
32 Knoll, N., Kienle, R., Bauer, K., Pfüller, B., & Luszczynska, A. (2007). Affect and enacted
33 support in couples undergoing in-vitro fertilization: When providing is better than
34 receiving. *Social Science & Medicine*, 64(9), 1789–1801. Retrieved from
35
36 <http://dx.doi.org/10.1016/j.socscimed.2007.01.004>
37
38
39
40
41 Korenromp, M. J., Page-Christiaens, G. C. M. L., van den Bout, J., Mulder, E. J. H., Hunfeld, J.
42 A. M., Potters, C. M. A. A., . . . Visser, G. H. A. (2007). Parental coping four months
43 after termination of pregnancy for fetal anomalies: A prospective study in 226 cases.
44
45 *Prenatal Diagnosis*, 27(8), 709–716. doi: 10.1002/pd.1763
46
47
48
49
50
51 Lakey, B., & Orehek, E. (2011). Relational regulation theory: A new approach to explain the link
52 between perceived social support and mental health. *Psychological Review*, 118(3), 482–
53 495. doi: 10.1037/a0023477
54
55
56
57
58
59
60

- 1
2
3 Larkin, M., & Thompson, A. R. (2012). Interpretative Phenomenological Analysis in mental
4 health and psychotherapy research. In D. Harper & A. R. Thompson (Eds.), *Qualitative*
5 *research method in mental health and psychotherapy: A guide for students and*
6 *practitioners* (pp. 101–116). Chichester, UK: John Wiley & Sons.
7
8
9
10
11
12 Lasker, J. N., & Toedter, L. J. (1991). Acute versus chronic grief: The case of pregnancy loss.
13 *American Journal of Orthopsychiatry*, 61(4), 510–522. doi: 10.1037/h0079288
14
15
16
17
18 Lazarus, R. S., & Folkman, S. (1984). *Stress, Appraisal, and Coping*. New York: Springer
19 Publishing Company.
20
21
22
23 Major, B., Appelbaum, M., Beckman, L., Dutton, M. A., Russo, N. F., & West, C. (2009).
24 Abortion and mental health: Evaluating the evidence. *American Psychologist*, 64(9),
25 863–890. doi: 10.1037/a0017497
26
27
28
29
30 Marshall, J., & Raynor, M. (2002). Conscientious objection: Legal and ethical issues. *British*
31 *Journal of Midwifery*, 10(6), 388–392.
32
33
34
35 McCoyd, J. L. M. (2007). Pregnancy interrupted: Loss of a desired pregnancy after diagnosis of
36 fetal anomaly. *Journal of Psychosomatic Obstetrics & Gynecology*, 28(1), 37–48. doi:
37 10.1080/01674820601096153
38
39
40
41
42 McCoyd, J. L. M. (2008). “I’m not a saint”: Burden assessment as an unrecognized factor in
43 prenatal decision making. *Qualitative Health Research*, 18(11), 1489–1500. doi:
44 10.1177/1049732308325642
45
46
47
48
49 McCoyd, J. L. M. (2009). Discrepant feeling rules and unscripted emotion work: Women coping
50 with termination for fetal anomaly. *American Journal of Orthopsychiatry*, 79(4), 441–
51 451. doi: 10.1037/a0010483
52
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- McCoyd, J. L. M. (2010). Women in no man's land: The abortion debate in the USA and women terminating desired pregnancies due to fetal anomaly. *British Journal of Social Work*, 40(1), 133–153. doi: 10.1093/bjsw/bcn080
- McDonald, S., Hetrick, S., & Green, S. (2004). Pre-operative education for hip and knee replacement. *The Cochrane Library*, 4. doi: 10.1002/14651858.CD003526.pub2
- Mulveen, R., & Hepworth, J. (2006). An Interpretative Phenomenological Analysis of participation in a pro-anorexia internet site and its relationship with disordered eating. *Journal of Health Psychology*, 11(2), 283–296. doi: 10.1177/1359105306061187
- National Collaborating Centre for Mental Health. (2011). *Induced abortion and mental health: A systematic review of the mental health outcomes of induced abortion including their prevalence and associating factors* (Report). London: Academy of Medical Royal Colleges, 2011.
- National Institute for Health and Clinical Excellence. (2007). *Antenatal and postnatal mental health. Clinical management and service guidance*. Retrieved from <http://www.nice.org.uk/Guidance/CG45/QuickRefGuide/pdf/English.p9>.
- Palmer, M., Larkin, M., de Visser, R., & Fadden, G. (2010). Developing an interpretative phenomenological approach to focus group data. *Qualitative Research in Psychology*, 7(2), 99–121. doi: 10.1080/14780880802513194
- Pennebaker, J. W. (2004). Theories, therapies, and taxpayers: On the complexities of the expressive writing paradigm. *Clinical Psychology: Science and Practice*, 11(2), 138–141. doi: 10.1093/clipsy/bph063
- Rillstone, P., & Hutchinson, S. A. (2001). Managing the reemergence of anguish: Pregnancy after a loss due to anomalies. *Journal of Obstetric, Gynecologic, & Neonatal Nursing*, 30(3), 291–298. doi: 10.1111/j.1552-6909.2001.tb01547.x

- 1
2
3 Robson, F. M. (2002). "Yes!—A chance to tell my side of the story": A case study of a male
4
5 partner of a woman undergoing termination of pregnancy for fetal abnormality. *Journal*
6
7 *of Health Psychology*, 7(2), 183–193. doi: 10.1177/1359105302007002457
8
9
- 10 Roth, S., & Cohen, L. J. (1986). Approach, avoidance, and coping with stress. *American*
11
12 *Psychologist*, 41(7), 813–819. doi: 10.1037/0003-066X.41.7.813
13
14
- 15 Royal College of Obstetricians and Gynaecologists. (2009). *RCOG statement on later maternal*
16
17 *age*. Retrieved from [www.rcog.org.uk/what-we-do/campaigning-and](http://www.rcog.org.uk/what-we-do/campaigning-and-opinions/statement/rcog-statement-later-maternal-age)
18
19 [opinions/statement/rcog-statement-later-maternal-age](http://www.rcog.org.uk/what-we-do/campaigning-and-opinions/statement/rcog-statement-later-maternal-age)
20
21
- 22 Salvesen, K. A., Oyen, L., Schmidt, N., Malt, U. F., & Eik-Nes, S. H. (1997). Comparison of
23
24 long-term psychological responses of women after pregnancy of termination due to fetal
25
26 anomalies and after perinatal loss. *Ultrasound in Obstetrics and Gynecology*, 9(2), 80–
27
28 85. doi: 10.1046/j.1469-0705.1997.09020080.x
29
30
- 31 Sandelowski, M., & Barroso, J. (2005). The travesty of choosing after positive prenatal
32
33 diagnosis. *Journal of Obstetric, Gynecologic, & Neonatal Nursing*, 34(3), 307–318. doi:
34
35 10.1177/0884217505276291
36
37
- 38 Schnider, K. R., Elhai, J. D., & Gray, M. J. (2007). Coping style use predicts posttraumatic stress
39
40 and complicated grief symptom severity among college students reporting a traumatic
41
42 loss. *Journal of Counseling Psychology*, 54(3), 344–350. doi: 10.1037/0022-
43
44 0167.54.3.344
45
46
47
- 48 Skinner, E. A., Edge, K., Altman, J., & Sherwood, H. (2003). Searching for the structure of
49
50 coping: A review and critique of category systems for classifying ways of coping.
51
52 *Psychological Bulletin*, 129(2), 216–269. doi: 10.1037/0033-2909.129.2.216
53
54
55
56
57
58
59
60

- 1
2
3 Shuldham, C. (1999). A review of the impact of pre-operative education on recovery from
4 surgery. *International Journal of Nursing Studies*, 36(2), 171–177. Retrieved from
5 [http://dx.doi.org/10.1016/S0020-7489\(99\)00010-3](http://dx.doi.org/10.1016/S0020-7489(99)00010-3),
6
7
8
9
10 Smith, J. A., Flowers, P., & Larkin, M. (2009). *Interpretative Phenomenological Analysis:*
11 *Theory, method and research*. Los Angeles, California: SAGE.
12
13
14
15 Statham, H., Solomou, W., & Green, J.M. (2001). When a baby has an abnormality: A study of
16 parents' experiences. Report to the NHS Exec (Mother and Child Health Initiative) of
17 grant no. MCH 4-12.
18
19
20
21
22 Strickland, S. (2011). Conscientious objection in medical students: a questionnaire survey.
23 *Journal of Medical Ethics*, 38(1), 22–25. doi:10.1136/jme.2011.042770.
24
25
26
27 Van de Mortel, T. F. (2008). Faking it: Social desirability response bias in self-report research.
28 *Australian Journal of Advanced Nursing*, 25(4), 40–48.
29
30
31
32 Warner, L. M., Schüz, B., Wurm, S., Ziegelman, J. P., & Tesch-Römer, C. (2010). Giving and
33 taking—differential effects of providing, receiving and anticipating emotional support on
34 quality of life in adults with multiple illnesses. *Journal of Health Psychology*, 15(5),
35 660–670. doi: 10.1177/1359105310368186
36
37
38
39
40
41 White-Van Mourik, M. C. A., Connor, J. M., & Ferguson-Smith, M. A. (1992). The psychosocial
42 sequelae of a second-trimester termination of pregnancy for fetal abnormality. *Prenatal*
43 *Diagnosis*, 12(3), 189–204. doi: 10.1002/pd.1970120308
44
45
46
47

48 Bios

49
50 **Caroline Lafarge**, MSc, MA, is a researcher in health psychology in the School of Psychology,
51 Social Work and Human Sciences, University of West London, London, UK.
52

53 **Kathryn Mitchell**, PhD, CPsychol, is deputy vice-chancellor, University of West London,
54 London, UK
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Pauline Fox, PhD, CPsychol, is a principal lecturer in health psychology in the School of Psychology, Social Work and Human Sciences, University of West London, London, UK.

For Peer Review