

The misuse and overuse of clinical gloves: a validated audit tool to define the problem

Journal:	<i>Journal of Infection Prevention</i>
Manuscript ID:	Draft
Manuscript Type:	Original Article
Keyword:	Behaviour, Hand Hygiene, Infection Control, Standard Precautions
Abstract:	<p>Background: The use of clinical gloves has become routine in the delivery of healthcare, often for procedures for which they are not required, their use may increase the risk of cross-contamination, and is generally not integrated into hand hygiene audit. This paper describes a small-scale application and validation of an observational audit tool devised to identify inappropriate glove-use and potential for cross-contamination.</p> <p>Methods: Two observers simultaneously observed the glove-use during episodes of care in an acute hospital setting. The inter-rater reliability (IRR) of the audit tool was measured corrected for chance agreement using kappa.</p> <p>Results: A total of 22 of episodes of care using gloves were observed. In 68.6% (24/35) of procedures there was no contact with blood/body fluid; in 54.3% (19/35) glove-use was inappropriate. The IRR was 100% for eight of 12 components of the tool. For hand hygiene before and after glove removal it was 82% (Kappa = 0.72) and 95% (Kappa = 0.87) .</p> <p>Conclusions: In this small-scale application of a glove-use audit tool we demonstrated over-use and misuse of gloves and the potential for cross transmission on gloved hands. The audit tool provides an effective mechanism for integrating glove-use into the audit of hand hygiene behaviour.</p>

SCHOLARONE™
Manuscripts

Clinical glove use audit
Commence observation when healthcare worker observed to be commencing a task/procedure and continue until the procedure is completed and gloves are removed or hands decontaminated

Ward:		Date:		Time start:			
				Time stop:			
1. Discipline	<input type="checkbox"/> Senior nurse	<input type="checkbox"/> Domestic	2. Location of task performed?	<input type="checkbox"/> Clean utility			
	<input type="checkbox"/> Sister	<input type="checkbox"/> Phlebotomist		<input type="checkbox"/> Sluice			
	<input type="checkbox"/> Staff nurse	<input type="checkbox"/> Porter		<input type="checkbox"/> Room/bay			
	<input type="checkbox"/> Agency Nurse	<input type="checkbox"/> AHP		<input type="checkbox"/> At bedside			
	<input type="checkbox"/> HCA	<input type="checkbox"/> Junior doctor		<input type="checkbox"/> Other			
<input type="checkbox"/> Student nurse	<input type="checkbox"/> Senior doctor		<input type="checkbox"/> Unknown				
3. Sequence of items/objects touched in this episode of care with points of hand hygiene/glove use <i>Use to categorise the risk of cross-contamination in one or more of 'My 5 moments of hand hygiene' at end of the observation</i>							
	Item	HH	G		Item	HH	G
1				7			
2				8			
3				9			
4				10			
5				11			
6				12			
4. If gloves used?							
Location put on?		Location removed?					
<input type="checkbox"/> Clean utility		<input type="checkbox"/> Clean utility					
<input type="checkbox"/> Sluice		<input type="checkbox"/> Sluice					
<input type="checkbox"/> Room/bay		<input type="checkbox"/> Room/bay					
<input type="checkbox"/> At bedside		<input type="checkbox"/> At bedside					
<input type="checkbox"/> Other		<input type="checkbox"/> Other					
<input type="checkbox"/> Unknown		<input type="checkbox"/> Unknown					
Was this as close to the point of use as possible/practical? <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> U/K		Was this as close to the point of use as possible/practical? <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> U/K					
5. Adequate hand hygiene after gloves removed? <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> U/K <i>According to local policy</i>							
Analysis							
Procedure(s) performed during this care episode							
Procedure <i>Check with healthcare worker if cannot observe</i>	Glove used?	Risk of contact with BBF? <i>Was/highly likely to be contact</i>	Glove use appropriate*?				
1	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> U/K	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> U/K	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> U/K				
2	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> U/K	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> U/K	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> U/K				
3	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> U/K	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> U/K	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> U/K				
4	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> U/K	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> U/K	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> U/K				
<small>* risk of contact with BBF, mucous membranes, hazardous substances (e.g. chemicals, cytotoxic drugs) or patient under isolation precaution.</small>							
Was there a risk of cross contamination? <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> U/K		If yes, which 'moments of hand hygiene'?					
		1 <input type="checkbox"/> Before contact with patient zone					
		2 <input type="checkbox"/> Before contact with susceptible site					
		3 <input type="checkbox"/> After contact with blood/body fluid					
		4 <input type="checkbox"/> After contact with patient zone					
		5 <input type="checkbox"/> After contact with healthcare zone					
Comment <i>(indicate if gloves required by local policy e.g. Isolation Precautions, handling chemicals etc)</i>							

HCA = healthcare assistant; AHP = Allied health professional e.g. physiotherapist, occupational therapist, pharmacist.

University of West London

Figure 1: The glove use audit tool
216x212mm (96 x 96 DPI)



The misuse and overuse of clinical gloves: a validated audit tool to define the problem

Word count: 4046

Abstract

Background: The use of clinical gloves has become routine in the delivery of healthcare, often for procedures for which they are not required, their use may increase the risk of cross-contamination, and is generally not integrated into hand hygiene audit. This paper describes a small-scale application and validation of an observational audit tool devised to identify inappropriate glove-use and potential for cross-contamination.

Methods: Two observers simultaneously observed the glove-use during episodes of care in an acute hospital setting. The inter-rater reliability (IRR) of the audit tool was measured corrected for chance agreement using kappa.

Results: A total of 22 of episodes of care using gloves were observed. In 68.6% (24/35) of procedures there was no contact with blood/body fluid; in 54.3% (19/35) glove-use was inappropriate. The IRR was 100% for eight of 12 components of the tool. For hand hygiene before and after glove removal it was 82% (Kappa = 0.72) and 95% (Kappa = 0.87) .

Conclusions: In this small-scale application of a glove-use audit tool we demonstrated over-use and misuse of gloves and the potential for cross transmission on gloved hands. The audit tool provides an effective mechanism for integrating glove-use into the audit of hand hygiene behaviour.

Background

The hands of staff are recognised to play a major role in the transmission of infection in healthcare settings (World Health Organisation, 2009). Hand hygiene is now accepted as an essential component of

infection control programmes, and education on hand hygiene is commonly based on the promotion of the '5 moments of hand hygiene' (5MHH) (Sax *et al*, 2007; World Health Organisation, 2009).

The routine use of gloves in clinical care emerged in the late 1980s when the Centre for Disease Control (CDC) recommended 'universal precautions' in response to the emerging AIDS epidemic (CDC, 1987).

These precautions required the use of protective clothing for direct contact with blood and some body fluids from any patient on the basis that it was not possible to discriminate those carrying blood borne viruses (CDC, 1988). Subsequently the recommendations were extended to all body fluids to address the increased risks of contamination associated with such contact and became known as standard precautions (Lynch *et al*, 1987; Wilson & Breedon, 1990; Pratt *et al*, 2001). Thus standard precautions indicate that personal protective equipment (PPE) should be used for any procedure where a risk of direct contact with blood and body fluids (BBF) is anticipated (Loveday *et al* 2014a; RCN 2012). The type of PPE selected should be dictated by the extent of potential exposure to BBF and as most contact involves hands, disposable gloves are the most commonly used form of PPE. However, adequate hand hygiene after glove use and the need to change gloves between procedures on the same patient and between patients is required to minimise the risk of infection transmission (World Health Organisation, 2009; Loveday *et al*, 2014a). If gloves are worn during the delivery of care, but not removed at the points in care where hand hygiene is indicated by the 5MHH, then their use will increase rather than decrease the risk of transmission of infection between patients. Although most studies focus on hand

1
2
3 hygiene rather than glove use behaviour, a few have indicated an emerging problem with an overuse of
4 gloves, with staff wearing them in for procedures that do not involve exposure to BBF and not removing
5 them in a timely way (Thompson *et al*, 1997; Prieto & Macleod Clark, 2005). Other studies have
6 suggested that the use of clinical gloves may have an adverse effect on compliance with the 5MHH
7 (Fuller *et al*,2011, Flores & Pevalin, 2006).
8
9
10
11
12
13
14
15
16

17 A mechanism of feeding back of data on the misuse of gloves is required in order to educate clinical staff
18 about using gloves appropriately and recognising the importance of hand hygiene in the context of glove
19 use. Most hospitals in the UK measure compliance with 5MHH using simple audit tools but these do not
20 account for the use of gloves in the delivery of care and are not able to capture data on the specific risks
21 of cross contamination associated with the use of gloves. In a previous study we designed and tested an
22 observational audit tool to identify inappropriate glove-use and potential for cross contamination
23 associated with the use of gloves (Loveday *et al*, 2014b). We found glove use was inappropriate in 42%
24 (69 of 163) of episodes and a risk of cross contamination in 37% (60 of 163) because an indication for
25 hand hygiene was missed. This study we describe components of the glove-use audit tool and report on
26 the inter-rater reliability associated with its use.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 **Method**

44 The audit tool has been designed to capture the detail of clinical procedures performed whilst wearing
45 gloves, the items that were touched during their use and the point when gloves were removed and hand
46 washed. This approach enables glove use behaviour to be analysed and the risk of cross contamination
47 associated with glove use to be measured. Although the audit tool was initially designed to be used for
48 recording events related to the use of a single pair of gloves (Loveday *et al*, 2014), we have subsequently
49 adapted it in order to record points during an episode of care where gloves are put on or removed and
50
51
52
53
54
55
56
57
58
59
60

1
2
3 hands decontaminated (Figure 1). The advantage of this approach is that it enables the extent of misuse
4
5 of gloves to be estimated by measuring the proportion of episodes of care in which gloves are used. The
6
7 audit episode commences once the healthcare worker (HCW) is observed to be preparing to undertake
8
9 an episode of care and ends once the HCW completes the episode. Each item touched with the hands
10
11 during the procedure/s, the order that they are touched and the point at which hand hygiene occurs or
12
13 gloves put on/taken off is recorded. On completion of the observation, the information is used to
14
15 categorise the potential for cross-contamination during each procedure observed using the criteria
16
17 shown in Table 1. These criteria have been adapted from the 5MHH for use in the context of the gloved
18
19 hand with a 'moment of HH' translating to the requirement to remove or change gloves and/or
20
21 decontaminate hands. Lack of hand hygiene before putting on gloves was not considered as a risk of
22
23 cross contamination (Rock *et al*, 2013). Hand hygiene after glove removal was considered adequate if
24
25 performed according to the hospital protocol.
26
27
28
29
30

31
32 Appropriateness of glove use was also assessed for each procedure. Glove use was considered
33
34 appropriate if the procedure was in the high-risk category of the Fulkerson scale and therefore involved
35
36 contact with, or a risk of contact with BBF (or hazardous substances, mucous membranes) or gloves
37
38 were required because of local policy (e.g. patient under isolation precautions) (McLaws *et al*, 2009).
39
40
41
42

43
44 We tested the tool in three wards in a large, acute teaching hospital. In order to minimise the
45
46 Hawthorne effect, the purpose of the observation would be made obscure. In general staff will assume
47
48 the observation is related to hand hygiene rather than the use of gloves and they are therefore less
49
50 likely to change their glove use behaviour during the observation. In addition, auditors would be
51
52 present on the ward for at least 10 minutes prior to collecting data the staff become more familiar with
53
54 their presence and are more likely to exhibit their normal behaviour. Positioning of the auditor is
55
56 important to ensure that observations can be carried out un-obtrusively, however, often clinical tasks or
57
58
59
60

1
2
3 procedures require the patient to be given privacy and care may be delivered behind curtains. Where
4 possible, observers would deduce procedures being undertaken by the equipment being used and
5 confirm by questioning the healthcare worker or patient. Ethical approval was not required for this
6 study as the observations constituted audit of practice and were a normal part of infection control
7 activity.
8
9
10
11
12
13
14
15
16

17 To determine the validity of the audit tool paired observations were captured simultaneously by two
18 observers. The inter-rater reliability (IRR) for each of 11 items documented in the tool was assessed
19 using percentage agreement between observers and corrected for chance using Cohen's kappa
20 coefficient in Excel. Kappa values of 0.41 to 0.6 are considered to demonstrate moderate agreement,
21 0.61 to 0.8 good and 0.81 or more very good agreement (Viera & Garrett ,2005).
22
23
24
25
26
27
28
29
30

31 **Results**

32
33 Twenty-two episodes of care were observed in a medical ward (9), high dependency unit (6) and
34 neurosurgical ward (7), all of which involved the use of gloves. These were performed by staff nurses (9
35 episodes), healthcare assistants (9 episodes), student nurses (2 episodes) and allied health
36 professionals/phlebotomists (2 episodes). There were 35 procedures performed during these episodes
37 of which 34 were performed using gloves. The number of procedures where there was contact with
38 blood or body fluid and where glove use was deemed appropriate is shown in Table 2.
39
40
41
42
43
44
45
46
47
48

49 Gloves were put on in the bay or outside the patient's room in 17 of the 22 episodes (77%). A total of
50 54 items were observed to be touched using gloves. On one of the 22 episodes (5%) a risk of cross
51 contamination was observed after Moment 4. In a further six of the 22 episodes (27%) the curtains
52
53
54
55
56
57
58
59
60

1
2
3 surrounding the patient's bed were touched before contact with the patient and, since this is deemed to
4
5 be outside the patient zone, it contravened moment 1.
6
7

8
9
10 The validation of the observational audit found high IRR in the paired observations for eight of 12
11
12 variables documented in the tool, including the appropriateness of glove use and risk of cross
13
14 contamination (see Table 3). In the five discrepant observations related to hand hygiene before or after
15
16 glove removal four related to hand hygiene being documented as 'unknown' rather than 'yes' or 'no'. In
17
18 2 of the 22 episodes there was a minor difference in documentation of the exact point where gloves
19
20 were removed (IRR 91%; Kappa 0.48); although this Kappa coefficient is likely to be underestimated
21
22 because disagreement with this variable was rare (Viera & Garrett 2005) (Table 3). In addition, there
23
24 was one minor variation in the description of the procedure being performed (Hygiene needs vs. tidying
25
26 bed/bedspace) and nine differences noted in specific items touched out of a total of 54 documented
27
28 items touched (IRR 83%) (Table 4).
29
30
31
32
33
34
35
36

37 Discussion

38
39
40 Our validation study demonstrated a high level of agreement between observers in documenting the
41
42 use of gloves using this audit tool, with an inter-observer agreement of 100% for the majority of
43
44 variables, including appropriateness of use and risk of cross-contamination, and Kappa scores of more
45
46 than 0.7. In this small set of observations we found that gloves were worn for more than 50% of
47
48 procedures but not required for 67% of these as there was no risk of contact with BBF or indications for
49
50 their use. In 27% of episodes of care we also found evidence that glove use was associated with a risk of
51
52 cross contamination, between the environment and patients or between patients, because gloves were
53
54 put on at a point distant to the patient and therefore contaminated outside the patient zone prior to
55
56
57
58
59
60

1
2
3 contact. The actual risk of cross-infection associated with observed glove-use practice has been
4
5 demonstrated by Girou *et al* who sampled gloves after use and recovered pathogens from 86%, even
6
7 after the application of alcohol hand rub (Girou *et al*, 2004). Snyder *et al* also found that gloves become
8
9 readily contaminated with antibiotic-resistant pathogens as a result of contact with patients or their
10
11 environment (Snyder *et al*, 2008). The widespread use of gloves was observed by Fuller *et al* (2011) who
12
13 found that 26% of hand hygiene opportunities involved the use of gloves. Other studies have found
14
15 poor compliance with timely removal of gloves where healthcare episodes which involve more than one
16
17 procedure on the same patient, particularly in the context of patient isolation for infection (Patterson *et*
18
19 *al*, 1991; Johnstone *et al*, 1990; Prieto & Macleod Clark, 2005).
20
21
22
23
24
25

26 An earlier version of the tool reported in this paper was used in a larger study in a different acute care
27
28 hospital, where we found that the decision of healthcare workers to use gloves appeared to be
29
30 influenced by feelings of disgust and misjudgement about the risk to self, and that glove-use behaviour
31
32 was often influenced by co-workers (Loveday *et al*, 2014b). Whilst we cannot be sure about the origin of
33
34 these sentiments, they may have emerged from the widespread publicity about risk of HCAI, strong
35
36 promotion of hand hygiene as fundamental to preventing infection, and from the promotion of personal
37
38 protective clothing as part of standard precautions.
39
40
41
42
43
44

45 Whilst promotion the 5MHH framework and the use of alcohol hand gel has become the standard
46
47 approach to education and training on hand hygiene in healthcare settings (WHO, 2009), the use of
48
49 gloves does not fit easily with these principles. Guidance on use of 5MHH suggests that indications for
50
51 hand hygiene are independent of those that justify the use of gloves and glove use should not replace or
52
53 alter the performance of hand hygiene. However, when 5MHH was conceived it was assumed that
54
55 gloves would be used as '*a second skin to prevent exposure of hand to body fluids*' and that '*glove*
56
57
58
59
60

1
2
3 *removal represents a strong cue to hand hygiene'* (Sax *et al* 2007). Unfortunately, our observations
4
5 suggest that the use of gloves has now been extended to a wide range of clinical activities that do not
6
7 involve exposure to body fluids and their use compromises the principles of the 5MHH because they are
8
9 frequently donned outside the patient zone and, in the absence of exposure to body fluid, the trigger to
10
11 remove them and perform hand hygiene may be lost. For example, we observed that it was common
12
13 practice to put on gloves in the bay where the glove dispenser was situated or outside patient's room
14
15 rather than at the bedside. As a result curtains and other equipment outside the patient zone were
16
17 touched by the gloved hand before contact with the patient. Similarly, donning gloves in the bay
18
19 precluded the application of alcohol hand gel immediately prior to contact with the patient. Therefore,
20
21 in using gloves to reduce the risk of infection, HCWs may actually increase the risk of transmission
22
23 between the environment and patient and between patients through lack of their timely application and
24
25 removal.
26
27
28
29
30
31
32

33 The audit tool also highlights other inconsistencies with 5MHH that are difficult to reconcile, for example
34
35 a commode moves from outside to inside the patient zone and then returns outside after use, which
36
37 makes it difficult to categorise the relevant moments of hand hygiene and/or glove removal. In addition,
38
39 the surfaces within the patient zone are considered to be continuous with the patient as they are likely
40
41 to become readily contaminated with their micro-organisms. However, the true microbiological risks
42
43 associated with this approach are unknown.
44
45
46
47
48

49 There are few examples in the literature of studies evaluating inter-observer agreement of hand hygiene
50
51 behaviour. McAteer *et al* used a simplified audit tool based on the assignment of hand hygiene
52
53 opportunities to six groups (before and after low risk contact, before and after high risk contact and
54
55 before and after unobserved contact). They found a *kappa* for hand hygiene opportunities and hand
56
57
58
59
60

1
2
3 hygiene actions of 0.68 and 0.77 respectively, but commented that use of the tool required clear
4 standard operating procedures and between 4 and 6 hours of training. In most healthcare settings
5
6 monitoring of compliance with hand hygiene is based on the WHO hand hygiene observation method,
7
8 although the recommendations to train and validate observers to ensure consistency is probably not
9
10 commonly performed (Sax *et al* 2009). Data on the IRR of this method is limited (Huis *et al* 2013, Steed
11
12
13
14
15 *et al* 2011).

16
17
18
19
20
21 We suggest that the standard approach to hand hygiene audit needs to be developed to address
22
23 inappropriate use of gloves and to more accurately reflect non-compliance with 5MHH in situations
24
25 where gloves are being used. We have shown that our audit tool can be used to provide consistent data
26
27 about the misuse of gloves and their potential impact on infection control, which infection prevention
28
29 teams can use to challenge the over-use of gloves and increase the knowledge and understanding of
30
31 healthcare workers about the hazards associated with their misuse. It does have limitations, in
32
33 particular it requires some expertise to interpret the risk of cross contamination, and the findings may
34
35 be subject to the Hawthorne effect (Holden 2001).
36
37
38

39
40 In conclusion, systematic audit of glove use behaviour indicates the lack of integration between glove-
41
42 use and 5MHH and a significant potential for cross transmission on gloved hands. Further work is
43
44 required to determine how widespread the observed glove-use practice is among clinical staff,
45
46 understand the drivers of this behaviour and to develop effective interventions to improve appropriate
47
48 use of clinical gloves.
49
50

51 52 53 54 55 56 **References** 57 58 59 60

Centers for Disease Control (1987). Recommendations for prevention of HIV transmission in health-care settings. *MMWR* **36** (suppl no. 2S).

Centers for Disease Control (1988). Update: universal precautions for prevention of transmission of human immunodeficiency virus, hepatitis B virus and other bloodborne pathogens in healthcare settings. *MMWR* **37**: 377-88

Chau JP, Thompson DR, Twinn S, Lee DT, Pang SW. (2011). An evaluation of hospital hand hygiene practice and glove use in Hong Kong. *J Clin Nurs*; **20** :1319-1328.

Flores A, Pevalin DJ. (2006). Healthcare workers' compliance with glove use and the effect of glove use on hand hygiene compliance. *Brit. J. Infect. Control.* **7**:15-19.

Fuller C, Savage J, Besser S, *et al.* (2011). "The dirty hand in the latex glove": A study of hand hygiene compliance when gloves are worn. *Infect. Control Hosp. Epidemiol.* **32**:1194-1199.

Holden JD. (2001). Hawthorne effects and research into professional practice. *J Eval Clin Pract.* **7**: 65-70.

Hripcsak G, Heitjan DF. (2002). Measuring agreement in medical informatics reliability studies. *J Biomed Inform.* **35**: 99-110.

1
2
3 Huis A, Schoonhoven L, Grol R *et al* (2013). Impact of a team and leaders-directed strategy to improve
4 nurses' adherence to hand hygiene guidelines: A cluster randomised trial. *Int. J. Nurs. Studies* **50**: 464–
5
6 474
7
8

9
10
11
12 Johnson S, Gerding DN, Olson MM, *et al.* (1990). Prospective, controlled study of vinyl glove use to
13 interrupt *Clostridium difficile* nosocomial transmission. *Am. J. Med.* **88**:137-140.
14
15

16
17
18
19 Loveday HP, Wilson JA, Pratt RP *et al* (2014a). epic3: National Evidence-Based Guidelines for Preventing
20 Healthcare-Associated Infections in NHS Hospitals in England. *J. Hosp. Infect.* **86S1** S1–S70
21
22

23
24
25
26 Loveday HP, Lynam S, Singleton J, Wilson J. (2014b). Clinical glove use: healthcare workers actions and
27 perceptions. *J. Hosp. Infect.* **86**: 110-116
28
29

30
31
32
33 Lynch P, Jackson M, Cummings MJ, Stamm WE. (1987). Rethinking the role of isolation practices in the
34 prevention of nosocomial infections. *Ann Int. Med.* **107**: 243-46.
35
36

37
38
39
40 McAteer J, Stone S, Fuller C, Charlett A, Cookson B, Slade R, Michie S. (2008) Development of an
41 observational measure of healthcare worker hand-hygiene behavior: the hand-hygiene observation tool
42 (HHOT). *J. Hosp. Infect.* **48**: 222-29
43
44
45

46
47
48
49 McLaws M-L, Pantle AC, Fitzpatrick KR, Hughes CF. (2009). Improvements in hand hygiene across New
50 South Wales public hospitals: Clean hands save lives, Part III. *Med. J. Aust.* **191**: S18-25.
51
52
53
54
55
56
57
58
59
60

1
2
3 Patterson JE, Vecchio J, Pantelick E, et al. (1991). Association of contaminated gloves with transmission
4 of *Acinetobacter calcoaceticus* var. *anitratus* in an intensive care unit. *Am J Med*;91:479-483.
5
6
7

8
9
10 Pratt RJ, Pellowe CM, Wilson JA, et al. (2007). Epic2 National Evidence-Based Guidelines for Preventing
11 Healthcare-Associated Infections in NHS Hospitals in England. *J. Hosp. Infect.* 65S: 1-64.
12
13
14

15
16
17 Prieto J and Macleod Clark J (2005). Contact precautions for *Clostridium difficile* and Methicillin-resistant
18 *Staphylococcus aureus* (MRSA): Assessing the impact of a supportive intervention to improve practice.
19 *Journal of Research in Nursing.* 10(5): 511-526.
20
21
22
23

24
25
26 Rock C, Harris AD, Reich NG et al (2013). Is hand hygiene before putting on nonsterile gloves in the
27 intensive care unit a waste of healthcare worker time? – a randomised controlled trial. *Am. J. Infect.*
28 *Contro.l* 41: 994-6
29
30
31
32

33
34
35 Royal College of Nursing (2012). Tools of the trade: RCN guidance for healthcare staff on glove use and
36 the prevention of contact dermatitis. Royal College of Nursing, London.
37
38
39

40
41
42 Sax H, Allegranzi B, Uckay L, Larson E, Boyce J, Pittet D (2007). “My five moments for hand hygiene”: a
43 user-centred design approach to understand, train, monitor and report hand hygiene. *J. Hosp. Infect.*
44 67:9-21.
45
46
47
48

49
50
51 Sax H, Allegranzi B, Chraiti M-N, Boyce J, Larson E, Pittet D. (2009) The World Health Organisation hand
52 hygiene method. *Am. J. Infect. Control* 37: 827-34
53
54
55
56
57
58
59
60

1
2
3 Snyder GM, Thom KA, Furono JP *et al.* (2008). Detection of methicillin-resistant *Staphylococcus aureus*
4 and vancomycin-resistant enterococci on the gowns and gloves of healthcare workers. *Infect. Control*
5
6
7
8 *Hosp. Epidemiol.* **29**: 583-89.

9
10
11
12 Steed C, Kelly JW, Blackhurst D, Boeker S, Diller T, Alper P, Larson E. (2011) Hospital hand hygiene
13 opportunities: where and when(HOW2)? The HOW2 benchmark study. *Am. J. Infect. Control.* **39**: 19-26
14
15

16
17
18
19 Thompson BL, Dwyer DM, Ussery XT, Denman S, Vacek P, Schwartz B. (1997). Handwashing and glove
20 use in a long-term-care facility. *Infect Control Hosp Epidemiol.***19**: 97-103.
21
22

23
24
25
26 Viera AJ, Garrett JM (2005). Understanding Interobserver Agreement: The Kappa Statistic. *Family*
27
28
29 *Medicine* 35: 360-3
30

31
32
33 Wilson J, Breedon P. (1990). Universal precautions. *Nurs. Times* **86 (37)**: 67-70.
34
35

36
37
38 World Health Organization (2009). WHO Patient Safety. *WHO guidelines on hand hygiene in health care.*
39

40 World Health Organization. Geneva.
41
42

43
44
45 WHO (2009) Hand hygiene technical reference manual. World Health Organization. Geneva.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

1
2
3 **Figure legends**
4
5
6

7 Figure 1: The glove use audit tool
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Table 1: Classification of the risk of cross contamination to incorporate the use of gloves (adapted from 'My 5 moments for hand hygiene' Sax *et al*, 2007)

Moment for hand decontamination	Risk of cross contamination	Definition
1	A patient touched by a contaminated glove/hand	Gloves/hands contaminated if they had contact with any part of the environment outside the patient's zone before direct contact with the patient's intact skin. <i>If the HCW touches their own clothing, skin or hair this is not considered part of the 'patient zone'</i>
2	A contaminated glove/hand touched a susceptible site e.g. wound, IV access site, phlebotomy	Gloves/hands contaminated if they had touched any other non-sterile objects or patient sites before the aseptic task e.g. patient skin, bed linen.
3	A glove/hand touched a surface or patient after contact with BBF	Gloves/hands contaminated if used for handling urine or assisting a patient with toileting then touched other surfaces or patient.
4	Gloves used for contact within patient zone not removed or hand hygiene not performed before contact with an object outside patient zone	Gloves/hands contaminated if touched another patient/objects outside patient zone; hand hygiene not performed after glove removal; or one glove/outer glove (where double-gloves used) removed part way through procedure.
5	Failure to remove gloves and/or perform hand hygiene after contact with patient	Gloves not removed or adequate hand hygiene not performed on leaving the healthcare zone.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

surroundings

For Peer Review

Table 2: Appropriateness of glove use for 34 procedures undertaken during episodes of care

	No		Yes		Unknown		Total
	No.	%	No.	%	No.	%	
Contact with BBF	23	67.6%	7	20.1%	4	11.4%	34 (100%)
Glove use appropriate	18	52.9%	15	44.1%	1	2.9%	34 (100%)

Table 3: Inter-observer agreement for variables in glove-use audit tool**a) Variables related to 22 episodes of care**

Item	Description	No. (%) agreement
1	Discipline of staff	22 (100%)
2	Location gloves put on	22 (100%)
3	Location task performed	22 (100%)
4	Gloving location appropriate	22 (100%)
5	Location gloves removed	22 (100%)

b) Variables related to glove removal and hand hygiene pre and post gloving during 22 episodes of care

Item	Description	No. in agreement			Total no. (%) agreement	Kappa
		Yes	No	UK		
6	Hand hygiene pre gloving	6	9	3	18 (82%)	0.72
7	Hand hygiene after removal	17	0	4	21 (95%)	0.87
8	Point of glove removal	20	2	-	20 (91%)	0.48

c) Variables related to 35 procedures observed during 22 episodes of care

Item	Description	No. (%) agreement
9	Procedure performed	35* (100%)
10	Risk of contact with BBF	35 (100%)
11	Glove use appropriate	35 (100%)
12	Risk of cross contamination	35 (100%)

*minor variation in description for one procedure

Table 4: Discrepancy in items observed to be touched during procedure

	Observer 1	Observer 2
1	Call button	<i>Not recorded</i>
2	<i>Not recorded</i>	Slide sheet
3	Shower chair	Commode
4	<i>Not recorded</i>	Toothbrush
5	Needle & syringe	<i>Not recorded</i>
6	<i>Not recorded</i>	Table
7	<i>Not recorded</i>	Sink
8	<i>Not recorded</i>	Patient property bag
9	Crockery	Zimmer frame