



UWL REPOSITORY

repository.uwl.ac.uk

Prevalence of overweight and obesity among adolescents in Bangladesh: do eating habits and physical activity have a gender differential effect?

Khan, Mostaued Ali, Karim, Masud, Islam, Ahmed Zohirul, Islam, Rafiqul, Khan, Hafiz T.A. ORCID logo ORCID: <https://orcid.org/0000-0002-1817-3730> and Khalilullah, Ibrahim (2019) Prevalence of overweight and obesity among adolescents in Bangladesh: do eating habits and physical activity have a gender differential effect? *Journal of Biosocial Science*, 51 (6). pp. 843-856. ISSN 0021-9320

<http://dx.doi.org/10.1017/S0021932019000142>

This is the Accepted Version of the final output.

UWL repository link: <https://repository.uwl.ac.uk/id/eprint/5785/>

Alternative formats: If you require this document in an alternative format, please contact: open.research@uwl.ac.uk

Copyright:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy: If you believe that this document breaches copyright, please contact us at open.research@uwl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

**PREVALENCE OF OVERWEIGHT AND OBESITY AMONG
ADOLESCENTS IN BANGLADESH: DOES FOOD HABITS AND
PHYSICAL ACTIVITIES HAVE GENDER DIFFERENTIAL
EFFECT?**

Journal:	<i>Journal of Biosocial Science</i>
Manuscript ID	JBS-4581.R2
Manuscript Type:	Research Article
Date Submitted by the Author:	29-Jan-2019
Complete List of Authors:	Khan, Md. Mostaufed; University of Rajshahi, Population Science and Human Resource Development Karim, Masud ; University of Rajshahi, Population Science and Human Resource Development Islam, Ahmed Zohirul; University of Rajshahi, Population Science & Human Resource Development Islam, Md. Rafiqul; University of Rajshahi, Population Science and Human Resource Development Khan, Hafiz ; University of West London, The Graduate School Khalilullah, Md. Ibrahim ; University of Rajshahi, Population Science and Human Resource Development
Keywords:	Obesity, Child Health and Nutrition

SCHOLARONE™
Manuscripts

PREVALENCE OF OVERWEIGHT AND OBESITY AMONG ADOLESCENTS IN BANGLADESH: *DOES FOOD HABITS AND PHYSICAL ACTIVITIES HAVE GENDER DIFFERENTIAL EFFECT?*

Authors: Md. Mostaured Ali Khan¹, Masud Karim¹, Ahmed Zohirul Islam¹, Md. Rafiqul Islam^{1*}, Hafiz T. A. Khan² and Md. Ibrahim Khalilullah¹

¹ Department of Population Science and Human Resource Development,
University of Rajshahi, Rajshahi-6205, Bangladesh.

² The Graduate School,
University of West London,
St Mary's Road, Ealing, London W5 5RF, United Kingdom.

****Corresponding Author:***

Md. Rafiqul Islam
Professor, Department of Population Science and Human Resource Development,
University of Rajshahi, Rajshahi-6205, Bangladesh.

Email address of the corresponding author: rafique_pops@yahoo.com

Keywords: Obesity; Adolescent; Food habit; Physical activities; Bangladesh.

Abstract

The aim of the study is to examine the gender differential outcomes of food habits and physical activities on obesity among school-aged adolescents in Bangladesh. Nationally representative data extracted from the 2014 Global School-based Student Health Survey (GSHS) were utilized. The information related to physical and mental health was collected from 2989 school-aged adolescents in Bangladesh. To fulfill the study aims, an exploratory data analysis and multivariate logistic regression model were employed. Female adolescents were at a lower risk of being overweight or obese (AOR = 0.573) with a prevalence of 7.4% compared to males (9.9%). The results showed that high consumption of vegetables (both: AOR = 0.454; males: AOR = 0.504; and females: AOR = 0.432), high soft drink consumption (both: AOR=2.357; males: AOR = 2.929; and females: AOR = 1.677), high fast food consumption (both: AOR = 2.777; males: AOR = 6.064; and females: AOR = 1.695), sleep disturbance (both: AOR = 0.675; males: AOR = 0.590; and females: AOR = 0.555), and regular walking or cycling to school (both: AOR = 0.472; males: AOR = 0.430; and females: AOR = 0.557) were vital influencing factors for being overweight or obese among adolescents across both sexes. Sedentary activities during leisure time were also identified as significant predictors of being overweight and obesity for males. Regular fruit and vegetable consumption, the avoidance of soft drinks and fast food, an increase in vigorous physical activity, regular attendance at physical education classes and less sedentary activities in leisure time could all help reduce the risk of being overweight or obese for both sexes.

Introduction

Obese and overweight children and adolescents represents one of the biggest challenges to face public health in the 21st century and is greatly affecting many low and middle-income countries (LMICs) (De Onis *et al.*, 2010; Peng *et al.*, 2017). The prevalence of obesity across the world has nearly trebled since 1975 (WHO, 2018). In 2016, over 1.9 billion adults and 340 million adolescents worldwide, including children, were found to be overweight or obese (WHO, 2018). A high risk of obesity was observed particularly for Asians and Pacific Islanders (Young *et al.*, 2017) although in South Asian countries, malnutrition (stunting, wasting and underweight) among children is a more hazardous situation. Problems with obesity are also a matter of vital concern in many developing countries including Bangladesh due to its flourishing economy (Shafique *et al.*, 2007). Since the year 2000, the increase in Body Mass Index (BMI) in East and South Asian countries in particular has accelerated swiftly for both sexes (Collaboration, 2017). Rapid urbanization and industrialization, plus economic development and globalization of food production are some of the important causal factors for this situation emerging in the developing world.

Previous research has identified the many negative aspects of being overweight or obese on the health and growth of children and adolescents that can extend into adulthood and increase the risk of developing chronic diseases such as cardiovascular disease (Singh *et al.*, 2013), chronic kidney disease (Singh *et al.*, 2013), diabetes, many cancers (Lauby-Secretan *et al.*, 2016), and disabilities (Dereń *et al.*, 2018). Furthermore, being overweight and obesity are significantly related to mortality (Di Angelantonio *et al.*, 2016; Flegal *et al.*, 2013).

Although there is a growing body of studies that have examined the various risk factors of being overweight and obese, there is no specific study that is focused on the gender

differential of obesity as a whole. Some studies have mentioned that the diverse food habits and physical activities of children have a significant impact on their weight (Virtanen *et al.*, 2015), as do other metabolic and socio-demographic factors (Hossain *et al.*, 2018). These factors include insufficient physical activity (Li *et al.*, 2017), shortened duration of sleep at night (Brug *et al.*, 2012), physical education (PE) at school (Naiman *et al.*, 2015) and physical activity (PA) facilities (Hood *et al.*, 2014). Diverse food habits involving the consumption of fast food (Davis & Carpenter, 2009; Rosenheck, 2008), low level of fruit and vegetable intake and high fat and sugar intake (Epstein *et al.*, 2012), food insecurity (Lyons *et al.*, 2008; Robaina & Martin, 2013) and poor diet quality (Robaina & Martin, 2013), were also found to be important determinants for overweight and obese children and adolescents.

In most developing countries, epidemiological studies on school-level risk factors for obesity are still inadequate and any differences in terms of gender are unknown. Males and females display differences in fat stores, dissimulation in anatomical fat distribution, and also in high food intake and low physical activity (Reue, 2017) that gives strength to this study. In Bangladesh, gender discrimination exists in all sectors including health and nutrition (Hossain *et al.*, 2018; Shafique *et al.*, 2007). A number of attempts have been made to uncover the risk factors of being overweight or obese but there has not been any research on the gender differential risk factors among children and adolescents. This study focuses on the prevalence of obesity and on ascertaining the gender differential outcomes of food habits and physical activity on overweight and obese school-aged adolescents in Bangladesh.

Methods

Study design and Sampling procedure

This study has used data extracted from the Global School-based Student Health Survey (GSHS) 2014. The survey collected data from school-age adolescents (usually aged

11-17 years) in 43 developing countries including Bangladesh and was administrated by the World Health Organization (WHO) in collaboration with the Center for Disease Control (CDC). Data were collected using a clustered sampling technique and a standardized scientific sample selection process, conventional school-based methodology, and a combination of core questionnaire modules with expanded questions plus country-specific questionnaires utilized by the survey. The school response rate was 90–100% with the student response rate ranged between 76–96% and the overall response rate at 69–96% for each of the countries. In Bangladesh, information related to dietary behaviors, hygiene, drug, tobacco and alcohol use, sexual behaviors, mental health, physical activity etc. and was collected by GSHS in 2014 from 2989 adolescents. Full clarifications of the study, including the core questionnaire used with items selected from pertinent modules, are available at the websites of CDC and WHO (C. WHO, 2017).

Calculation of BMI

The respondent’s BMI was calculated applying the following formula:

$$BMI = \frac{Weight\ (kg)}{Height^2\ (m)}$$

As all the respondents were up to 18 years of age, they were classed as being overweight if their calculated BMI exceeded the standardized value for age and sex at +1SD of Z scores of BMI (equivalent to BMI 25 kg/m² at 19 years of age). They were classed as being obese if their calculated BMI exceeded the standardized value for age and sex at +2SD of Z scores of BMI (equivalent to BMI 30 kg/m² at 19 years of age) on the basis of BMI interpretation provided by WHO (Onis *et al.*, 2007; WHO, 2015).

Outcome and Explanatory variables

To achieve the objective of the study, being overweight or obese were considered as dependent or outcome variables. The outcome variable was addressed as follows:

$$Y = \text{Overweight or obese} = \begin{cases} 1, & \text{Yes} \\ 0, & \text{No} \end{cases}$$

Several explanatory variables related to food insecurity, food habits, depression and physical activities were treated as risk factors for being overweight and obese with variables selected in accordance with their importance based on previous research. The information was categorized according to the recommendation provided by WHO (WHO, 2012). A complete list of explanatory variables is shown in table 1. (**Table 1**)

Statistical analysis

Any association between the state of being overweight and obese and different explanatory variables were assessed by Chi-square tests (usually, set at $p < 0.05$ level of significance). As the outcome variable of this study had two categories, the binary logistic regression model was fitted to measure the impact of selected explanatory variables on the outcome variable. In this study, the odds ratios (ORs) were estimated to assess the strength of association between the outcome variable and the explanatory variables, and 95% confidence intervals (CIs) were exerted to examine the level of significance. The data were analyzed using the computer program SPSS in Windows version 23.0 (SPSS Inc., Chicago, IL).

Results

Table 2 represents the characteristics of the respondents. The mean age was 14.2 (± 0.98) years, mean height was 1.563 meters (± 0.087), mean weight was 45.88 kilograms (± 7.868) and mean calculated BMI was 18.78 kg/m² (± 2.87). The prevalence of being overweight and obese was 9.9% for males and 7.4% for females. (**Table 2**)

Table 3 illustrates the association between being overweight and obesity and selected explanatory variables were pursued by applying a Chi-square test to observe the significance. In this study, the frequency of respondents to experience hunger, fruit and vegetable eating, consumption of soft drinks, fast food eating, sleep disturbance, physical activity (PA), physical education (PE) attendance were significantly related to being overweight or obesity for adolescents of both sexes. A high consumption of fast food had the highest prevalence of being overweight and obesity for males (25.3%), while the highest prevalence observed for females (13.1%) was among those that never attended PE classes. Male respondents with higher fruit (2.4%) and vegetable (5.4%) eating habits displayed a lower prevalence of being overweight or obese. Similarly, there were only 2.5% and 4.6% of females with high fruit and vegetable eating habits that were either overweight or obese. Alternatively, 18.7% of males and 12.4% of females who consumed soft drinks at a high frequency were overweight and obese. Only 5.9% of male and 4.9% of female respondents who were vigorously physically active were found to be overweight or obese. The frequency of being overweight and obese was lower among male and female respondents who walked or cycled to school (male: 5.5% and female: 4.8%) and attended PE classes regularly (male: 8.2% and female: 7.4%). There was a significant association among male respondents between a high amount of sitting or sedentary activities leading to a high prevalence of being overweight and obesity (19.2%). (Table 3)

Effect of food habits and physical activities on being overweight and obesity

Table 4 illustrates the effects of adolescent food habits and physical activities for being overweight and obese in Bangladesh. The occurrence of either state was decreased for female adolescents (AOR = 0.573, CI: 0.403-0.816) compared to male adolescents. Regular feelings of hunger at 2.789 (AOR = 2.789, CI: 1.733-4.489) times, highly accelerates the risk of being overweight and obese than among those that never feel hunger. A high consumption

of fruit (AOR = 0.454, CI: 0.205-0.997) and vegetables (AOR = 0.475, CI: 0.294-0.768) significantly diminished the risk of adolescents being overweight or obese. However, a high consumption of soft drinks (AOR = 2.357, CI: 1.544-3.597) and fast food (AOR = 2.777, CI: 1.755-4.392) significantly increased the risks. Adolescents with frequent sleep disturbances (AOR = 0.675, CI: 0.481-0.947) were found less likely to be overweight or obese. This was also the case for those adolescents that walked or cycled to school (AOR = 0.472, CI: 0.327-0.682) and attended regular PE classes (AOR = 0.592, CI: 0.327-0.682) when compared to those that never walked or cycled or attended PE.

The fitted model of Cox and Snell R^2 , and Nagelkerke R^2 was shown to be 61.0% and 81.3%, respectively and was estimated from the linear relationship between the independent variables. The overall model was significant when all independent variables were controlled for age. (Table 4)

Gender differential effect of food habits on being overweight and obesity

Table 5 shows the results of the logistic regression model of the gender differential influence of food patterns on being overweight and obesity among school-aged adolescents in Bangladesh. The likelihood of either of these states was decreased for males who sometimes went hungry (AOR = 1.399, CI: 1.036-1.891) or went hungry most of the time (AOR = 2.759, CI: 1.846-4.125) than it was for respondents that never went hungry. The risk of being overweight or obese was also decreased for males that ate a lot of fruit (AOR = 0.372, CI: 0.203-0.683). The occurrence was decreased for males with a high frequency of vegetable (AOR = 0.504, CI: 0.333-0.764) eating and reduced for females with an average (AOR = 0.582, CI: 0.372-0.910) or high vegetable eating habit (AOR = 0.432, CI: 0.248-0.753) compared to males and females with a low vegetable eating habit. However, males with an average soft drink consumption habit were at a higher risk of being overweight or obese (AOR = 2.583, CI: 1.855-3.597) as were adolescents with a high soft drinks' consumption

168 habit (male: AOR = 2.929, CI: 2.086-4.112; female: AOR = 1.677, CI: 1.022-2.753)
169 compared to adolescents whose weekly consumption of soft drinks was lower. High
170 consumption of fast food significantly increased the chances of ending up overweight or
171 obese for both sexes (male: AOR = 6.064, CI: 4.327-8.499; and female: AOR = 1.695, CI:
172 1.011-3.174) as it was for males with an average fast food eating habit (AOR = 1.503, CI:
173 1.084-2.083).

174 The fitted model of Cox and Snell R^2 , and Nagelkerke R^2 was shown to be 56.1%
175 and 74.7% respectively of the variance for males, and 58.9% and 78.6% respectively of the
176 variance for females and was estimated from the linear relationship between the independent
177 variables. The overall model was significant when all independent variables were controlled
178 for age. (Table 5)

179 *Gender differential effect of physical activities on being overweight and obesity*

180 The results of the logistic regression model shown in Table 6 illustrate the effect of
181 PA on being overweight and obesity among school-aged adolescents. Sleep disturbance was
182 found to have a significantly decreased association with obesity (male: AOR = 0.590, CI:
183 0.455-0.766; and female: AOR = 0.555, CI: 0.369-0.837). As would be expected, being
184 overweight or obese decreased among vigorously physically active males (AOR = 0.751, CI:
185 0.592-0.991) as it was for respondents who took part in moderate PA. The risk of being
186 overweight or obese was reduced for both males and females who occasionally walked or
187 cycled to school (male: AOR = 0.265, CI: 0.171-0.410; and female: AOR = 0.453, CI: 0.205-
188 0.924) or who regularly walked or cycled to school (male: AOR = 0.430, CI: 0.322-0.576;
189 and female: AOR = 0.557, CI: 0.359-0.866) compared to respondents that never walked or
190 cycled to school. The likelihood of being overweight or obese decreased among males and
191 females that occasionally attended PE classes (male: AOR = 0.420, CI: 0.281-0.627; and
192 female: AOR = 0.445, CI: 0.266-0.745) or for males that regularly attended such classes

(male: AOR = 0.488, CI: 0.330-0.722) compared to males and females that never attended PE. The risk of being overweight or obese was increased by 3.404 (AOR = 3.404, CI: 2.343-4.945) times higher for males with high sitting or sedentary activities compared to males with moderate sitting activities.

In the fitted model, the Cox and Snell R^2 , and Nagelkerke R^2 was 54.1% and 77.2% respectively. The variances of males and for females was 60.0% and 80.0% respectively, that can be estimated from the linear relationship between the independent variables. The overall model was significant when all explanatory variables were included. (Table 6)

Discussion

The prevalence of being overweight and obesity is showing an increasing trend in Bangladesh (Biswas *et al.*, 2017). It has not yet become an alarming situation for adolescents but it is increasing day-by-day. The findings of this study indicate that the risk of males being overweight or obese is notably higher than it is for females. This study findings also show that male adolescents with high food insecurity are at an increased risk of being overweight or obese that is consistent with earlier studies (Robaina & Martin, 2013; Sanjeevi *et al.*, 2018). As well as Sanjeevi *et al.* (2018) showing a uniformity with the results of our study, these authors concluded that food insecurity is associated with a less conducive multidimensional home environmental subscale score and poor diet quality that, in turn, is related to greater BMI. Lohman *et al.* (2016) also identified a gender differential outcome of household food insecurity for being overweight or obese. In Bangladesh, less importance is generally given to female children than to males in all sectors.

Dietary behavior and different food patterns have a diverse impact on being overweight or obesity (Rautiainen *et al.*, 2015; Virtanen *et al.*, 2015). This study has identified a significantly lower risk of both these states in adolescents with a high fruit and vegetable eating habit. A gender differential effect was seen again when such eating habits

significantly decreased the risk for male adolescents, but in the case of females, no significant effect due to fruit-eating was found but a highly significant effect was identified with average to higher vegetable consumption. Previous research supports this hypothesis by identifying that there is a significant positive impact of regular fruit and vegetable intake among children and adolescents (Epstein *et al.*, 2012; Field *et al.*, 2003). According to Rohde *et al.* (2017), responsible intake of macronutrients, energy, fruit and vegetables can help restrain excessive weight gain among children. More precisely, fruit and vegetables provide fiber, are low in calories and rich in minerals and vitamins that help to keep a person healthy and energized.

This study shows that a high consumption of soft drinks and fast food increases the risk for both male and female adolescents of becoming overweight or obese. As a consequence, adolescents are at a high risk of experiencing problems with their weight regardless of their intake of junk food. A number of previous studies have shown that high consumption of soft drinks and fast food has a highly negative effect on obesity in adolescents and young children (Davis & Carpenter, 2009; Moore *et al.*, 2009; Rosenheck, 2008). These types of food and drink contain more fat and sugar, and fewer vitamins and minerals than healthier alternatives and therefore can lead to poor weight management and body metabolism leading further to a risk of obesity (Lucan & DiNicolantonio, 2015). In recent experiments, researchers have shown that reducing soft drink and fast food consumption in adolescents has been successful in lessening the prevalence of obesity (Cantoral *et al.*, 2016; Hu, 2013; Laxy *et al.*, 2015). In addition, a high intake of artificially sweetened soft beverages enhances the risk of obesity-related cancers (Hodge *et al.*, 2018). In Bangladesh, the quality of soft drinks and fast food is much poorer than in developed countries that is perhaps a similar situation in other developing countries.

The study results show there is a lower risk of obesity among adolescents of both sexes who often face sleep problems due to depression, that is inconsistent with findings from

several previous studies (Brug *et al.*, 2012; Mannan *et al.*, 2016; Nielsen *et al.*, 2011). In addition, (Nielsen *et al.*, 2011) observed a significant link between short duration of sleep and being overweight or obesity among young adults including children. A meta-analysis and systematic review of longitudinal studies conducted by Mannan *et al.* (2016) revealed a 70% greater risk of depressed male and female adolescents being overweight or obese.

Physical activity (PA) is an emerging determinant of weight for both children and adults. The study found there was a much lower risk of male adolescents being overweight or obese if they were vigorously active compared to those who were moderately active. This hypothesis is supported by a few earlier studies such as Ogden *et al.* (2016). Also showing a consistency with the study findings, Chaput *et al.* (2018) noticed there was a lower risk of obesity by level of sedentary behavior in children who were vigorously physically active but found no significant effect of PA in case of overweight or obese males or females. Males and females that regularly walked or cycled to school were at a very low risk of being overweight or obese. Walking and cycling carry twofold advantages: they help protect the environment and prevent excessive weight gain by increasing body metabolism. Responsible parents should therefore encourage their children to regularly walk or cycle to school.

The attendance of adolescents at PE has also been identified as a feasible predictor of being overweight and obesity in Bangladesh. The respondents of both sexes that regularly attended PE had a very low risk of developing weight problems compared to those that never attended PE (Naiman *et al.*, 2015). Spending time at PE can help reduce the gap between actual and recommended physical activity for children and adolescents (Fernandes & Sturm, 2010) and also help increase the number of days per week spent in vigorous PA (Jinsook, 2012). So, it can be seen that PE indirectly affects excessive weight gain among adolescents and children. Unfortunately, however, PE facilities in Bangladesh are very poor and there is poor awareness of the benefits of PE among parents plus a lack of strict regulations.

268 Sedentary activities of adolescents show a negative impact on being overweight or obesity for
269 male adolescents but show no significant effects on female adolescents. Those male
270 adolescents with high levels of sitting activity per day are almost at three times higher risk of
271 becoming overweight or obese. This finding is supported by previous research related to
272 adults (Chau *et al.*, 2012; Ng *et al.*, 2017). In Australia, the risk of being overweight or obese
273 increases significantly among workers with mostly sitting jobs than workers with mostly
274 standing jobs (Chau *et al.*, 2012). However, there are no studies that describe the effect of
275 sitting behaviors as a cause of weight problems or obesity among children and adolescents.
276 Adolescent leisure time activities per day such as watching TV, gossiping, playing computer
277 games etc. increase their risk of becoming overweight or obese especially for male
278 adolescents.

279 This study had several limitations. For example, a secondary source of data was used
280 for analysis and thus some important variables are missing in relation to being overweight
281 and obesity. Nevertheless, an attempt has been made to depict a compact demonstration of
282 the effect of adolescent eating behaviors and PA on being overweight and obese. Future
283 studies could be undertaken to collect data covering variables involved in differences
284 between rural-urban areas.

285 In conclusion, findings from the study suggest that the levels of being overweight or
286 obesity among school-aged adolescents in Bangladesh have yet to be improved. This research
287 discovered gender differences in food practice and physical activity among adolescents that
288 affect their overweight or obesity levels and therefore indicates improvements are needed in
289 their behavior around eating and physical activity. A regular consumption of healthy food,
290 particularly a diet rich in fruit and vegetables and avoidance of soft drinks and fast food
291 especially for males, are necessary components for helping to lessen the risk of adolescents in
292 Bangladesh being overweight or obese. Increasing levels of physical activity, cutting back on

high levels of leisure time sitting activities, especially among males, and encouraging adolescents of both sexes to regularly walk or cycle to school, can all help to cut the risk of developing weight problems. Policy in this area should focus on the need for regular attendance at PE classes to help improve the health of school-age adolescents. The implementation of such policies would help decrease the risk of adolescent obesity in Bangladesh and, in turn, be helpful for prolonging their good health.

Abbreviations

GSHS, Global School-based Student Health Survey; BMI, Body Mass Index; SD, Standard Deviation; PE, Physical Education; PA, Physical Activity; WHO, World Health Organization; CDC, Center for Disease Control; CI, Confidence Interval; OR, Odds Ratio.

Acknowledgments

The authors are thankful to the Office of the Global School-based Student Health Survey (GSHS), Department of Chronic Diseases and Health Promotion, World Health Organization, jointly worked with Center for Disease Control (CDC), for permitting us to use the datasets for this analysis.

Ethical Approval

The study was ethically approved and the statement was obtained from the Ministry of Health and Family Welfare, Dhaka, Bangladesh. The World Health Organization (WHO) financially and technically supports this survey with the collaboration of the Center for Disease Control (CDC). Further, written permission was obtained from each participating school and from all classroom teachers.

Conflicts of interest

The authors have no competing interests to declare.

316 **Funding**

317 The authors received no funding grant from any funding agency, commercial entity or not-
318 for-profit organization for this study.

319 **References**

320 **Biswas T, Uddin MJ, Mamun AA, Pervin S and Garnett SP** (2017). Increasing prevalence of overweight and
321 obesity in Bangladeshi women of reproductive age: Findings from 2004 to 2014. *PLOS ONE*, **12**(7),
322 e0181080.

323 **Brug J, van Stralen MM, te Velde SJ, Chinapaw MJ, De Bourdeaudhuij I et al.** (2012). Differences in
324 weight status and energy-balance related behaviors among schoolchildren across Europe: the
325 ENERGY-project. *PLOS ONE*, **7**(4), e34742.

326 **Cantoral A, Téllez-Rojo MM, Ettinger A, Hu H, Hernández-Ávila M and Peterson K** (2016). Early
327 introduction and cumulative consumption of sugar-sweetened beverages during the pre-school period
328 and risk of obesity at 8–14 years of age. *Pediatric Obesity*, **11**(1), 68-74.

329 **Chaput JP, Barnes J, Tremblay M, Fogelholm M, Hu G et al.** (2018). Thresholds of physical activity
330 associated with obesity by level of sedentary behaviour in children. *Pediatric Obesity*, **13**, 450–457.

331 **Chau JY, van der Ploeg HP, Merom D, Chey T and Bauman AE** (2012). Cross-sectional associations
332 between occupational and leisure-time sitting, physical activity and obesity in working adults.
333 *Preventive Medicine*, **54**(3), 195-200.

334 **Collaboration NRF** (2017). Worldwide trends in body-mass index, underweight, overweight, and obesity from
335 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128·9 million
336 children, adolescents, and adults. *The Lancet*, **390**(10113), 2627-2642.

337 **Davis B and Carpenter C** (2009). Proximity of fast-food restaurants to schools and adolescent obesity.
338 *American Journal of Public Health*, **99**(3), 505-510.

339 **De Onis M, Blössner M and Borghi E** (2010). Global prevalence and trends of overweight and obesity among
340 preschool children–. *The American journal of clinical nutrition*, **92**(5), 1257-1264.

341 **Dereń K, Nyankovskyy S, Nyankovska O, Łuszczki E, Wyszynska J et al.** (2018). The prevalence of
342 underweight, overweight and obesity in children and adolescents from Ukraine. *Scientific Reports*,
343 **8**(1), 3625.

344 **Di Angelantonio E, Bhupathiraju SN, Wormser D, Gao P, Kaptoge S et al.** (2016). Body-mass index and
345 all-cause mortality: individual-participant-data meta-analysis of 239 prospective studies in four
346 continents. *The Lancet*, **388**(10046), 776-786.

347 **Epstein LH, Gordy CC, Raynor HA, Beddome M, Kilanowski CK and Paluch R** (2012). Increasing fruit
348 and vegetable intake and decreasing fat and sugar intake in families at risk for childhood obesity.
349 *Obesity*, **9**(3), 171-178.

350 **Fernandes M and Sturm R** (2010). Facility provision in elementary schools: Correlates with physical
351 education, recess, and obesity. *Preventive Medicine*, **50** (Suppl 1), S30-S35.

352 **Field AE, Gillman MW, Rosner B, Rockett HR and Colditz GA** (2003). Association between fruit and
353 vegetable intake and change in body mass index among a large sample of children and adolescents in
354 the United States. *International journal of obesity*, **27**(7), 821-826.

355 **Flegal KM, Kit BK, Orpana H and Graubard BI** (2013). Association of all-cause mortality with overweight
356 and obesity using standard body mass index categories: A systematic review and meta-analysis. *JAMA*,
357 **309**(1), 71-82.

358 **Hodge AM, Bassett JK, Milne RL, English DR and Giles GG** (2018). Consumption of sugar-sweetened and
359 artificially sweetened soft drinks and risk of obesity-related cancers. *Public health nutrition*, **21**(9),
360 1618-1626.

- Hood NE, Colabianchi N, Terry-McElrath YM, O'Malley PM and Johnston LD** (2014). Physical activity breaks and facilities in US secondary schools. *Journal of School Health*, **84**(11), 697-705.
- Hossain MM, Islam MR, Sarker ASR, Khan MMA and Taneepanichskul S** (2018). Prevalence and determinants risk factors of underweight and overweight among women in Bangladesh. *Obesity Medicine*, **11**, 1-5
- Hu FB** (2013). Resolved: there is sufficient scientific evidence that decreasing sugar-sweetened beverage consumption will reduce the prevalence of obesity and obesity-related diseases. *Obesity reviews*, **14**(8), 606-619.
- Jinsook K** (2012). Are Physical Education-Related State Policies and Schools' Physical Education Requirement Related to Children's Physical Activity and Obesity? *Journal of School Health*, **82**(6), 268-276.
- Lauby-Secretan B, Scoccianti C, Loomis D, Grosse Y, Bianchini F and Straif K** (2016). Body fatness and cancer—viewpoint of the IARC Working Group. *New England Journal of Medicine*, **375**(8), 794-798.
- Laxy M, Malecki KC, Givens ML, Walsh MC and Nieto FJ** (2015). The association between neighborhood economic hardship, the retail food environment, fast food intake, and obesity: findings from the Survey of the Health of Wisconsin. *BMC public health*, **15**(1), 237.
- Li M, Xue H, Wen M, Wang W and Wang Y** (2017). Nutrition and physical activity related school environment/policy factors and child obesity in China: a nationally representative study of 8573 students in 110 middle schools. *Pediatric Obesity*, **12**(6), 485-493.
- Lohman B J, Gillette MT and Neppel TK** (2016). Harsh Parenting and Food Insecurity in Adolescence: The Association With Emerging Adult Obesity. *Journal of Adolescent Health*, **59**(1), 123-127.
- Lucan SC and DiNicolantonio JJ** (2015). How calorie-focused thinking about obesity and related diseases may mislead and harm public health. An alternative. *Public health nutrition*, **18**(4), 571-581.
- Lyons AA, Park J and Nelson CH** (2008). Food Insecurity and Obesity: A Comparison of Self-Reported and Measured Height and Weight. *American Journal of Public Health*, **98**(4), 751-757.
- Mannan M, Mamun A, Doi S and Clavarino A** (2016). Prospective Associations between Depression and Obesity for Adolescent Males and Females- A Systematic Review and Meta-Analysis of Longitudinal Studies. *PLOS ONE*, **11**(6), e0157240.
- Moore LV, Diez Roux AV, Nettleton JA, Jacobs DR and Franco M** (2009). Fast-food consumption, diet quality, and neighborhood exposure to fast food: the multi-ethnic study of atherosclerosis. *American journal of epidemiology*, **170**(1), 29-36.
- Naiman DI, Leatherdale ST, Gotay C and Mâsse LC** (2015). School factors associated with the provision of physical education and levels of physical activity among elementary school students in Ontario. *Can J Public Health*, **106**(5), 290-296.
- Ng N, Santosa A and Kowal P** (2017). Sitting time and obesity among older adults in low- and middle-income countriesNawi Ng. *European Journal of Public Health*, **27**(suppl_3), Doi: 10.1093/eurpub/ckx186.042
- Nielsen L, Danielsen K and Sørensen T** (2011). Short sleep duration as a possible cause of obesity: critical analysis of the epidemiological evidence. *Obesity reviews*, **12**(2), 78-92.
- Ogden CL, Carroll MD and Lawman HG** (2016). Trends in obesity prevalence among children and adolescents in the united states, 1988-1994 through 2013-2014. *JAMA*, **315**(21), 2292-2299.
- Onis Md, Onyango AW, Borghi E, Siyam A, Nishida C and Siekmann J** (2007). Development of a WHO growth reference for school-aged children and adolescents. *Bulletin of the World health Organization*, **85**(9), 660-667.
- Peng W, Goldsmith R and Berry EM** (2017). Demographic and lifestyle factors associated with adherence to the Mediterranean diet in relation to overweight/obesity among Israeli adolescents: findings from the Mabat Israeli national youth health and nutrition survey. *Public health nutrition*, **20**(5), 883-892.
- Rautiainen S, Wang L, Lee IM, Manson JE, Buring JE and Sesso HD** (2015). Higher Intake of Fruit, but Not Vegetables or Fiber, at Baseline Is Associated with Lower Risk of Becoming Overweight or Obese in Middle-Aged and Older Women of Normal BMI at Baseline. *The Journal of Nutrition*, **145**(5), 960-968.
- Reue K** (2017). Sex differences in obesity: X chromosome dosage as a risk factor for increased food intake, adiposity and co-morbidities. *Physiology & Behavior*, **176**, 174-182.

Robaina KA and Martin KS (2013). Food insecurity, poor diet quality, and obesity among food pantry participants in Hartford, CT. *Journal of nutrition education and behavior*, **45**(2), 159-164.

Rohde JF, Larsen SC, Ängquist L, Olsen NJ, Stougaard M, Mortensen EL et al. (2017). Effects of the Healthy Start randomized intervention on dietary intake among obesity-prone normal-weight children. *Public health nutrition*, **20**(16), 2988-2997.

Rosenheck R (2008). Fast food consumption and increased caloric intake: a systematic review of a trajectory towards weight gain and obesity risk. *Obesity reviews*, **9**(6), 535-547.

Sanjeevi N, Freeland-Graves J and Hersh M (2018). Food insecurity, diet quality and body mass index of women participating in the Supplemental Nutrition Assistance Program: The role of intrapersonal, home environment, community and social factors. *Appetite*, **125**, 109-117.

Shafique S, Akhter N, Stallkamp G, de Pee S, Panagides D and Bloem MW (2007). Trends of under-and overweight among rural and urban poor women indicate the double burden of malnutrition in Bangladesh. *International journal of epidemiology*, **36**(2), 449-457.

Singh GM, Danaei G, Farzadfar F, Stevens GA, Woodward M, Wormser D et al. (2013). The Age-Specific Quantitative Effects of Metabolic Risk Factors on Cardiovascular Diseases and Diabetes: A Pooled Analysis. *PLOS ONE*, **8**(7), e65174.

Virtanen M, Kivimäki H, Ervasti J, Oksanen T, Pentti J, Kouvonen A et al. (2015). Fast-food outlets and grocery stores near school and adolescents' eating habits and overweight in Finland. *European Journal of Public Health*, **25**(4), 650-655.

WHO (2012). Global recommendations on physical activity for health. *World Health Organization*. WHO, Geneva, Switzerland.

WHO (2015). BMI-for-age (5-19 years). *World Health Organization*. WHO, Geneva, Switzerland. URL: http://www.who.int/growthref/who2007_bmi_for_age/en/ (accessed 12th May, 2018)

WHO (2018). Obesity and overweight. *World Health Organization*, WHO, Geneva, Switzerland. URL: <http://www.who.int/mediacentre/factsheets/fs311/en/> (accessed 15th May, 2018)

WHO and CDC (2017). URL: <http://www.cdc.gov/GSHS/>; <http://www.who.int/chp/gshs/en/>

Young DR, Koebnick C and Hsu JWY (2017). Sociodemographic associations of 4-year overweight and obese incidence among a racially diverse cohort of healthy weight 18-year-olds. *Pediatric Obesity*, **12**(6), 502-510.

Table 1. The complete list of explanatory variables.

Variables	Category	Measurements	Duration	Variable Type
How often went hungry	1 = Never		During the past 30 days	Food insecurity
	2 = Rarely			
	3 = Most of the time			
Fruit eating	1 = Low	less than one time per day	During the past 30 days	Food habits
Vegetable eating	2 = Average	1-2 times per day		
Soft drinks	3 = High	More than 2 times per day		
Fast food eating				
Sleep disturbance	1 = Never		During the past 30 days	Depression
	2 = Often			
Physical activity	1 = Moderate	≤2 days	60 Minutes per day	Physical activity
	2 = Vigorous	> 2 days	During the past 7 days	
Walk or bike to school	1 = Never	0 days	During the past 7 days	
Physical education attendance	2 = occasionally	1-3 days		
	3 = Regularly	> 3 days		
Sitting activities (includes TV watching, playing computer games etc.)	1 = Moderate	< 5 hours per day		
	2 = High	≥ 5 hours per day		

Table 2. The characteristics of study population.

Characteristics	n	Minimum	Maximum	Mean (SD)
Age (in years)	2980	11	17	14.2 (±0.98)
Height (m)	2703	1.27	1.9	1.563 (±0.087)
Weight (kg)	2703	28	102	45.88 (±7.868)
BMI	2703	13.05	39.67	18.78 (±2.87)
Gender	n	Overweight and obese (%)		
Male	1192	9.9%		
Female	1788	7.4%		

Note: All percentages are weighted.

Table 3. Percentage distribution of overweight/obesity among school-aged adolescents in Bangladesh according to their food habits and physical activities.

Risk Factors	Overweight or obese					
	Male		χ^2_{cal} (<i>p-value</i>)	Female		χ^2_{cal} (<i>p-value</i>)
	No (%)	Yes (%)		No (%)	Yes (%)	
How often went hungry:						
Never	333(93.8%)	22(6.2%)	15.419	573(93.2%)	42(6.8%)	8.840
Sometimes	507(92.5%)	41(7.5%)	(0.000)	728(94.4%)	43(5.6%)	(0.012)
Most of the time	75(81.5%)	17(18.5%)		226(89.0%)	28(11.0%)	
Fruit eating:						
Low	440(91.3%)	42(8.7%)	8.720	754(93.2%)	55(6.8%)	11.295
Average	322(90.2%)	35(9.8%)	(0.013)	561(91.1%)	55(8.9%)	(0.004)
High	160(97.6%)	5(2.4%)		1552(97.5%)	6(2.5%)	
Vegetable eating:						
Low	218(90.1%)	24(9.9%)	5.656	319(88.1%)	43(11.9%)	18.840
Average	318(90.4%)	41(9.6%)	(0.05)	666(93.5%)	46(6.5%)	(<0.0001)
High	318(94.6%)	18(5.4%)		561(95.4%)	27(4.6%)	

Soft drinks:						
Low	488(94.6%)	28(5.4%)	28.429	842(93.0%)	63(7.0%)	13.590
Average	299(92.6%)	24(7.4%)	(<0.0001)	515(95.2%)	26(4.8%)	(0.001)
High	126(81.3%)	29(18.7%)		183(87.6%)	26(12.4%)	
Fast food eating:						
Low	533(94.2%)	33(5.8%)	42.451	1009(92.7%)	79(7.3%)	9.912
Average	314(92.6%)	25(7.4%)	(<0.0001)	409(95.8%)	18(4.2%)	(0.007)
High	74(74.7%)	25(25.3%)		131(88.5%)	17(11.5%)	
Sleep disturbance:						
Never	356(88.6%)	46(11.4%)	9.502	645(90.8%)	65(9.2%)	9.072
Often	577(94.0%)	37(6.0%)	(0.002)	917(94.6%)	52(5.4%)	(0.003)
Physical activity:						
Moderate	281(87.0%)	42(13.0%)	14.509	567(90.4%)	60(9.6%)	13.772
Vigorous	635(94.1%)	40(5.9%)	(<0.0001)	959(95.1%)	49(4.9%)	(<0.0001)
Walk or bike to school:						
Never	175(81.8%)	39(18.2%)	35.484	486(89.3%)	58(10.7%)	19.068
Occasionally	181(94.3%)	11(5.7%)	(<0.0001)	161(94.7%)	9(5.3%)	(<0.0001)
Regularly	564(94.5%)	33(5.5%)		886(95.2%)	45(4.8%)	
PE attendance:						
Never	61(80.3%)	15(19.7%)	16.001	106(86.9%)	16(13.1%)	8.673
Occasionally	401(93.9%)	26(6.1%)	(<0.0001)	783(94.1%)	49(5.9%)	(0.013)
Regularly	436(91.8%)	39(8.2%)		615(92.6%)	49(7.4%)	
Sitting activities:						
Moderate	831(92.4%)	68(7.6%)	12.567	1451(93.1%)	107(6.9%)	0.000
High	63(80.8%)	15(19.2%)	(<0.0001)	68(93.2%)	5(6.8%)	(0.995)

Note: Significant at $p < 0.05$ level; PE, Physical Education

Table 4. Logistic regression analysis to estimate the effect of adolescents' food habits and physical activities on the state of overweight/obesity in Bangladesh, 2014.

Risk Factors	Overweight or obese	
	AOR	95% CI
Gender		
Male ^(RC)	1.00
Female	0.573**	(0.403-0.816)
How often went hungry:		
Never ^(RC)	1.00
Sometimes	1.475*	(1.001-2.175)
Most of the time	2.789**	(1.733-4.489)
Fruit eating:		
Low ^(RC)	1.00
Average	1.070	(0.749-1.530)
High	0.454*	(0.205-0.997)
Vegetable eating:		
Low ^(RC)	1.00
Average	0.625*	(0.420-0.930)
High	0.475**	(0.294-0.768)
Soft drinks:		
Low ^(RC)	1.00
Average	1.244	(0.814-1.900)
High	2.357**	(1.544-3.597)
Fast food eating:		
Low ^(RC)	1.00
Average	1.244	(0.804-1.923)
High	2.777**	(1.755-4.392)

Sleep disturbance:		
Never ^(RC)	1.00
Often	0.675*	(0.481-0.947)
Physical activity:		
Moderate ^(RC)	1.00
Vigorous	0.850	(0.597-1.209)
Walk or bike to school:		
Never ^(RC)	1.00
Occasionally	0.334**	(0.185-0.601)
Regularly	0.472**	(0.327-0.682)
PE attendance:		
Never ^(RC)	1.00
Occasionally	0.483**	(0.284-0.822)
Regularly	0.592*	(0.327-0.682)
Sitting activities:		
Moderate ^(RC)	1.00
High	1.551	(0.860-2.797)
Model summary:		
Model Chi-Square (p-value)	2220.687 (<0.001)	
-2Log Likelihood	1048.196	
Cox and Snell R-square	0.610	
Nagelkerke R-square	0.813	

Note: Sample are weighted and controlled by age. “(RC)” denotes reference category; AOR, adjusted odds ratio; Significant at ** $p < 0.01$, and * $p < 0.05$ level.

Table 5. Logistic regression analysis to estimate the gender differential effect of food habits on adolescents' overweight/obesity in Bangladesh, 2014.

Risk Factors	Male		Female	
	AOR	95% CI	AOR	95% CI
How often went hungry:				
Never ^(RC)	1.00	1.00
Sometimes	1.399*	(1.036-1.891)	1.235	(0.791-1.928)
Most of the time	2.759**	(1.846-4.125)	1.540	(0.904-2.623)
Fruit eating:				
Low ^(RC)	1.00	1.00
Average	1.201	(0.901-1.601)	1.163	(0.763-1.773)
High	0.372**	(0.203-0.683)	0.645	(0.258-1.613)
Vegetable eating:				
Low ^(RC)	1.00	1.00
Average	1.043	(0.753-1.446)	0.582*	(0.372-0.910)
High	0.504**	(0.333-0.764)	0.432**	(0.248-0.753)
Soft drinks:				
Low ^(RC)	1.00	1.00
Average	2.583**	(1.855-3.597)	0.785	(0.470-1.312)
High	2.929**	(2.086-4.112)	1.677*	(1.022-2.753)
Fast food eating:				
Low ^(RC)	1.00	1.00
Average	1.503*	(1.084-2.083)	0.737	(0.411-1.321)
High	6.064**	(4.327-8.499)	1.695*	(1.011-3.174)
Model summary:				
Model Chi-Square (p-value)	2327.11 (<0.001)		1401.33 (<0.001)	
-2Log Likelihood	1596.663		781.801	
Cox and Snell R-square	0.561		0.589	
Nagelkerke R-square	0.747		0.786	

Table 6. Logistic regression analysis to estimate the gender differential effect of physical activities on adolescents’ overweight/obesity in Bangladesh, 2014.

Risk Factors	Male		Female	
	AOR	95% CI	AOR	95% CI
Sleep disturbance:				
Never ^(RC)	1.00	1.00
Often	0.590**	(0.455-0.766)	0.555**	(0.369-0.837)
Physical activity:				
Moderate ^(RC)	1.00	1.00
Vigorous	0.751*	(0.562-0.991)	0.809	(0.532-1.229)
Walk or bike to school:				
Never ^(RC)	1.00	1.00
Occasionally	0.265**	(0.171-0.410)	0.453*	(0.205-0.924)
Regularly	0.430**	(0.322-0.576)	0.557**	(0.359-0.866)
PE attendance:				
Never ^(RC)	1.00	1.00
Occasionally	0.420**	(0.281-0.627)	0.445**	(0.266-0.745)
Regularly	0.488**	(0.330-0.722)	0.754	(0.461-1.234)
Sitting activities:				
Moderate ^(RC)	1.00	1.00
High	3.404**	(2.343-4.945)	0.469	(0.154-1.427)
Model summary:				
Model Chi-Square (p-value)	2102.016 (<0.001)		1403.618 (<0.001)	
-2Log Likelihood	1638.223		718.459	
Cox and Snell R-square	0.541		0.600	
Nagelkerke R-square	0.772		0.800	

Note: Sample are weighted and controlled by age. “(RC)” denotes reference category; AOR, adjusted odds ratio; Significant at ** $p<0.01$, and * $p<0.05$ level.