



UWL REPOSITORY

repository.uwl.ac.uk

Exploring the determinants of global life expectancy in an ecological perspective

Mondal, Nazrul Islam, Khan, Nuruzzaman, Baki, Abu Naser Muhammad Abdul, Hoque, Nazrul and Khan, Hafiz T.A. ORCID: <https://orcid.org/0000-0002-1817-3730> (2019) Exploring the determinants of global life expectancy in an ecological perspective. Turkish Journal of Public Health, 17 (3). pp. 314-325. ISSN 1304-1096

10.20518/tjph.452721

This is the Published Version of the final output.

UWL repository link: <https://repository.uwl.ac.uk/id/eprint/6487/>

Alternative formats: If you require this document in an alternative format, please contact: open.research@uwl.ac.uk

Copyright: Creative Commons: Attribution-Noncommercial 4.0

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy: If you believe that this document breaches copyright, please contact us at open.research@uwl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Research article

Exploring the determinants of global life expectancy from an ecological perspective

Md. Nazrul Islam Mondal^{a,b}, Abu Naser Muhammad Abdul Bakic, Md. Nazrul Hoque^d, Hafiz T.A. Khan^e, Md. Nuruzzaman Khan^f

^aProfessor, Department of Population Science and Human Resource Development, University of Rajshahi Rajshahi; Bangladesh

^bVisiting Scholar, Department of Sociology, Middle East Technical University, Ankara; Turkey

^cPh. D student, Department of Population Science and Human Resource Development, University of Rajshahi Rajshahi; Bangladesh

^dHobby Center for Public Policy, University of Houston, 306 McElhinney Hall, Houston, TX 77204-5021; USA

^eProfessor, The Graduate School, University of West London, St Mary's Road, Ealing, London W5 5RF; UK

^fDepartment of Population Science, Jatiya Kabi Kazi Nazrul Islam University, Mymensingh 2200; Bangladesh

Submitted date: 10.08.2018, Accepted date: 06.07.2019


Abstract

Objective: This study is designed to understand the impacts of demographic events, socioeconomic differentials, health factors' availability, and environmental reasons which influence life expectancy (LE) globally. **Methods:** Data of 183 countries were taken from the United Nations agencies. The predicted variable was LE, and the determinants were demographic events, socioeconomic factors, health-related factors, and environmental issues. Descriptive statistics, correlation analysis, and backward multiple regression analysis were used to reach the research objectives. **Results:** The lowest LEs were found in the African countries; and LE was found significantly associated ($p < 0.05$) with a wide range of demographic, socioeconomic, health, and environmental factors. The necessity of full coverage of immunization, higher income, and improved sanitation are expected to raise LE. However, LE may be increased by way of decreased fertilities, Human Immunodeficiency Virus (HIV) prevalence, and carbon dioxide (CO_2) emissions. **Conclusions:** LE is influenced by different demographic, socioeconomic, health, and environmental factors. Country-level and global efforts should be taken to raise LE throughout the reduction of HIV infection, births, and CO_2 emissions. The policy-makers should focus on advancing reproductive decisions, increasing immunization coverage, and upturning improved sanitation usage.

Keywords: Life expectancy; determinants of life expectancy; ecological analysis; HIV infection

Corresponding Author: Md. Nazrul Islam Mondal, Department of Population Science and Human Resource Development, University of Rajshahi, Rajshahi 6205; Bangladesh. Phone: +88-01716922856 E-mail: nazrulupm@gmail.com

Copyright holder Turkish Journal of Public Health

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.  This is an open Access article which can be used if cited properly

Küresel doğumda beklenen yaşam beklentisinin ekolojik perspektifle incelenmesi

Öz

Amaç: Bu çalışma, yaşam beklentisini (YB) küresel olarak etkileyen demografik olayların, sosyoekonomik farklılıkların, sağlık hizmetlerinin mevcudiyetinin ve çevresel nedenlerin etkilerini anlamak için tasarlanmıştır. **Yöntem:** 183 ülkenin verileri Birleşmiş Milletler ajanslarından alınmıştır. Tahmin edilen (bağımlı) değişken YB'dir; belirleyiciler ise demografik olaylar, sosyoekonomik faktörler, sağlıkla ilgili faktörler ve çevresel konulardır. Araştırma amaçlarını değerlendirmek için, tanımlayıcı istatistikler, korelasyon analizi ve multiple regresyon analizleri kullanılmıştır. **Bulgular:** En düşük YB Afrika ülkelerine ait olup, çok çeşitli demografik, sosyoekonomik, sağlık ve çevresel faktörlerle anlamlı düzeyde ilişkilidir ($p < 0.05$). Tam kapsayıcı bağışıklama, daha yüksek gelir ve iyileştirilmiş sanitasyonun YB'yi yükseltmesi beklenmektedir. Bununla birlikte, doğurganlık, HIV yaygınlığının ve karbondioksit emisyonlarının azaltılması yoluyla YB artırılabilir. **Sonuç:** YB farklı demografik, sosyoekonomik, sağlık ve çevresel faktörlerden etkilenmektedir. HIV enfeksiyonu, doğumlar ve CO_2 emisyonlarının azaltılması yoluyla YB'yi yükseltmek için ülke düzeyinde ve küresel olarak çaba gösterilmelidir. Politika yapıcılar üreme kararlarını iletmeye, aşı kapsamını artırmaya ve sanitasyonu iyileştirmeye odaklanmalıdır.

Anahtar sözcükler: Yaşam beklentisi; yaşam beklentisinin belirleyicileri, ekolojik analiz, HIV enfeksiyonu

Introduction

Life expectancy (LE) at birth reflects the overall mortality level of a population. It is defined as “the years a newborn infant would live if prevailing patterns of age-specific mortality rates at the time of birth were to stay the same throughout the child’s life”.¹ The LE is considered as a commonly used indicator to measure the overall improvement of a country. It is of actual importance for the low- and lower middle-income countries and especially for the African countries since these countries are struggling to reach socioeconomic advancements by significant investments in the social and health areas. In most countries of the world, LEs have been increasing during the last decade. In 2015, the global LE was 71.40 years (female, 73.80 years; male, 69.10 years), extending from 60 years in the World Health Organization (WHO) African region to 76.80 years in the WHO European region, giving a ratio of 1.30 between the two regions. Globally, it is observed that females live

longer than males. In 1990, the difference in LEs between the sexes was observed to be 4.50 years and it had remained almost the same by 2015 (4.60 years). Optimistically, the global average LE increased by 5 years between the years 2000 and 2015, the fastest rise since the 1960s. From 2000 to 2015 the increase of LE was observed greatest in the WHO African region, where LE raised by 9.40 years to 60 years, mostly dominated by the developments in child survival and extended access to antiretroviral therapy of Human Immunodeficiency Virus (HIV).¹

An ecologic study focuses on the comparison of groups, rather than individuals which are often used to measure the prevalence and incidence of diseases. In an ecologic analysis, the variables may be aggregate measures, environmental measures, or global measures. It is inexpensive and easy to carry out, using routinely collected data, but they are prone to bias and confounding.² The contributing factors in increasing LE are one of the principal interests among demographic and health researchers. Today, people across the

globe enjoy better health, are economically solvent, and live longer than ever before. The LE is expected to increase by seven years for the period of 1998-2025, and it would be greater than 80 years in 26 countries.³ Until now, the mentionable disparities in LE were observed between high- and low-income countries. The significant determinants were the development of technology, drugs, environmental development, and international cooperation. Moreover, the rise in LE was initiated by higher income and education, access to improved sanitation, and clean water; developed healthcare facilities; and enormous rises in agricultural production. The changeability of LE has significant effects on personal and combined human activities since they affect fertility performance, socioeconomic development, human capital investment, intergenerational transmissions, and encouragements for pension subsidy rights.⁴⁻⁵ LE focuses the population's physical condition of a country and the healthcare services that people generally receive when they fall ill.⁶ The demographic and economic factors of LE include sex, age, residence, schooling, and income [gross national income (GNI) per capita].⁷⁻⁹ Research also shows that income plays an important role for increasing recent LE. For example, enhanced income contributed an affirmative influence on LE in South Korea.¹⁰ Older people in Thailand with improved earnings and innovative education had better health outcomes.¹¹ In recent times, economic solvency and schooling inequalities were observed to explain the regional dissimilarities in LE with other health indicators.¹² The lower earnings and joblessness were seen to undesirably affect health outcomes.¹³ Significant associations were observed between LE and education in Brazil,¹⁴ Finland, Sweden, Norway, Denmark,¹⁵⁻¹⁶ and some European countries.¹⁷ Again, LE was found to be significantly associated with lower infant mortality rates and higher schooling rates.¹⁸ When and where death risks are higher, the women usually give birth more often to raise the possibility that at any rate, some offspring will stay alive to adulthood.¹⁹ The contributing factors of LE which were taken into account were: healthcare expenses and

resources, death rates, HIV prevalence, and health consequences.²⁰ The services related to healthcare, e.g., the increased numbers of healthcare persons, hospital supplies, and prenatal checkups diminished deaths and raised LE.²¹ Moreover, some researchers have pointed at the effects of demographic measures, socioeconomic variability, and accessibility of healthcare resources on LE.²²⁻²⁴ The previous studies on global inequalities in LE have usually explored the overall mortality. Until recently no study has ever investigated on a global scale, considering demographic, socioeconomic, health, and environmental factors together. This study attempts to fill in this research gap. Therefore, the main objective of this study is to explore the impacts of demographic events, socioeconomic differentials, health factors' availability, and environmental reasons that influence LE globally. It is believed that the study will help understanding the factors that have significant effects on LE globally.

Method

Data and necessary information of 183 WHO member countries were taken from the different United Nations (UN) agencies.^{1, 25-27} Due to the inavailability of LE data; some countries were not included in this study. The factors that had significant effects on LE in earlier researches were considered.^{8-10, 22-23, 28} The factors considered in this study, along with their definitions and sources are shown in Table 1. The contributing factors of LE (Y) are classified into 4 main groups: demographic events, social and economic status, health factors, and environmental issues. The demographic events were the total fertility rate (TFR) (X_1), adolescent birth rate (ABR) (X_2), and population density (X_3); social and economic determinants were mean years of schooling (X_9), and GNI (X_7); health-related variables were HIV prevalence rate (X_5), physicians density (X_4), and immunization coverage rate (X_6); and the environmental issues were improved sanitation usage rate (X_8) and carbon dioxide (CO_2) emissions rate (X_{10}).

Table 1: Variables, their descriptions and sources

Selected Variables	Descriptions
Life expectancy	The average number of years a newborn infant can expect to live under current mortality levels.
Total fertility rate (TFR)	The average number of children a woman would have assuming that current age-specific birth rates remain constant throughout her childbearing years (usually considered to age 15 to 49).
Adolescent fertility rate (AFR)	The number of births to women ages 15-19 per 1000 women ages 15-19.
Human Immunodeficiency Virus (HIV) prevalence rate	Estimated number of adult population aged 15-49 years per 100 populations.
Physicians density	The number of physicians per ten thousand populations.
Gross National Income (GNI) per capita	GNI PPP per capita is gross national income in purchasing power parity (PPP) divided by midyear population. GNI PPP refers to gross national income converted to "international" dollars using a purchasing power parity conversion factor. International dollars indicate the amount of goods and services one could buy in the USA with a given amount of money.
Mean year of schooling	An average number of years of education received by people 25 and older, converted from education attainment levels using official durations of each level.
Population density	Population density is a measurement of population per unit area or unit volume. It was obtained by the mid-year 2016 population divided by the square kilometers of arable land.
Diabetes prevalence rate	The diabetes prevalence rate is the number of people per 100 in a given group or population who are reported to have diabetes.
Immunization rate	The number of children per 100 ages 12-30 months takes vaccines for Diphtheria, Tetanus, and Pertussis (DTP)
Improved sanitation rate	The number of population per 100 using Improved sanitation
Carbon dioxide (CO ₂) emissions	Total carbon emissions from fossil-fuel burning, cement production, and gas flaring. Emissions are expressed in million metric tons of carbon.

Source: (WHO, 2016)

To analyze the collected data, descriptive statistics, correlation analysis, and stepwise (backward) regression analysis were used. Descriptive statistics were used to see the maximum, minimum, mean, median values, and standard deviations (SDs). The Pearson correlation analysis was used to set up the relationships among variables. Backward stepwise regression is a stepwise regression approach, which begins with a full (saturated) model and at each step gradually eliminates variables from the regression model to find a reduced model that best explains the data. The stepwise approach is useful because it reduces the number of predictors, reducing the multicollinearity problem and it is one of the ways to resolve the over fitting. Assume the regression model considering all the covariates is:

$$Y = \beta_0 + \sum_i \beta_i X_i + \varepsilon. \quad (1)$$

In Eq. (1), Y is the outcome variable (LE), X_i ($i = 1, 2, 3, \dots, 10$) are the explanatory variables, β_0 is the constant, β_i ($i = 1, 2, 3, \dots, 10$) are the unknown regression coefficients, and ε is the error term with an $N(0, \sigma^2)$ distribution. To check the multicollinearity problem among the predictors, the variance inflation factor (VIF) was used. The VIF for the explanatory variables, X_j is:

$$VIF_j = (1 - R_j^2)^{-1}, \quad j = 1, 2, 3, \dots, l. \quad (2)$$

In Eq. (2), l is the number of predictors, and R_j^2 is the square of the multiple correlation coefficient of the j th variable with the remaining $(l-1)$ variables. The VIF is always positive and if it is less than 5, then

there is no multicollinearity; if it is less than 10, then there exists a reasonable multicollinearity; and if it is greater than 10, there exists considerable multicollinearity among the variables.²⁹ The analyses were performed by using the Statistical Package for Social Sciences (SPSS) (SPSS Inc, Chicago, IL; USA).

Results

The study included a total of 183 WHO member countries. The influences of TFR, ABR, population density, schooling, GNI,

physicians' density, immunization rate, HIV prevalence rate, improved sanitation, and CO₂ emissions on LE are explored here sequentially.

The descriptive statistics of dependent and independent variables are presented in Table 2. The extreme values (maximum and minimum) of all the factors as well as the variables' mean, median, and SDs are presented in this table to explore their main features. This type of analysis is considered very meaningful since the diverse factors are frequently calculated in diverse units.

Table 2. Descriptive statistics for dependent and independent variables from 194 countries

Selected Variables	N	Minimum	Maximum	Mean	SE mean	Median	SD
Life expectancy	183	50.1 (Sierra Leone)	83.7(Japan)	71.29	0.59	73.50	7.98
Total fertility rate	182	1(Republic of Korea)	8(Niger)	2.85	0.105	2.31	1.41
Adolescent birth rate	186	1(Democratic People's Republic of Korea)	229(Central African Republic)	55.54	3.521	44.00	48.02
Population density	189	2(Mongolia)	18866(Monaco)	292.77	108.15	82.53	1486.79
Physicians density	161	0(Liberia)	77(Qatar)	17.44	1.25	14.20	15.81
HIV prevalence rate	111	0(Azerbaijan)	28(Swaziland)	2.46	0.48	60.00	5.10
Immunization rate	194	24(Equatorial Guinea)	99(Qatar)	88.36	0.97	94.00	13.40
GNI per capita	163	600(Central African Republic)	140720(Qatar)	18604.91	1549.92	11960.00	19788.08
Improved sanitation	179	7(South Sudan)	100(Singapore)	72.59	2.191	85.60	29.31
Mean years of schooling	157	1(Niger)	14(Finland)	8.59	0.246	9.20	3.08
CO ₂ emissions	183	0(Burundi)	44(Qatar)	4.65	0.458	2.36	6.26

Note: 'N, number of countries', 'SE, standard error', 'SD, standard deviation'

Table 3. Pearson correlation coefficient between variables

	Y	X ₁	X ₂	X ₃	X ₄	X ₅	X ₆	X ₇	X ₈	X ₉	X ₁₀	X ₁₁
Y	1	-.839**	-.783**	.173*	.688**	-.397**	.347**	.586**	.638**	.849**	.745**	.447**
X ₁		1	.782**	-.142	-.656**	.175	-.366**	-.556**	-.511**	-.819**	-.734**	-.427**
X ₂			1	-.138	-.611**	.219*	-.273**	-.591**	-.557**	-.760**	-.712**	-.449**
X ₃				1	.259**	-.120	.008	.096	.259**	.113	.066	.033
X ₄					1	-.282**	.065	.366**	.694**	.674**	.689**	.494**
X ₅						1	-.233*	-.022	-.135	-.273**	-.028	-.084
X ₆							1	.148*	.322**	.385**	.238**	.371**
X ₇								1	.342**	.535**	.470**	.281**
X ₈									1	.582**	.587**	.790**
X ₉										1	.785**	.501**
X ₁₀											1	.473**
X ₁₁												1

Note: * $p < 0.05$, ** $p < 0.01$, 'Y, Life expectancy', 'X₁, Total fertility rate', 'X₂, Adolescent birth rate', 'X₃, Population density', 'X₄, Physicians density', 'X₅, HIV prevalence rate', 'X₆, Diabetes prevalence rate', 'X₇, Immunization', 'X₈, GNI per capita', 'X₉, Improved sanitation', 'X₁₀, Mean years of schooling', 'X₁₁, CO₂ emissions'

The LEs of the African countries are found remarkably lower than that of other countries in the globe. Among these countries, the LE of Sierra Leone is found the lowest (50.1 years). The LEs are found 52.4, 53.5, 53.1, 53.3, and 53.7 years of Angola, Central African Republic, Chad, Côte d'Ivoire, and Lesotho respectively; which are around three-fifths compared to countries like Japan (83.7 years), Switzerland (83.4 years), Singapore (83.1 years), etc. The TFR and ABR are found the highest in the African countries. The TFR of Nigeria is 8, in Somalia, Mali, Chad, Angola, the Democratic Republic of the Congo, Burundi, Uganda, Nigeria, and Burkina Faso it is 6. The average TFR is found 1.70 in the more developed countries. A comparable trend is found for ABR. The highest ABR (229) is observed in the Central African Republic. It is the second highest in Niger (206) followed by Chad (203), Angola (191), and some other African countries. The highest densely populated countries are Singapore (7,829 population/km² (pop./ km²), Bahrain (1,789 pop./km²), Maldives (1,364 pop. /km²), Malta (1,348 pop. /km²), Bangladesh (1,237 pop./km²), etc. On the other hand, a good number of the lowest population density countries are found, e.g., Mongolia (2 pop. /km²), Namibia (3 pop/km²), Australia (3 pop./km²), Iceland (3 pop./km²), Suriname (3 pop. /km²), etc. The physicians' density is observed very low in the countries of the African region. It is mentionable that there are only 1 physician per 10,000 people in the Central African Republic, Togo, Burkina Faso, etc.; 2 physicians per 10,000 people in Madagascar, Angola, Zambia, etc. The HIV prevalence rates are observed the maximum in the countries of the African region (Swaziland 28.7%; Botswana 22.3%; Lesotho 22.8%; South Africa 19.3%, Zimbabwe 14.7%; etc.) while it is <0.10% in the most developed countries. Immunization covers almost 100% children in the globe except for some countries e.g., Equatorial Guinea (24%), South Sudan (39%), Somalia (42%), etc. The economic factor, GNI was found very low of the countries where the LEs are found lower (e.g., GNI of Central African Republic is 600\$). The lower improved sanitation facilities were observed in the African countries (e.g., South

Sudan, 7%; Niger, 11%; Togo, 12%, etc.) where the LEs are found also lower. The mean years of schooling were found very few in the countries of the African region (Niger, 1 year; Mali, 2 years; etc.). The highest CO₂ emission rates were found in the middle-income countries e.g., Qatar (40.5), Trinidad and Tobago (34.5), Kuwait (27.3), etc.

In bivariate analysis, the results of Pearson correlation coefficients (*r*) are presented in Table 3. The results revealed that the LE was found significantly positively correlated with physician density, immunization, income, sanitation, schooling, and CO₂ emissions; and significantly negatively correlated with fertilities (TFR and ABR), and HIV prevalence rate.

The results of the stepwise regression analysis are presented in Table 4. In this analysis, five sets of regression models (Models 1-5) were performed. There were no multicollinearity among the variables because the VIFs were less than 5 for all the cases except TFR (VIF = 5.116) and improved sanitation using rate (VIF = 5.858). Model 1 ($R_a^2 = 0.837$) considered all the explanatory variables, among these predictors TFR, HIV prevalence, and CO₂ emissions indicated significant negative effects; and income and sanitation indicated significant positive effects on LE. Model 2 ($R_a^2 = 0.840$) retained the predictors except population density where TFR, HIV prevalence rate, income, sanitation, and CO₂ emissions were identified as the significant factors. Model 3 ($R_a^2 = 0.843$) retained all the predictors except population density and schooling where TFR, HIV prevalence rate, GNI per capita, sanitation, and CO₂ emissions were identified as the significant factors. Model 4 ($R_a^2 = 0.845$) excluded ABR, population density, and mean years of schooling; and among the retained predictors TFR, HIV prevalence, GNI per capita, sanitation, and CO₂ emissions were identified as the significant predictors of LE. Model 5 ($R_a^2 = 0.845$) excluded the predictors ABR, population density, physician density, mean years of schooling; and among the retained predictors TFR, HIV prevalence, immunization, GNI per capita, sanitation, and CO₂ emissions were identified as the significant factors of LE.

Table 4: Results of backward regression analysis

Model 1			Model 2		Model 3		Model 4		Model 5		Model 6		Model 7	
	β	VIF	β	VIF	β	VIF	β	VIF	β	VIF	β	VIF	β	VIF
X ₁	-.297**	5.335	-.299**	4.888	.295**	4.699	.305**	4.476	.325**	3.971	.321**	3.959	.329**	3.932
X ₂	-.051	3.968	-.051	3.967	-.052	3.958	-.055	3.937						
X ₃	-.011	1.230	-.012	1.162										
X ₄	-.081	4.054	-.079	3.694	-.076	3.596	-.063	3.275	-.063	3.274				
X ₅	-.209**	1.529	-.207**	1.262	-.206**	1.243	-.209**	1.226	-.213**	1.200	-.205**	1.151	-.209**	1.146
X ₆	.057	2.113	.057	2.113	.058	2.108	.059	2.106	.050	1.987	.065	1.793		
X ₇	.108	2.011	.110	1.889	.108	1.860	.103	1.816	.116*	1.580	.122*	1.550	.117*	1.536
X ₈	.361**	3.710	.362**	3.664	.364**	3.608	.383**	2.865	.385**	2.860	.370**	2.670	.357**	2.592
X ₉	.301*	7.711	.302*	7.672	.304*	7.650	.306*	7.643	.322**	7.291	.283**	6.003	.329**	5.087
X ₁₀	.008	4.930												
X ₁₁	-.172*	3.217	-.171*	3.170	-.171*	3.170	-.184*	2.840	-.175*	2.742	-.184*	2.685	-.186*	2.682
R _a ²	0.834		0.837		0.840		0.842		0.844		0.845		0.845	

Note: * $p < 0.05$, ** $p < 0.01$, ' β , Standardized Coefficients,' 'VIF, variance inflation factor', 'X₁, Total fertility rate', 'X₂, Adolescent birth rate', 'X₃, Population density', 'X₄, Physicians density', 'X₅, Prevalence of HIV rate', 'X₆, Diabetes prevalence rate', 'X₇, Immunization', 'X₈, GNI per capita', 'X₉, Improved sanitation', 'X₁₀, Mean years of schooling', 'X₁₁, CO₂ emission', ' R_a^2 , adjusted R squared'

Discussion

The univariate analysis clearly explains the global scenario of LE. The correlation analysis established the significant relationships between LE and demographic, socioeconomic, health, and environmental factors. Finally, the regression model (Model 5) identified TFR, HIV prevalence, income, sanitation, and CO₂ emissions as the significant predictors of global LE.

In 2015, the global LE was 71.40 years and it is improved at a rate of more than 3 years per 10 years from 1950. In the 1990s, the growth on LE stalled in the countries of African region because of the tremendous increase of HIV prevalence rates; and increased death rates in many former Soviet countries following the collapse of the Soviet Union. In most of the regions, the LEs have increased from 2000 onwards. The overall global increment of LE was 5.00 years from 2000 to 2015. The largest increment (9.40 years) was observed in the African countries. Now, the LE is more than 82 years in 12 countries (Switzerland, Spain, Italy, Iceland,

Israel, France, Sweden, Japan, Singapore, Australia, the Republic of Korea, and Canada) and it is on an average 80 years in 29 countries. On the other hand, there are 22 countries with LEs less than 60 years.¹

Globally the fertility patterns have changed dramatically over the last few decades. In 2015, the global TFR was 2.50 which masks wide regional variations. Africa remains the region with the highest TFR (4.70) and Europe has the lowest TFR (1.60).³⁰ The higher TFR may also have a negative effect on LE. Clearly, the families having the higher number of births, the lower duration between births, and insufficient resources allotted for those children may reduce breastfeeding duration and risk malnutrition in those children. The consistent results are observed in this study. In addition, LE was inversely correlated with fertilities and from the regression analysis; the TFR is identified as the significant determinant of LE.

In the developing countries, the TFR has fallen remarkably since 1950. The higher

fertility ($TFR \geq 5$) and closely-spaced births characterized 33 countries among them 29 are in Sub-Saharan Africa. The higher fertility creates the higher health burden, which risks for infants and mothers, derails wealth investment, downturns economic growth, and worsens ecological coercions. The significant downturn in births may contribute to raising the productive output. The present study identified that higher fertility contributed to lower LE. Obviously, the declining trend of TFR assists to raise economic growth through the positive changes in age-structure. The ABR and coverage of modern contraceptives are two proposed indicators of the Sustainable Development Goal Target 3.7 on general entrance to reproductive healthcare services. It was estimated that the global ABR= 44 which was 5 times more prevalent in the low-income countries compared to the high-income countries.³⁰ Early sexual activities (marital or extramarital) contribute to ABR. In 2014, it was estimated globally 700 million girls were married before 18 years among them 250 million had experienced extramarital sexual relations.³¹ Globally, adolescents are exposed to excess reproductive health hazards. Adolescent fertility results in adverse mother and infant health outcomes and it is strongly associated with obstructed delivery, low birth weight, fetal growth retardation, and higher maternal and infant deaths.²⁸ The higher availability of unsafe sexual activities among unmarried girls threatens them to unexpected pregnancies, risky abortions, and sexually transmitted diseases. On the other hand, delayed motherhood beyond adolescence raises survivorship of offspring.³²

The availability of physicians contributes to raise LE. In an area or in a country, if we find an inadequate number of medical personnel that treats the general population, it would more likely not to have enough healthcare services. Availability and easy access to healthcare services are considered an important issue to protect from diseases and to hasten recovery from sickness and disabilities. In this study, the physician density is significantly associated with LE which is consistent with the previous studies.^{12, 20-21} On the other hand, an

inadequate number of healthcare personnel, lack of skilled health workers, and not having easy access to healthcare services may contribute to raise the disease prevalence.³³ Moreover, the physicians may contribute to increasing LE by reducing fertilities.⁸

The prevalence of HIV is found significantly negatively correlated with LE and in all the regression models (Models 1-5), it was retained as the significant predictor of LE. HIV leads to untreatable disease and hits the immune system. The HIV infected person may survive 9-11 years without treatment. This means that HIV-infected individuals have shorter lifetimes. The higher HIV prevalence in an area or a country may cause to increase more HIV infected persons. As a result, the higher prevalence of HIV infection lowers LE. Globally, the differences of LEs among regions are also seen, as the numerous countries in the African region with the higher prevalence of HIV experienced a fall in LE and established a relationship between higher prevalence of HIV and lower LE.³⁴

Immunization was found significantly positively correlated with LE. In all the regression models (Models 1-5) this factor retained as the significant determinant. Vaccination greatly reduces diseases, disabilities, deaths, and increases LE worldwide. It has significantly reduced the burden of infectious diseases. Today, vaccines are considered an excellent safety tool. In a previous study, it was identified that vaccination significantly contributes to increasing LE by lowering morbidity and mortality.³⁵

National income (GNI per capita) has shown significant positive correlation with LE. In regression analysis, GNI is identified as the significant determinant of LE in all the models (Models 1-5). The present study identified that financial development increases LE. The populations of a country live longer when they have quality livelihoods and have lower death rates.¹¹ The quality livelihood of a country's population is measured by that country's national income, which has also been established in this study. There is significant evidence linking income dissimilarity to unfortunate health outcomes. In the low- and lower middle-income

countries, a smaller amount of money is allotted for healthcare services which might contribute to having lower LE. The findings of this study authenticate the previous study results on the effects of national income on LE.^{8,13}

Improved sanitation was found significantly positively correlated with LE and identified as the significant determinant of LE (Models 1-5). An increase in the lifespan was determined mostly by the improvements in sanitation systems during the 1990s and early 2000s. The transmission of most infectious diseases (e.g., cholera, typhoid, infectious hepatitis, polio, cryptosporidiosis, ascariasis, etc.) is through human waste. Each year, around 1.80 million people die due to diarrheal diseases, among them 90% are less than 5 years old, and they are mostly in the low- and lower-income countries.³⁶ Usually, the unhealthy sanitation systems may contribute to having various infections. In the tropical and subtropical areas in terms of socioeconomic and public health concern, the ranks of malaria and schistosomiasis were placed first and second respectively among the human parasitic diseases. Globally, around 0.6 million deaths occur per year due to severe *Ascaris* infections.³⁷

Education is another influential factor found to be significantly positively correlated with LE. This finding has likewise important implications. An educated nation might contribute increasing LE. The higher schooling rates are strongly correlated with timely receipt of healthcare services and better healthcare knowledge. The educated population is more likely to have improved prenatal care and they can be promoted to optimize the use of mothers' healthcare services, thus keeping away from childbirth-related difficulties.³⁸ Usually, more educated populations can earn more money, which contributes to having higher household incomes, enabling them buy quality healthcare services. Besides these, the educated individuals tend to better understand the importance of personal hygiene, nutritional information, knowledge regarding illness, etc.³⁹

The CO_2 emission was significantly positively correlated with LE, and it was

retained all in the regression models (Models 1-5). The regression analysis identified it is a significant determinant of LE. The higher energy usage is essential for economic development, which is the prerequisite for human development. The higher energy usage leads to higher CO_2 emissions. The higher LE is companionable with low carbon emissions. From the perspective of a sustainable development, the countries would reach both higher GNI and higher LEs at low levels of CO_2 emissions.⁴⁰

A limitation of this study is that we analyzed data for the most common determining factors, *i.e.*, those which were found to be significantly associated with LE in the previous studies. In addition, the analysis was limited to those countries whose data were available. The present study identified associations and determinants of LE but did not investigate the determinants that explain gender and urban-rural differences in LE in these countries. Data on the 183 countries were obtained from the different UN agencies. However, the sources and quality of data vary according to country. Some countries have comprehensive civil registration and vital statistics systems and regular censuses of the entire population. However, many countries have incomplete or dysfunctional birth and death registration systems, and therefore lack continuous empirical data on mortality and LE.

Conclusion

The study analyzed how demographic, socioeconomic, health, and environmental factors affect LE globally, and empirically identified the determinants of LE. All the selected factors were found to be statistically correlated with LE. Among these factors, physician density, immunization coverage, national income, and improved sanitation were positively; but the higher fertilities (TFR and ABR), and HIV prevalence rate were negatively associated with LE. However, TFR, HIV prevalence rate, immunization rate, national income, improved sanitation, and CO_2 emissions were identified as influential factors of LE. The findings of this study have some important policy implications for all countries particularly for the African

countries. To raise LEs of those countries substantive steps should be taken. National and international programs may be taken to raise LE by raising schooling, immunization coverage, improved sanitation, and the number of physicians; and limiting fertilities and HIV infection in the countries where LEs are still very low. We analyzed cross-sectional data of 183 countries, of 10 demographic, socioeconomic, health, and environmental determinants. To identify the factors that influence LE, further research should evaluate panel data sets with a broader range of predictors.

Abbreviations: ABR: Adolescent birth rate; CO₂: Carbon dioxide; GNI: Gross national income; HIV: Human Immunodeficiency Virus; LE: Life expectancy; SD: Standard deviation; TFR: Total fertility rate; UN: United Nations; VIF: Variance inflation factor; WHO: World Health Organization

Ethics approval and consent to participate: Ethics approval for this study was not required since the data is secondary and is available in the public domain.

Consent for publication: Not applicable.

Availability of data materials: Data for this study were sourced from:

1. World Health Organization (WHO) 2016; and available here: http://apps.who.int/iris/bitstream/10665/206498/1/9789241565264_eng.pdf
2. World Health Statistics 2015; and available here: http://apps.who.int/iris/bitstream/handle/10665/170250/9789240694439_eng.pdf?sequence=1
3. World Population Data Sheet 2016; and available here: <https://assets.prb.org/pdf16/prb-wpds2016-web-2016.pdf>
4. Human Development Report 2016; and available here: http://hdr.undp.org/sites/default/files/2016_human_development_report.pdf

Competing interests: The authors declare that they have no competing interests.

Funding: The authors have no support or funding to report.

Authors' contributions: MNIM and ANMAB drafted the manuscript and were responsible for data collection and analyses. MNH, HTAK, and MNK contributed to the design of the study and editing the manuscript. In the end, the authors read and approved the final manuscript.

Acknowledgements: One of the authors (MNIM) is very grateful to the authority of ERASMUS+ Fellowship Program for granting a Visiting Scholar Fellowship to complete this study. The authors are very thankful to the Department of Sociology, Middle East Technical University (METU), Ankara; Turkey where the research has been conducted. Thanks also to the Editors and the referees for their valuable comments and criticism, which greatly improved this article.

References

1. WHO. World Health Statistics 2016. World Health Organization (WHO); Geneva, Switzerland; 2016.
2. Wakefield J. Ecologic Studies Revisited. *Annu Rev Public Health* 2008; 29:75-90.
3. WHO. World Health Report 2008: Primary health care now more than over. World Health Organization (WHO); Geneva, Switzerland; 2008.
4. Coile C, Diamond P, Gruber J and Jousten A. Delays in claiming social security benefits. *J Public Econ* 2002; 84:357-85.
5. Zhang J, Zhang J, Lee RD. Mortality decline and long-run economic growth. *J Public Econ* 2001; 80:485-507.
6. Shkolnikov VM, Andreev EM, Jasilionis D, Leinsalu M, Antonova OI, McKee M. The changing relation between education and life expectancy in central and Eastern Europe in the 1990s. *J Epidemiol Community Health* 2006; 60(10):875-81.
7. Mondal MNI, Shitan M. Impact of Socio-Health Factors on Life Expectancy in the Low and Lower Middle-Income Countries. *Iranian J Publ Health* 2013; 42(12):1354-62.
8. Mondal MNI, Shitan M. Relative Importance of Demographic,

- Socioeconomic and Health Factors on Life Expectancy in Low- and Lower-Middle-Income Countries. *J Epidemiol* 2014; 24(2):117-24.
9. Mondal MNI, Ullah MMMN, Islam MR, et al. Sociodemographic and Health Determinants of Inequalities in Life Expectancy in Least Developed Countries. *International Journal of MCH and AIDS* 2015; 4(1):01-10.
10. Khang YH, Yang S, Cho HJ, Jung-Choi K, Yun SC. Decomposition of socioeconomic variations in life expectancy at birth by age and cause of death among 4 million South Korean public servants and their dependents. *Int J Epidemiol* 2010; 39:1656-66.
11. Zimmer Z, Amornsirisomboon P. Socioeconomic status and health among older adults in Thailand: an examination using multiple indicators. *Soc Sci Med* 2001; 52:1297-311.
12. Hoi le V, Phuc HD, Dung TV, Chuc NT, Lindholm L. Remaining life expectancy among older people in a rural area of Vietnam: trends and socioeconomic inequalities during a period of multiple transitions. *BMC Public Health* 2009; 9:471.
13. Karlsson M, Nilsson T, Lyttkens CH, Leeson G. Income inequality and health: Importance of a cross-country perspective. *Soc Sci Med* 2010; 70(6):875-85.
14. Camargos MC, Machado CJ, do Nascimento Rodrigues R. Disability life expectancy for the elderly, city of Sao Paulo, Brazil, 2000: gender and educational differences. *J Biosoc Sci* 2007; 39(3):455-63.
15. Silventoinen K, Lahelma E. Health inequalities by education and age in four Nordic countries, 1986 and 1994. *J Epidemiol Community Health* 2002; 56:253-8.
16. Valkonen T, Sihvonen AP, Lahelma E. Health expectancy by level of education in Finland. *Soc Sci Med* 1997; 44:801-8.
17. Mackenbach JP, Kunst AE, Cavelaars AE, Groenhof F, Geurts JJ. Socioeconomic inequalities in morbidity and mortality in Western Europe. The EU Working Group on Socioeconomic Inequalities in Health. *The Lancet* 1997; 349:1655-9.
18. Lin CC, Rogot E, Johnson NJ, Sorlie PD, Arias E. A further study of life expectancy by socioeconomic factors in the National Longitudinal Mortality Study. *Ethn Dis* 2003; 13: 240-7.
19. Koons DN, Metcalf CJ, Tuljapurkar S. Evolution of delayed reproduction in uncertain environments: a life-history perspective. *Am Nat* 2008; 172:797-805.
20. Shaw JW, Horrace WC, Vogel RJ. The determinants of life expectancy: an analysis of the OECD health data. *South Econ J* 2005; 71:768-83.
21. Seubsman SA, Kelly MJ, Yiengprugsawan V, Sleigh AC. Thai Cohort Study Team. Gender, socioeconomic status, and self-rated health in a transitional middle-income setting: evidence from Thailand. *Asia Pac J Public Health* 2001; 23:754-65.
22. Lin RT, Chen YM, Chien LC, Chan CC. Political and Social Determinants of Life Expectancy in Less Developed Countries: A longitudinal Study. *BMC Public Health* 2012; 12:85.
23. Anderson KG. Life expectancy and the timing of historical events in developing countries. *Hum Nat* 2010; 21(2):103-23.
24. Tang JL, Griffiths S. Review paper: epidemiology, evidence-based medicine, and public health. *Asia Pac J Public Health* 2009; 21:244-51.
25. WHO. World Health Statistics 2015. World Health Organization (WHO); 2015, Geneva; Switzerland.
26. PRB. World Population Data Sheet 2016. Population Reference Bureau (PRB), 2016; Washington DC, USA.
27. UNDP. Human Development Report 2016. United Nations Development Program (UNDP), UN Plaza, New York; USA.
28. Alemayehu T, Haider J, Dereje H. Determinants of adolescent fertility in Ethiopia. *Ethiop J Health Dev* 2010; 24(1):30-7.
29. Chatterjee S, Hadi AS. Regression Analysis by Examples. Fifth Edition 2012. John Wiley & Sons, Inc., New Jersey; USA.
30. UN. World Population Prospects 2015-Data Booklet; 2015. United Nations

- (UN), Department of Economic and Social Affairs, Population Division; 2015.
31. UNICEF. Ending child marriage-Progress and prospects. United Nations Children's Fund (UNICEF), New York, USA; 2014.
 32. Marston C, Cleland J. The effects of contraception on obstetric outcomes. World Health Organization, Geneva; Switzerland; 2004.
 33. Romanelli F, Matheny SC. HIV Infection: The Role of Primary Care. *Am Fam Physician* 2009; 80(9):946-52.
 34. Low BS, Hazel A, Parker N, Welch KB. Influences on women's reproductive lives: unexpected ecological underpinnings. *Cross-Cultural Res* 2008; 42:201-19.
 35. Andre FE, Booy R, Bock HL, Clemens J, et al. Vaccination greatly reduces disease, disability, death and inequity worldwide. *Bull World Health Organ* 2008; 86(2):140-6.
 36. WHO. The World Health Report 2004. World Health Organization (WHO), Geneva; Switzerland; 2004.
 37. Yung-Tse H, Lawrence KW, Nazih KS. Handbook of Environment and Waste Management: Air and Water Pollution Control. World Scientific Publishing Ltd., Singapore; 2012.
 38. Silvestrina S, Homrich da Silva BC, Hirakatac VN André AS, et al. Maternal education level and low birth weight: a meta-analysis. *J Pediatr* 2013; 89(4):339-45.
 39. Cervellati M, Sunde U. Life expectancy, schooling, and lifetime labor supply: Theory and evidence revisited. *Econometrica* 2013; 81: 2055-86.
 40. Tariq S, Ul-Haq Z, Imran A, Mehmood U, Aslam MU, Mahmood K. CO₂ emissions from Pakistan and India and their relationship with economic variables. *Applied Ecology and Environmental Research* 2017; 15(4):1301-12.