



UWL REPOSITORY

repository.uwl.ac.uk

Cost-benefit analysis of surveillance for surgical site infection following caesarean section

Wloch, Catherine, van Hoek, Albert Jan, Green, Nathan, Conneely, Joanna, Harrington, Pauline, Sheridan, Elizabeth, Wilson, Jennie ORCID: <https://orcid.org/0000-0002-4713-9662> and Lamagni, Theresa (2020) Cost-benefit analysis of surveillance for surgical site infection following caesarean section. *BMJ Open*, 10 (7).

<http://dx.doi.org/10.1136/bmjopen-2020-036919>

This is the Accepted Version of the final output.

UWL repository link: <https://repository.uwl.ac.uk/id/eprint/7079/>

Alternative formats: If you require this document in an alternative format, please contact: open.research@uwl.ac.uk

Copyright: Creative Commons: Attribution-Noncommercial 4.0

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy: If you believe that this document breaches copyright, please contact us at open.research@uwl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Cost-benefit analysis of surveillance for surgical site infection following caesarean section

Figure 1. Change in surgical site infection (SSI) risk between consecutive 3 month surveillance periods for 7 hospitals during the multi-centre caesarean section study





Figure 2. Balance of surveillance cost versus savings from reductions of 10, 20 and 30% per surveillance period for surveillance strategies of one quarter a year, two quarters a year and continuous surveillance for starting surgical site infection (SSI) risk of 5%

Model assumes reductions in infection risk are achieved in conjunction with improvement programmes during surveillance periods and maintained between each surveillance period. No further reductions in risk of infection were included in the model once a postulated minimum SSI risk of 3% was reached.

Key: Surveillance strategy Reductions in SSI risk



i) SSI risk (%)



ii) balance of discounted cost versus savings





Figure 3. Balance of surveillance cost versus savings from reductions of 10, 20 and 30% per surveillance period for surveillance strategies of one quarter a year, two quarters a year and continuous surveillance for starting surgical site infection (SSI) risk of 10%

Model assumes reductions in infection risk are achieved in conjunction with improvement programmes during surveillance periods and maintained between each surveillance period. No further reductions in risk of infection were included in the model once a postulated minimum SSI risk of 3% was reached.

Key: Surveillance strategy Reductions in SSI risk



i) SSI risk (%)



ii) balance of discounted cost versus savings





Figure 4. Balance of surveillance cost versus savings from reductions of 10, 20 and 30% per surveillance period for surveillance strategies of one quarter a year, two quarters a year and continuous surveillance for starting surgical site infection (SSI) risk of 15%

Model assumes reductions in infection risk are achieved in conjunction with improvement programmes during surveillance periods and maintained between each surveillance period. No further reductions in risk of infection were included in the model once a postulated minimum SSI risk of 3% was reached.

Key: Surveillance strategy Reductions in SSI risk



i) SSI risk (%)



ii) balance of discounted cost versus savings





Figure 5. Balance of surveillance cost versus savings from reductions in surgical site infection risk of 10, 20 and 30% per surveillance period for baseline surgical site infection (SSI) risk of 10 or 15% using a variable surveillance strategy (continuous surveillance when the infection risk is above 10%, two quarters per year surveillance for infection risk between 5 and 10% and one quarter per year surveillance for infection risk below 5%)


Model assumes reductions in risk of infection are achieved in conjunction with improvement programmes during surveillance periods and maintained between each surveillance period. No further reductions in risk of infection were included once a postulated minimum SSI risk of 3% was reached.

Key: Reductions in risk

Baseline SSI risk



i) SSI risk (%)



ii) balance of discounted cost versus savings




Figure 6 Cumulative discounted prevented costs against costs of surveillance after 5-year surveillance programme - 15% Baseline surgical site infection risk





Figure 7 Cumulative discounted prevented costs against costs of surveillance after 5-year surveillance programme - 10% Baseline surgical site infection risk





Figure 8 Cumulative discounted prevented costs against costs of surveillance after 5-year surveillance programme - 5% Baseline surgical site infection risk



*Variable surveillance strategy is equivalent to once-a-year surveillance where SSI risk is <5%