

UWL REPOSITORY

repository.uwl.ac.uk

Antibiotic resistance and molecular characteristics of methicillin-resistant
Staphylococcus epidermidis recovered from hospital personnel in China

Xu, Zhen, Cave, Rory, Chen, Liqin, Yangkyi, Tsetan, Liu, Yan, Li, Ke, Meng, Ge, Niu, Kaijun, Zhang, Wanqi, Tang, Naijun, Shen, Jun and Mkrtchyan, Hermine (2020) Antibiotic resistance and molecular characteristics of methicillin-resistant Staphylococcus epidermidis recovered from hospital personnel in China. *Journal of Global Antimicrobial Resistance*, 22. pp. 195-201. ISSN 2213-7165

<http://dx.doi.org/10.1016/j.jgar.2020.02.013>

This is the Published Version of the final output.

UWL repository link: <https://repository.uwl.ac.uk/id/eprint/7298/>

Alternative formats: If you require this document in an alternative format, please contact:
open.research@uwl.ac.uk

Copyright: Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy: If you believe that this document breaches copyright, please contact us at open.research@uwl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Antibiotic resistance and molecular characteristics of methicillin-resistant *Staphylococcus epidermidis* recovered from hospital personnel in China

Zhen Xu^a, Rory Cave^b, Liqin Chen^a, Tsetan Yangkyi^c, Yan Liu^d, Ke Li^a, Ge Meng^a, Kaijun Niu^a, Wanqi Zhang^a, Naijun Tang^a, Jun Shen^a, Hermine V. Mkrtchyan^{b,*}

^a Tianjin Key Laboratory of Environment, Nutrition and Public Health, Department of Toxicology and Sanitary Chemistry, School of Public Health, Tianjin Medical University, Tianjin, China

^b School of Health, Sport and Biosciences, University of East London, London, UK

^c CDC of Biru County, Tibet, China

^d Tianjin Central Hospital of Gynecology Obstetrics, Tianjin, China

ARTICLE INFO

Article history:

Received 30 July 2019

Received in revised form 9 December 2019

Accepted 17 February 2020

Available online 22 February 2020

Keywords:

Staphylococcus epidermidis

Hospital personnel

mecA

SCCmec

MLST

Sequence type

ABSTRACT

Objectives: *Staphylococcus epidermidis* is a major nosocomial pathogen predominantly associated with indwelling medical device infections. Studies reporting on *S. epidermidis* recovered from hospital personnel in China are scarce. The aim of this study was to evaluate the carriage and antibiotic resistance of *S. epidermidis* among the hospital personnel in Tianjin, China and provide insights into their genetic diversity.

Methods: One hundred and seven *S. epidermidis* isolates were recovered from 68 hospital personnel in two public hospitals in Tianjin between March 2018 and May 2018. Staphylococcal cassette chromosome *mec* (SCCmec) types were determined by the combination of *mec* and *ccr* complexes. Multi-locus sequence typing was used to determine the sequence types (ST) of *S. epidermidis* isolates.

Results: Sixty-two (76.5%) isolates were determined to be methicillin-resistant *S. epidermidis* (MRSE). Thirty-five (51%) of 68 hospital personnel carried *S. epidermidis*, of which 32 (91%) were carriers of MRSE. All 62 MRSE isolates had high levels of resistance to penicillin (90%) and cefoxitin (100%). Thirty-seven (60%) isolates carried SCCmec type IV, followed by 15 (24%) carrying SCCmec V, and 4 (6%) SCCmec II. Novel STs were assigned to four *S. epidermidis* isolates (ST832, ST833, ST834 and ST835).

Conclusions: In this study, the majority of MRSE belonged to cluster II domain of CC2. The ST59-IV was a dominant clone among isolates recovered from hospital personnel. Determination of new MLST types confirmed the genetic diversity of these isolates. These observations highlight the need to review the infection control strategies to reduce the carriage of MRSE among hospital personnel.

© 2020 The Authors. Published by Elsevier Ltd on behalf of International Society for Antimicrobial Chemotherapy. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Staphylococcus epidermidis is a common colonizer of the human skin [1], but also one of the major opportunistic pathogens responsible for medical device associated infections (MDAI). However, the role of *S. epidermidis* as a nosocomial pathogen has been underestimated until late 1980s [2]. Recently, several studies reported that the distribution of *S. epidermidis* in human clinical samples was higher than infections caused by *S. aureus* in Greece and India [1,3]. The ever increasing antibiotic resistance

aggravates the problem, and poses a greater challenge for the control of hospital acquired infections [4].

Methicillin resistance is mediated by the *mecA* gene, which encodes penicillin binding protein 2a that has low affinity to β -lactam antibiotics, and thus confers methicillin resistance [5]. The *mecA* gene is located on a mobile genetic island named staphylococcal cassette chromosome *mec* (SCCmec). Up to date 11 SCCmec types have been assigned based on combinations of *mec* and *ccr* complexes [6]. Molecular typing methods, including multi-locus sequence typing have been instrumental to identify highly diverse genetic lineages of *S. epidermidis* [7]. *S. epidermidis* species have been assigned onto one major clonal complex (CC2), 8 minor clonal complexes and 13 singletons [8].

While there are many studies reporting the molecular epidemiology of *S. aureus*, studies reporting the molecular

* Corresponding author.

E-mail address: h.mkrtchyan@uel.ac.uk (H.V. Mkrtchyan).

characteristics of *S. epidermidis* are fragmentary. It has been documented that hospital personnel are often carriers of methicillin-resistant staphylococci and aid the dissemination of hospital-acquired infections [9]. However, little is known about the genotypic diversity of methicillin-resistant *S. epidermidis* (MRSE) among hospital personnel in China. To our knowledge, only two studies have reported acquisition of MRSE among hospital personnel with 89% of hospital personnel carrying MRSE in Sweden [10] and 30% in Shanghai, China [11].

In this study, we evaluate the carriage and susceptibility patterns of MRSE among the hospital personnel recovered from two hospitals in Tianjin, China and provide insights into their genetic diversity.

2. Material and methods

2.1. Study protocol

As a part of a small surveillance study to assess the carriage of MRSE among the hospital personnel, 107 samples were recovered from the hospital personnel ($n=68$) in two public hospitals in Tianjin city, North China, between March 2018 and May 2018. Samples were taken from doctors and nurses' hands ($n=68$) and nasal cavity ($n=39$).

The research protocol and informed consent was approved by the Ethics committee of Tianjin Science and Technology Commission (approval No TMUaMEC2017017). All research was performed in accordance with the relevant guidelines and regulations. Informed consent was obtained from all participants.

All specimens were inoculated onto mannitol salt agar (MSA, Oxoid, Basingstoke, UK) and incubated aerobically at 37 °C for 24–48 h.

2.2. Identification

S. epidermidis isolates were subjected to partial 16S ribosomal RNA (rRNA) gene sequencing using the primers and polymerase chain reaction (PCR) conditions as described previously [12]. Amplified PCR products were sequenced by Sangon Biotech (Shanghai, China). Homology searches were carried out using BLAST tool (NCBI: <https://www.ncbi.nlm.nih.gov/>) [13].

2.3. Antimicrobial susceptibility test

Susceptibility to 10 antibiotics was tested for all isolates using standard disk diffusion method. The antibiotics tested included: CN: gentamicin (10 µg), E: erythromycin (15 µg), FOX: cefoxitin (30 µg), P: penicillin (10 units), T: tetracycline (30 µg), TEC: teicoplanin (30 µg), DA: clindamycin (2 µg), CHL: chloramphenicol (30 µg), LZD: linezolid (30 µg), and RD: rifampin (5 µg). The isolates were categorized as susceptible, intermediate resistant, or resistant according to the recommendations of Clinical and Laboratory Standard Institute (CLSI) [14].

2.4. *mecA* gene determination and *SCCmec* typing

The *mecA* gene was determined for all isolates using PCR method as described previously [15]. *SCCmec* types were determined for all *mecA* positive isolates by evaluating the *mec* and *crr* complexes [15].

2.5. Multi-locus sequence typing of *S. epidermidis*

The *mecA* positive isolates ($n=60$) were analysed for seven housekeeping genes using MLST as described previously [16]. In brief, seven house-keeping genes of each isolate were amplified by

using *arcC*, *aroE*, *gtr*, *mutS*, *pyr*, *tpi*, and *yqil* primers and then sequenced by Sangon Biotech. The sequence type (ST) for each isolate was assigned using the *S. epidermidis* MLST database (<https://pubmlst.org/>).

2.6. Statistics

The goeBURST algorithm (<http://www.phyloviz.net/goeburst/>) was used to build goeBURST diagram tree [11]. A hierarchy clustering heatmap was constructed using the MLST data, isolation source and *SCCmec* type and resistance profile of the *mecA* positive isolates using the R package Heatmap.plus (<https://www.rdocumentation.org/packages/heatmap.plus/versions/1.3/topics/heatmap.plus.package>). Principal component analysis (PCA) was performed to distinguish between antibiotic resistance profiles by site using the R packages 'FactoMineR' (<https://cran.r-project.org/web/packages/FactoMineR/index.html>) and factextra (<https://cran.r-project.org/package=factextra>).

The χ^2 test was used to determine the quantitative variables. A *P*-value <0.05 was considered statistically significant.

3. Results

3.1. Study protocol and identification of isolates

Samples were taken from hands of hospital personnel ($n=68$) and nasal cavity ($n=39$). Thirty-five of 68 (51%) hospital personnel were identified as carriers of *S. epidermidis*, including 24 of 68 (35%) carried *S. epidermidis* on their hands. Nineteen of 39 (49%) hospital personnel were nasal carriers. In addition, 8 of 68 (12%) carried *S. epidermidis* both on their hands and nasal cavity. A total of 165 isolates were recovered from the hands or nasal cavity samples of the hospital personnel working in two public hospitals in Tianjin city, China between March 2018 and May 2018. Eighty-one (81/165; 49%) isolates were identified as *S. epidermidis*, including 40 isolates recovered from hands and 41 *S. epidermidis* isolated from nasal cavity.

3.2. *mecA* gene determination of *S. epidermidis*

The *mecA* gene was determined in 62 (62/81; 76.5%) isolates, including 30 (30/40; 75%) recovered from hospital personnel hands, and 32 (32/41; 78%) from nasal cavity. In addition, 33 (33/68; 49%) hospital personnel were carriers of MRSE, including 19 (19/68; 28%) carried MRSE on their hands and 19 (19/68; 28%) carried MRSE in their nasal cavity. Moreover, five (5/68; 7%) of the hospital personnel were carriers of MRSE both on their hands and in the nasal cavities (Table 1).

3.3. Antimicrobial susceptibility of MRSE

The disc diffusion method was used to test the susceptibility of all *S. epidermidis* isolates ($n=81$) against 10 antibiotics. Eighty-one (100%) isolates were resistant to at least one antibiotic. Resistance to penicillin and cefoxitin was observed in 70 of 81 (86%) and 62 of 81 (77%) isolates respectively. Thirty-four (42%) isolates were resistant to erythromycin ($R \leq 1.3$ cm), whereas 18 (26%) isolates showed intermediate resistance ($I = 1.4–2.2$ cm) towards erythromycin. Five (8%) isolates were resistant to gentamicin ($R \leq 1.2$ cm), and four (5%) to tetracycline ($R \leq 1.4$ cm) (Table 1). Thirty-one (38%) isolates were resistant to teicoplanin ($R \leq 1.0$ cm), 34 (42%) isolates to clindamycin ($R \leq 1.4$ cm), 6 (7%) to chloramphenicol ($R \leq 1.2$ cm), 6 (7%) to linezolid ($R \leq 2.0$ cm), and 13 (16%) to rifampin ($R \leq 1.6$ cm). All the *mecA*-negative *S. epidermidis* isolates were susceptible to cefoxitin. The majority of *mecA*-negative isolates were susceptible to gentamicin,

Table 1Antibiotic resistance and molecular characterization of *mecA* gene positive *Staphylococcus epidermidis*.

ID	Hospital	Personnel	Sites	<i>mecA</i>	<i>mec</i>	<i>ccr</i>	SCC <i>mec</i>	FOX ₃₀	Other antibiotics								
									CHL ₃₀	CN ₁₀	DA ₂	E ₁₅	LZD ₃₀	P ₁₀	RD ₅	TEC ₃₀	T ₃₀
1	A	A1	H	+	B	2	IV	R	S	S	R	S	S	R	S	I	S
2	A	A12	H	+	B	2	IV	R	R	S	I	R	S	R	S	I	S
3	A	A2	H	+	B	2	IV	R	S	S	I	I	S	R	S	R	S
4	A	A3	H	+	B	2	IV	R	S	S	S	S	S	S	S	R	S
5	A	A1	H	+	B	2	IV	R	S	S	I	S	R	R	S	I	S
6	A	A1	H	+	B	2	IV	R	S	S	S	S	S	R	R	R	I
7	A	A2	H	+	B	2	IV	R	S	S	I	S	S	R	S	R	S
8	A	A4	H	+	B	2	IV	R	S	S	S	I	R	R	S	R	S
9	A	A12	H	+	B	2	IV	R	R	S	I	R	S	R	S	I	S
10	A	A13	H	+	B	2	IV	R	S	S	I	R	S	R	S	R	S
11	A	A1	H	+	B	2	IV	R	S	S	I	S	S	R	S	R	S
12	B	B2	H	+	B	2	IV	R	R	R	R	R	R	R	R	R	S
13	B	B3	H	+	B	2	IV	R	R	R	R	I	S	R	S	R	S
14	B	B3	H	+	B	2	IV	R	R	I	I	S	R	R	R	R	S
15	B	B4	H	+	B	2	IV	R	S	S	I	S	S	R	S	I	S
16	B	B4	H	+	B	2	IV	R	S	S	S	S	S	R	S	I	S
17	B	B8	H	+	B	2	IV	R	S	S	I	I	R	R	S	R	I
18	B	B11	H	+	B	2	IV	R	S	S	S	R	S	R	S	I	S
19	B	B9	H	+	B	2	IV	R	S	S	S	S	S	R	S	I	S
20	B	B16	H	+	B	2	IV	R	S	S	S	I	S	R	S	I	S
21	B	B18	H	+	B	2	IV	R	S	R	S	S	S	S	S	I	S
22	B	B15	H	+	C	5	V	R	S	S	I	R	S	R	S	I	S
23	B	B5	H	+	C	5	V	R	S	S	S	R	S	R	S	R	I
24	B	B1	H	+	C	5	V	R	S	S	I	R	S	R	S	R	S
25	A	A5	H	+	C	5	V	R	S	S	S	R	S	R	R	I	I
26	A	A3	H	+	C	5	V	R	R	S	R	R	S	R	S	R	S
27	A	A4	H	+	C	5	V	R	S	S	R	R	S	R	R	I	S
28	A	A3	H	+	C	5	V	R	S	S	R	I	S	R	S	I	S
29	A	A5	H	+	C	2	C/2	R	S	I	R	R	S	R	R	I	S
30	B	B17	H	+	C	2	C/2	R	S	S	R	S	S	R	S	I	S
31	A	A11	NC	+	A	2	II	R	S	S	R	R	S	R	S	I	S
32	A	A11	NC	+	A	2	II	R	S	S	R	R	S	R	S	I	S
33	B	B19	NC	+	A	2	II	R	S	S	I	R	S	S	S	I	S
34	B	B20	NC	+	A	2	II	R	S	S	S	R	S	R	S	R	S
35	A	A8	NC	+	B	2	IV	R	S	S	I	S	S	R	S	R	S
36	A	A8	NC	+	B	2	IV	R	S	S	R	R	S	R	S	R	S
37	A	A12	NC	+	B	2	IV	R	S	S	S	R	S	R	S	R	S
38	A	A10	NC	+	B	2	IV	R	S	S	S	I	S	R	S	I	S
39	B	B6	NC	+	B	2	IV	R	S	R	S	S	S	R	S	I	S
40	B	B7	NC	+	B	2	IV	R	S	S	R	S	S	R	S	R	S
41	A	A6	NC	+	B	2	IV	R	S	S	I	S	R	R	S	I	S
42	A	A6	NC	+	B	2	IV	R	S	S	R	I	S	R	S	I	S
43	A	A10	NC	+	B	2	IV	R	S	S	S	I	S	R	S	R	S
44	B	B17	NC	+	B	2	IV	R	S	S	I	S	S	R	R	I	S
45	B	B16	NC	+	B	2	IV	R	S	S	I	S	S	R	S	I	S
46	B	B12	NC	+	B	2	IV	R	S	S	R	S	S	S	S	I	S
47	B	B12	NC	+	B	2	IV	R	S	S	R	I	S	S	S	S	S
48	B	B10	NC	+	B	2	IV	R	S	S	R	I	S	R	R	R	S
49	B	B16	NC	+	B	2	IV	R	S	S	I	S	S	R	S	I	S
50	B	B18	NC	+	B	2	IV	R	S	S	I	S	S	R	S	R	S
51	A	A7	NC	+	C	5	V	R	S	S	R	I	S	R	S	I	S
52	A	A7	NC	+	C	5	V	R	S	S	I	I	S	R	R	I	S
53	B	B13	NC	+	C	5	V	R	S	S	R	R	S	R	S	R	S
54	B	B13	NC	+	C	5	V	R	S	S	R	R	S	R	R	R	S
55	B	B10	NC	+	C	5	V	R	S	S	R	R	S	R	R	I	R
56	B	B10	NC	+	C	5	V	R	S	S	I	R	S	R	R	I	S
57	B	B14	NC	+	C	5	V	R	S	S	R	I	S	R	S	I	S
58	B	B18	NC	+	C	5	V	R	S	S	I	I	S	R	S	R	S
59	B	B1	NC	+	C	3	C/3	R	S	R	I	I	S	R	S	I	R
60	B	B6	NC	+	C	3	C/3	R	S	S	I	S	S	R	S	I	S
61	B	B6	NC	+	C	3	C/3	R	S	S	S	S	S	R	S	I	S
62	A	A9	NC	+	NT	3	NT	R	S	I	I	S	S	S	S	I	S

Penicillin ($S \geq 2.9$ cm, $R \leq 2.8$ cm), erythromycin ($S \geq 2.3$ cm, $I = 1.4$ – 2.2 cm, $R \leq 1.3$ cm), cefoxitin ($S \geq 2.5$ cm, $R \leq 2.4$ cm), gentamicin ($S \geq 1.5$ cm, $I = 1.3$ – 1.4 cm, $R \leq 1.2$ cm), tetracycline ($S \geq 1.9$ cm, $I = 1.5$ – 1.8 cm, $R \leq 1.4$ cm), teicoplanin ($S \geq 1.4$ cm, $I = 1.1$ – 1.3 cm, $R \leq 1.0$ cm), clindamycin ($S \geq 2.1$ cm, $I = 1.5$ – 2.0 cm, $R \leq 1.4$ cm), chloramphenicol ($S \geq 1.8$ cm, $I = 1.3$ – 1.7 cm, $R \leq 1.2$ cm), linezolid ($S \geq 2.1$ cm, $R \leq 2.0$ cm), rifampin ($S \geq 2.0$ cm, $I = 1.7$ – 1.9 cm, $R \leq 1.6$ cm).

^aMLST type not determined.

A, hospital 1; B, hospital 2; CHL, chloramphenicol; CN, gentamicin; DA, clindamycin; E, erythromycin; FOX, cefoxitin; H, hand; LZD, linezolid; NC, nasal cavity; NT, non-typeable; P, penicillin; RD, rifampin; T, tetracycline; TEC, teicoplanin.

Table 2
Antibiotic resistance of *mecA* gene negative *Staphylococcus epidermidis*.

ID	Hospital	Personnel	<i>mecA</i>	CHL ₃₀	CN ₁₀	DA ₂	E ₁₅	FOX ₃₀	LZD ₃₀	P ₁₀	RD ₅	TEC ₃₀	T ₃₀
63	A	A1	–	S	S	R	I	S	S	R	S	R	S
64	A	A1	–	S	S	I	R	S	S	S	S	R	S
65	A	A1	–	S	S	R	R	S	S	S	S	I	S
66	A	A5	–	S	S	R	S	S	S	S	S	I	S
67	A	A6	–	S	S	R	R	S	S	R	S	I	S
68	A	A7	–	S	S	R	I	S	S	R	S	I	R
69	A	A8	–	S	S	R	R	S	S	R	S	I	S
70	A	A8	–	S	S	R	R	S	S	R	S	I	S
71	A	A9	–	S	S	I	R	S	S	R	R	R	S
72	A	A10	–	S	S	I	S	S	S	R	S	I	S
73	A	A10	–	S	S	I	S	S	S	R	S	R	S
74	A	A11	–	S	S	R	R	S	S	R	S	I	S
75	A	A11	–	S	S	R	S	S	S	R	S	I	S
76	A	A11	–	S	S	R	R	S	S	R	S	I	S
77	A	A13	–	S	S	S	R	S	S	R	S	I	S
78	A	A14	–	S	S	R	S	S	S	R	S	R	R
79	A	A15	–	S	S	I	R	S	S	R	S	R	S
80	A	A15	–	S	S	R	R	S	S	S	S	I	S
81	B	B20	–	S	I	R	R	S	S	S	S	I	I

Penicillin ($S \geq 2.9$ cm, $R \leq 2.8$ cm), erythromycin ($S \geq 2.3$ cm, $I = 1.4$ – 2.2 cm, $R \leq 1.4$ cm) cefoxitin ($S \geq 2.5$ cm, $R \leq 2.4$ cm), gentamicin ($S \geq 1.5$ cm, $I = 1.3$ – 1.4 cm, $R \leq 1.2$ cm), tetracycline ($S \geq 1.9$ cm, $I = 1.5$ – 1.8 cm, $R \leq 1.4$ cm), teicoplanin ($S \geq 1.4$ cm, $I = 1.1$ – 1.3 cm, $R \leq 1.0$ cm), clindamycin ($S \geq 2.1$ cm, $I = 1.5$ – 2.0 cm, $R \leq 1.4$ cm), chloramphenicol ($S \geq 1.8$ cm, $I = 1.3$ – 1.7 cm, $R \leq 1.2$ cm), linezolid ($S \geq 2.1$ cm, $R \leq 2.0$ cm), rifampin ($S \geq 2.0$ cm, $I = 1.7$ – 1.9 cm, $R \leq 1.6$ cm).

A, hospital 1; B, hospital 2; CHL, chloramphenicol; CN, gentamicin; DA, clindamycin; E, erythromycin; FOX, cefoxitin; H, hand; LZD, linezolid; N, nose; NT non-typeable; P, penicillin; RD, rifampin; T, tetracycline; TEC, teicoplanin.

chloramphenicol, linezolid and rifampin. Fourteen (14/19; 73.7%) *mecA*-negative isolates were resistant to penicillin, followed by 13 (13/19; 68.4%) to clindamycin, 12 (12/19; 63.2%) to erythromycin, and 6 (6/19; 31.6%) to teicoplanin (Table 2).

3.4. SCCmec typing of MRSE

SCCmec types were determined for all 62 MRSE isolates. Thirty-seven (60%) isolates (from hands $n = 21$; from nasal cavity $n = 16$) harboured SCCmec type IV, followed by 15 (24%) SCCmec type V, and 4 (6%) SCCmec type II. In addition to this, six isolates were non-typeable, including three (5%) isolates harboured a combination of class C *mec* complex and the *ccr* type 3 complex, and two (3%) isolates carried class C *mec* complex and *ccr* type 2, whereas one (2%) isolate lacked *mec* complex but carried *ccr* type 3 complex (Table 1).

3.5. Multi-locus sequence typing of MRSE

MLST was performed to determine the STs of 60 MRSE, including 29 recovered from hospital personnel hands ($n = 18$) and 31 from nasal samples ($n = 18$). The ST59 ($n = 19$) was the most common ST, followed by ST35 ($n = 4$), ST14 ($n = 3$), ST57 ($n = 3$), ST218 ($n = 3$), ST20 ($n = 3$), ST49 ($n = 2$), ST69 ($n = 2$), ST110 ($n = 2$), ST152 ($n = 2$), ST227 ($n = 2$), ST466 ($n = 2$), ST5 ($n = 1$), ST6 ($n = 1$), ST17 ($n = 1$), ST50 ($n = 1$), ST84 ($n = 1$), ST130 ($n = 1$), ST190 ($n = 1$), ST192 ($n = 1$), and ST234 ($n = 1$). In addition, four isolates contained novel STs that were assigned as: ST832, ST833, ST834 and ST835. All 60 MRSE isolates were clustered into clonal complex 2 (CC2) by the goeBURST algorithm. In addition, no singleton was detected (Fig. 1).

3.6. Hierarchical clustering and PCA analysis

A hierarchical cluster heatmap was performed on 60 MRSE isolates based on their site of isolation, SCCmec and MLST types (Fig. 2A) to their antibiotic resistant profile. These analyses showed that there was no obvious clustering based on the site of isolation, SCCmec and MLST types. This was further showed by PCA analysis

on site isolates as the confidence ellipse overlapped each other (Fig. 2B).

3.7. Statistical analysis

The χ^2 test was used to analyse the quantitative variables. A *P*-value < 0.05 was considered statistically significant.

4. Discussion

Staphylococcus epidermidis is a major nosocomial pathogen responsible for device associated infections [1]. Hospital personnel have an important role of being in direct contact with patients and subsequently play a key role in cross-transmission of MRSE [9]. In this study, we report the antimicrobial resistance patterns and genetic diversity of MRSE among the hospital personnel in Tianjin, China.

The resistance of MRSE towards penicillin is well documented. Xin et al [11] reported that 90.4% of *S. epidermidis* isolates recovered from hospital environments were resistant to penicillin. Consistent with their results we have shown that 90% of MRSE isolates in our study were phenotypically resistant to penicillin. In addition, we have previously reported high levels of antibiotic resistance towards penicillin in environmental *S. epidermidis* (70%) isolates recovered from London [12]. Interestingly, in this study the resistance rates towards gentamicin (8%), and erythromycin (35%) were relatively low compared with others that demonstrated resistance towards gentamicin and erythromycin as 20% and 50% respectively [10,11]. Previously, it was shown that cefoxitin disk diffusion method was preferable for routine methicillin resistance screening of *S. aureus*, *S. epidermidis* and other CoNS isolates [17,18]. In this study, 62 (62/81; 77%) *S. epidermidis* isolates were resistant to cefoxitin, all of which carried the *mecA* gene. Li et al reported that resistance of clinical *S. epidermidis* isolates towards penicillin, cefoxitin, gentamicin and erythromycin were 100%, 100%, 77.8% and 72.2% respectively [19]. In this study, the resistance towards non- β -lactam antibiotics in *S. epidermidis* isolates recovered from hospital personnel was significantly lower than that of clinical *S. epidermidis* isolates, whereas the resistance towards β -lactam




Fig. 1. goeBURST analysis of 60 MRSE isolates recovered from hospital personnel in China.

ST nodes: dark green circles- sub-group founder; light blue circles- common nodes; light green- SCCmec II; dark yellow- SCCmec IV; turquoise- SCCmec V; red- SCCmec C/2; blue- SCCmec C/3; purple- non-typable SCCmec.

antibiotics was similar to clinical *S. epidermidis* isolates. In this study, 100% *S. epidermidis* isolates were phenotypically resistant to at least 1 antibiotic. Twenty-two out of 81 (27%) *S. epidermidis* isolates were resistant to 2 antibiotics, 27 isolates (33%) were resistant to 3 antibiotics, 18 isolates (22%) were resistant to 4

antibiotics, 5 (6%) isolates were resistant to 5 antibiotics, 5 (6%) isolates were resistant to 6 antibiotics and 1 (1.2%) isolate was resistant to 9 antibiotics.

In this study 32 (91%) of 35 *S. epidermidis* carriers were identified to carry MRSE. Although, in this study the carriage of MRSE among the hospital personnel was significantly higher than among the hospital personnel in Shanghai, China [11], our findings were consistent with the MRSE rates (89%) among the hospital personnel in Intensive Care Unit in Sweden [10]. Previously, it was reported that the rate of MRSE carriage among the volunteers and in the environment was 11% [20], which was significantly lower than the rate of MRSE carriage among the hospital personnel.

The size of SCCmec IV and V is smaller than those of SCCmec types I, II and III, and thus conferring SCCmec IV and V increased mobility and dissemination ability [21]. Li et al reported that 35% of clinical MRSE harboured SCCmec types IV and V in China [19]. In this study, 60% of MRSE harboured SCCmec type IV, followed by 24% harbouring SCCmec type V. We determined that 6% of MRSE that carried SCCmec type II were recovered from nasal samples. SCCmec II was reportedly identified in clinical *S. aureus*, *S. epidermidis* and other [19]. Interestingly, we did not identify SCCmec types I, and III, which in contrast have been identified by Xin et al in isolates recovered from hospital personnel in Shanghai [11]. Six previously unclassified SCCmec types were determined in this study, including three carrying class C *mec* complex and *ccr* 3, and two had a combination of class C *mec* complex and *ccr* 2. In addition, one MRSE was classified as SCCmec 12263 since it lacked *mec* complex.

For *S. epidermidis*, 1 major clonal complex (CC2), 8 minor clonal complexes and 13 singletons have been categorized [16]. In this study, all MLST types were classified into one major CC2 clonal complex, hence we did not detect isolates belonging to minor clonal complexes or singletons (Fig. 1). However, although belonging to one CC group, in our study, we detected a diversity of MLST types ($n = 25$). Miragaia et al. summarized the STs of CC2 that accounted for 74% of the *S. epidermidis* population, and divided them into clusters: cluster I: included the predicted ancestor (ST2) and cluster II: included several subgroup founders (ST5, 6, 57, 85, and 89) [16]. In this study, the majority of STs were categorized into cluster II, including those identified as subgroup founders: ST5, 49, 57, 110, 152, and 835 (Fig. 1). In addition, two isolates belonged to ST35. Miragaia et al. reported that ST2 of cluster I is the most widely disseminated ST that has been identified in different countries [16]. In this study, no ST2 was determined among the *S. epidermidis* isolates recovered from two hospital personnel in Tianjin, China. However, the ST59 of cluster II was the most frequently identified. Isolates belonging to ST59 were reported to be only the second to the ST2, a prominent cause of clinical infections in China [11]. Thus, the prevalence of ST59 among the hospital personnel is rather worrying.




Fig. 2. Hierarchy cluster heatmap and PCA analysis of MRSE isolates antibiotic resistance profile. (A) Hierarchy clustered heatmap. Red tile, resistant; black tile, intermediate; green tile, sensitive. (B) PCA analysis; 95% confidence ellipse.

Previously, whole genome sequencing studies revealed that environmental *S. epidermidis* ST59 isolate carried SCCmec type IV mobile genetic element [22]. To the best of our knowledge, this is the first study reporting that isolates belonging to ST59 harboured unclassified SCCmec types (C/2 and C/3) (Fig. 1).

S. epidermidis ST5 has been associated with clinical and animal infections [23], whereas *S. epidermidis* ST6, ST20, ST110, ST130 and ST152 were associated with clinical infections [8,24,25]. It is largely known that the main transmission route of staphylococci is person to person and hospital personnel play a major role in such transmission [1]. Thus, our study provides further insights on the colonization of the hospital personnel with *S. epidermidis*. In addition, Xin et al. reported a variety of *S. epidermidis* STs identified among hospital personnel (ST14, ST16, ST54, ST88, ST153, ST171, ST184, ST190, ST192, ST193, ST194, ST203, ST204, ST210, ST218, ST219, ST220, ST226, ST233, ST237, ST262, ST267, ST291, ST327, ST362, ST387, ST406, and ST466) [11]. Some of the STs (ST14, ST190, ST218, and ST466) identified in this study were consistent with their findings. However, in addition to this we also report STs (ST35, ST57, ST69, ST84, ST227, and ST234) that were not identified by Xin et al. [11]. In addition, four new MLST types were determined in this study.

We did not detect any obvious clustering based on site of isolation, SCCmec types and MLST types (Fig. 2A). This was further demonstrated by PCA analysis on site isolates as the confidence ellipse overlapped each other (Fig. 2B). Other authors reported discrepancies between the antibiotic resistance profile and different types of SCCmec elements [26]. This report shows that SCCmec type III and IV in *S. epidermidis* were more likely to be resistant to a larger number of antibiotics.

5. Conclusion

The main limitation of this study is that only samples recovered from the hospital personnel were included. The carriage of MRSE (91%) was unusually high among hospital personnel who were carriers of *S. epidermidis*. We also observed different MRSE colonization rate in hospital personnel in two hospitals. The MRSE isolates had high resistance rates towards β -lactam antibiotics, but low resistance rates against non- β -lactam antibiotics. Moreover, the majority of MRSE in this study belonged to cluster II domain of CC2. ST59-IV was the predominant clone among isolates recovered from hospital personnel, a ST that was reportedly associated with clinical infections. Moreover, new MLST types were determined, thus further confirming the genetic variability of these isolates. Our data demonstrate that the hospital personnel may well act as a reservoir of antimicrobial resistance pathogens. Undoubtedly, there is a need to review the infection control strategies and implement appropriate screening and monitoring measures to identify the high carriage of MRSE among hospital personnel on timely manner.

Funding

This work was funded by Tianjin Municipal Science and Technology Bureau, number: 17JCYBJC43000, China.

Ethical approval

The research protocol and informed consent was approved by the Ethics committee of Tianjin science and technology commission (approval No TMUaMEC2017017).

Conflict of interests

The authors declare they have no competing interests.

Consent for publication

Not applicable.

Contributions

ZX: conception and design of the study, samples collection, data analysis, manuscript preparation, study design, laboratory work. TY, RC: data analysis, manuscript preparation. LC, RC: data analysis, critically reviewing the paper. TYL, GM, KN: sample collection, critically reviewing the paper. YL, WZ, NT, JS: critically reviewing the paper. HVM: conception and design of the study, data analysis, manuscript preparation, critically reviewing the paper. All authors read and approved the final manuscript.

References

- [1] Becker K, Heilmann C, Peters G. Coagulase-negative staphylococci. Clin Microbiol Rev 2014;27:870–926.
- [2] Herwaldt LA, Geiss M, Kao C, Pfaller MA. The positive predictive value of isolating coagulase-negative staphylococci from blood cultures. Clin Infect Dis 1996;22:14–20.
- [3] Stefani S, Chung DR, Lindsay JA, Friedrich AW, Kearns AM, Westh H, et al. Methicillin-resistant *Staphylococcus aureus* (MRSA): global epidemiology and harmonisation of typing methods. Int J Antimicrob Agents 2012;39:273–82.
- [4] Dong F, Ji ZB, Chen CX, Wang GZ, Wang JM. Target gene and function prediction of differentially expressed microRNAs in lactating mammary glands of dairy goats. Int J Genomics 2013;917342.
- [5] Pinho MG, de Lencastre H, Tomasz A. An acquired and a native penicillin-binding protein cooperate in building the cell wall of drug-resistant staphylococci. Proc Natl Acad Sci 2001;98:10886–91.
- [6] IWG-SCC. Classification of staphylococcal cassette chromosome mec (SCCmec): guidelines for reporting novel SCCmec elements. Antimicrob Agents Chemother 2009;53:4961–7.
- [7] Thomas JC, Vargas MR, Miragaia M, Peacock SJ, Archer GL, Enright MC. Improved multilocus sequence typing scheme for *Staphylococcus epidermidis*. J Clin Microbiol 2007;45:616–9.
- [8] Miragaia M, Couto I, Lencastre H. Genetic diversity among methicillin-resistant *Staphylococcus epidermidis* (MRSE). Microb Drug Resist 2005;11:83–93.
- [9] Albrich WC, Harbarth S. Health-care workers: source, vector, or victim of MRSA? Lancet Infect Dis 2008;8:289–301.
- [10] Widerström M, Wiström J, Edebro H, Marklund E, Backman M, Lindqvist P, et al. Colonization of patients, healthcare workers, and the environment with healthcare-associated *Staphylococcus epidermidis* genotypes in an intensive care unit: a prospective observational cohort study. Infect Dis 2016;16:743.
- [11] Xin D, Zhu Y, Yan S, Li T, Tao L, Gang S, et al. Molecular analysis of *Staphylococcus epidermidis* strains isolated from community and hospital environments in China. PLoS One 2013;8:e62742.
- [12] Xu Z, Mkrtchyan HV, Russell CA, Wang N, Cutler RR. Antibiotic resistance and *mecA* characterization of coagulase-negative staphylococci isolated from three hotels in London, UK. Front Microbiol 2015;6:947.
- [13] Mkrtchyan HV, Russell CA, Wang N, Cutler RR. Could public restrooms be an environment for bacterial resistomes? PLoS One 2013;8:e54223.
- [14] CLSI. CLSI performance standard of antimicrobial susceptibility testing: twenty-fourth international supplement. Wayne, PA: CLSI; 2014.
- [15] Kondo Y, Ito T, Ma XX, Watanabe S, Kreiswirth BN, Etienne J, et al. Combination of multiplex PCRs for staphylococcal cassette chromosome *mec* type assignment: rapid identification system for *mec*, *ccr*, and major differences in junkyard regions. Antimicrob Agents Chemother 2007;51:264–74.
- [16] Miragaia M, Thomas JC, Couto I, Enright MC, de Lencastre H. Inferring a population structure for *Staphylococcus epidermidis* from multilocus sequence typing data. J Bacteriol 2007;189:2540–52.
- [17] Cauwelier B, Gordts B, Descheemaeker P, Landuyt H. Evaluation of a disk diffusion method with cefoxitin (30 μ g) for detection of methicillin-resistant *Staphylococcus aureus*. Eur J Clin Microbiol Infect Dis 2004;23:389–92.
- [18] Shah DA, Wasim S, Abdullah FE. Comparison of oxacillin and cefoxitin for the detection of *mecA* gene to determine methicillin resistance in coagulase negative staphylococci (CoNs). J Coll Physicians Surg Pak 2017;520–2.
- [19] Li M, Wang X, Gao Q, Lu Y. Molecular characterization of *Staphylococcus epidermidis* strains isolated from a teaching hospital in Shanghai, China. J Med Microbiol 2009;58:456–61.
- [20] Xu Z, Shah HN, Misra R, Chen J, Zhang W, Liu Y, et al. The prevalence, antibiotic resistance and *mecA* characterization of coagulase negative staphylococci recovered from non-healthcare settings in London, UK. Antimicrob Resist Infect Control 2018;7:73.
- [21] Trindade PA, McCulloch JA, Oliveira GA, Mamizuka EM. Molecular techniques for MRSA typing: current issues and perspectives. Braz J Infect Dis 2003;7:32–43.
- [22] Xu Z, Misra R, Jamroz D, Paterson GK, Cutler RR, Holmes MA, et al. Whole genome sequence and comparative genomics analysis of multi-drug resistant

- environmental *Staphylococcus epidermidis* ST59. G3 2018;8:2225–30 (Bethesda).
- [23] Weiß S, Kadlec K, Feßler AT, Schwarz S. Complete sequence of a multi-resistance plasmid from a methicillin-resistant *Staphylococcus epidermidis* ST5 isolated in a small animal clinic. J Antimicrob Chemother 2014;69:847–59.
- [24] Li M, Wang X, Gao Q, Lu Y. Molecular characterization of *Staphylococcus epidermidis* strains isolated from a teaching hospital in Shanghai, China. J Med Microbiol 2009;58:456–61.
- [25] Mendes RE, Deshpande LM, Costello AJ, Farrell DJ. Molecular epidemiology of *Staphylococcus epidermidis* clinical isolates from U.S. Hospitals. Antimicrob Agents Chemother 2012;56:4656–61.
- [26] Bouchami O, Achour W, Mekni MA, Rolo J, Hassen A. Antibiotic resistance and molecular characterization of clinical isolates of methicillin-resistant coagulase-negative staphylococci isolated from bacteremic patients in oncohematology. Folia Microbiol (Praha) 2011;56:122–30.