



## **UWL REPOSITORY**

**repository.uwl.ac.uk**

Microbiological hazards associated with food products imported from the Asia-Pacific region based on analysis of the rapid alert system for food and feed (RASFF) notifications

Dada, Ayokunle Christopher, Somorin, Yinka M., Ateba, Collins N., Onyeaka, Helen, Anyogu, Amarachukwu ORCID logoORCID: <https://orcid.org/0000-0001-9652-7728>, Kasan, Nor Azman and Odeyemi, Olumide A. (2021) Microbiological hazards associated with food products imported from the Asia-Pacific region based on analysis of the rapid alert system for food and feed (RASFF) notifications. Food Control, 129. p. 108243. ISSN 0956-7135

<http://dx.doi.org/10.1016/j.foodcont.2021.108243>

**This is the Accepted Version of the final output.**

**UWL repository link:** <https://repository.uwl.ac.uk/id/eprint/8240/>

**Alternative formats:** If you require this document in an alternative format, please contact: [open.research@uwl.ac.uk](mailto:open.research@uwl.ac.uk)

**Copyright:** Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy:** If you believe that this document breaches copyright, please contact us at [open.research@uwl.ac.uk](mailto:open.research@uwl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# Food Control

## Microbiological hazards associated with food products imported from the Asia-Pacific region based on analysis of the Rapid Alert System for Food and Feed (RASFF) notifications --Manuscript Draft--

Manuscript Number:	FOODCONT-D-21-00745R1
Article Type:	Research Paper
Keywords:	Microbiological safety; Supply chain; Foodborne pathogens; food products
Corresponding Author:	Olumide Adedokun Odeyemi Mowbray, AUSTRALIA
First Author:	Olumide A Odeyemi
Order of Authors:	Olumide A Odeyemi Ayokunle C. Dada Yinka M. Somorin Collins N. Ateba Helen Onyeaka Amarachukwu Anyogu Nor Azman Kasan
Abstract:	<p>The Rapid Alert System for Food and Feeds (RASFF) is a monitoring and notification tool developed by the European Commission to serve as a central database for collating food safety-related information for member states of the European Union (EU). This study evaluated the microbiological safety of foods originating from the Asia-Pacific region in the past two decades (2000-2020) by analysing incidences and trends of notifications and alerts on the RASFF database. The highest number of notifications were for foods exported from India and Thailand as foods exported from these two countries constituted more than half (54%) of the notifications recorded on the RASFF database for the entire Asia-Pacific region from 2000-2020, compared to ANZ (Australia and New Zealand), which had very low notifications (1.2% = 23/1873). Among the 2121 notifications of pathogenic microorganisms, consisting of 14 genera, <i>Salmonella</i> was the most predominant as approximately 7 out of every 10 pathogens isolated from exported foods from the Asia-Pacific Region in the past two decades (74%, 1560/2121) were <i>Salmonella</i> species. More than 95% of pathogen species notifications for fruits and vegetables exported from India and Bangladesh were associated with betel leaves. Among the nuts, nut products and seeds, sesame seeds were the main food item contaminated by <i>Salmonella</i>, and these accounted for 87% of total <i>Salmonella</i> notifications. Across the food categories, there was a statistically significant reduction in the number of notifications associated with seafoods over the years (<math>r = -0.73</math>, <math>p=0.0001</math>). Although a statistically significant increase in notifications for fruits and vegetables (<math>r = 0.66</math>, <math>p=0.008</math>) was recorded between 2000-2014, more recent years (2015-2020) have been associated with reducing trends in the number of notifications associated with this food type (<math>r = -0.85</math>, <math>p=0.03</math>). Results indicate that imported foods may be potential vehicles for the transmission of clinically relevant microorganisms.</p>

Office of Research Services,  
University of Tasmania,  
Australia  
19<sup>th</sup> April 2021

Editor,  
Food Control  
Dear Editor,

**RE: Response to Review Comments [Ref. No.: FOODCONT-D-21-00745]**

Thank you for the review of our manuscript " Microbiological hazards associated with food products exported from the Asia-Pacific region based on analysis of the Rapid Alert System for Food and Feed (RASFF) notifications [Ref. No.: FOODCONT-D-21-00745]".

All the editor's and reviewer's comments have now been addressed. Kindly find tabulated below point-by-point responses to each comment. Locations in the journals where changes have been effected are also stated, with changes in the manuscript set as tracked change, as per journal specification.

We thank you in advance for your time in considering this work for publication and look forward to hearing your response.

Yours sincerely,

Dr Olumide Odeyemi

## Highlights

- RASFF pathogen notifications analysed for Asia Pacific food imports (2000 – 2020).
- 1873 notifications reported over 2 decades across 29 notifying countries.
- Imported foods can potentially transmit clinically relevant microorganisms.
- *Salmonella* was the predominant pathogen (7 out of every 10 pathogens).
- Inconsistent actions in responses to notifications by different EU member states

**Response to Reviewers' comments:**

<b>Reviewer #1</b>		
<b>No</b>	<b>Comment</b>	<b>Response</b>
1	In this study, the authors evaluated the microbiological safety of foods being imported from the Asia-Pacific region from 2000 to 2020 by analyzing incidences and trends of notifications and alerts in the Rapid Alert System for Food and Feeds database. Overall, this is well-organized research article with some interesting findings.	Thank you
2	The scientific names of pathogens in latin should be in italic, no matter they are in the manuscript or the supplementary materials.	In agreement with the reviewer, issues with the pathogen names in the manuscript relate to the supplementary tables. We have now gone through each of the supplementary tables and have corrected the anomalies in the description of pathogen names. The names are now consistent with acceptable norms in literature (e.g. Jiang et al 2021, Food Control, <a href="https://doi.org/10.1016/j.foodcont.2020.107535">https://doi.org/10.1016/j.foodcont.2020.107535</a> and Bangtrakulnonth et al 2004, Emerging infectious diseases <a href="https://doi.org/10.3201/eid1001.02-0781">https://doi.org/10.3201/eid1001.02-0781</a> ), e.g. <i>Salmonella enterica</i> ser. Derby, <i>Salmonella</i> Typhimurium etc
3	It is confusing both "export" and "import" were used. It is more proper to use "import" since this study was conducted from the perspective of who imported foods from the Asia-Pacific region.	In agreement with the reviewer, issues with import/export clarifications have now been made. Given these changes, the manuscript now reflects that it was conducted from the perspective of who imported foods from the Asia-Pacific region. These changes are reflected in Lines 1,34-42,189,201,226,240, 308,369, 381,383,395,404, 412-414, 447-456, 482-484, 496-502,553,566, 583, 641, 667 and 714
<b>Reviewer #2</b>		
5	The MS entitled "Microbiological hazards associated with food products exported from the Asia-Pacific region based on analysis of the Rapid Alert System for Food and Feed (RASFF) notifications" describes microbial hazards associated with exported food items. The subject of the MS is relevant and this type of scientific investigation helps countries to implement policies to improve export quality. Authors have collected data for last 20 years which is significant amount of data analysis.	Thank you

6	Authors may want to break this data into 5 years slots such as 2000-2004, 2005-2009, and so on to see if there is any improvement in number of notifications reported. Alternatively, authors may want to consider analyzing this data over time to see any if number of notifications have gone up or down over 20 years.	Actually, this has already been done in the manuscript and results presented in Table 3. The data was split into distant years (2000-2014) versus most recent years (2015-2020) and assessed for trends in the different data timeframes (i.e. does the pathogen notification increase with years considering the entire period (2000-2020), most recent years (2015-2020) or distant years (2000-2014). However, we agree with the reviewer that this statement does not come out clearly in the methodology section so we have now added this into the methodology section.
<b>Reviewer #3</b>		
7	Table 2, it would be better to provide information on the total number of foods imported or sample for detection from each country or region.	It is difficult to get data on total number of foods imported or sample for detection from each country or region. One limitation of the RASSF database is that it does not record the number of detection per given number of food items tested. We have now included this as a limitation in the study and suggestion for RASSF database operators to consider including in the future (Line 713-716)
8	In the results part, brief analysis of observed results should be provided. More specific analysis should be conducted in the Discussion. For example, the author mentioned that "The highest number of notifications was made by the UK (28%)", why?.	Analysis and discussion of the results is already presented in the discussion section and would be repetitive if these analysis are presented again in the results section. However, as the reviewer as advised, we have now included more specific analysis....in the discussion section e.g. Line 519-527, 499-503, 594-597)
9	The significance of the research should be provided in both abstract and discussion section	This has now been added into the abstract (line 52-53...) and discussion (line 488-490).
<b>Editor</b>		
11	The highlights do not conform to the journal's requirements: maximum 85 characters, including spaces, per bullet point. Please revise.	The highlights have now been edited to conform with the journal's standard.

# Microbiological hazards associated with food products ~~imported~~~~exported~~ from the Asia-Pacific region based on analysis of the Rapid Alert System for Food and Feed (RASFF) notifications

Ayokunle C. Dada<sup>\*1</sup>, Yinka M. Somorin<sup>2</sup>, Collins N. Ateba<sup>3</sup>, Helen Onyeaka<sup>4</sup>, Amarachukwu Anyogu<sup>5</sup>, Nor Azman Kasan<sup>6</sup>,  
Olumide A. Odeyemi<sup>\*7</sup>

<sup>1</sup>QMRA Data Experts, P.O. Box 37, Waikato Mail Lobby, Hamilton 3204, New Zealand

<sup>2</sup>School of Pharmacy, Queen's University Belfast, Belfast, United Kingdom

<sup>3</sup>Food Security and Safety Niche Area, Faculty of Natural and Agricultural Sciences, North-West University, Mmabatho, South Africa

<sup>4</sup>Department of Chemical Engineering, University of Birmingham, Edgbaston, United Kingdom

<sup>5</sup>Food, Nutrition and Public Health, School of Life Sciences, University of Westminster, London, United Kingdom

<sup>6</sup>Higher Institution Centre of Excellence (HiCoE) Institute of Tropical Aquaculture and Fisheries, Universiti Malaysia Terengganu, 21030 Kuala Nerus, Terengganu, Malaysia.

<sup>7</sup>Office of Research Services, Research Division, University of Tasmania, Launceston, Australia

\*Corresponding ~~Author~~~~authors~~: [chris@qmradata.com](mailto:chris@qmradata.com), [olumide.odeyemi@utas.edu.au](mailto:olumide.odeyemi@utas.edu.au)

## Abstract

The Rapid Alert System for Food and Feeds (RASFF) is a monitoring and notification tool ~~and food safety-related database~~ developed by the European Commission ~~to serve as a central database for collating food safety-related information for member states of the European Union (EU)~~. This study evaluated the microbiological safety of foods originating from the Asia-Pacific region in the past two decades (2000-2020) by analysing incidences and trends of notifications and alerts on the RASFF database. The highest number of notifications were for foods ~~imported~~~~exported~~ from India and Thailand as foods ~~imported~~~~exported~~ from these two countries constituted more than half (54%) of the notifications recorded on the RASFF database for the entire Asia-Pacific region from 2000-2020, compared to ANZ (Australia and New Zealand), which had very low notifications (1.2% = 23/1873). Among the 2121 notifications of pathogenic microorganisms, consisting of 14 genera, *Salmonella* was the most predominant as approximately 7 out of every 10 pathogens isolated from ~~exported~~ foods ~~imported~~ from the Asia-Pacific Region in the past two decades (74%, 1560/2121) were *Salmonella* species. More than 95% of pathogen species notifications for fruits and vegetables ~~imported~~~~exported~~ from India and Bangladesh were associated with betel leaves. Among the nuts, nut products and seeds, sesame seeds were the main food item contaminated by *Salmonella*, and these accounted for 87% of total *Salmonella* notifications. Across the food categories, there was a statistically significant reduction in the number of notifications associated with seafoods over the years ( $r = -0.73$ ,  $p=0.0001$ ). Although a statistically significant increase in notifications for fruits and vegetables ( $r = 0.66$ ,  $p=0.008$ ) was recorded between 2000-2014, more recent years (2015-2020) have been associated with reducing trends in the number of notifications associated with this food type ( $r = -0.85$ ,  $p=0.03$ ). Results indicate that imported foods may be potential vehicles for the transmission of clinically relevant microorganisms. ~~Studies of this nature can potentially encourage -countries to implement policies that improve export quality.~~

54  
55 **Keywords:** Microbiological safety, supply chain, foodborne pathogens, food products  
56  
57  
58  
59  
60



## 1. Introduction

Food safety is everyone's business, and it is pivotal to public health protection. Many factors make food unsafe for human and animal consumption, ranging from chemical contaminants such as pesticides, and mercury in seafood, to biological hazards, including parasites and microorganisms (2010). It is imperative to maintain consumers' health by ensuring optimum standards for food from production/harvest through the food supply chain up to the point of consumption. However, the impact of globalization, leading to a long food supply chain coupled with the rapid distribution of foods across the world, has made it difficult to track the origin and history of most foods and/or food additives resulting in food spoilage, loss and waste, as well as foodborne disease outbreaks from the consumption of unsafe food (Rezaei & Liu, 2017). The major causes of foodborne illnesses include; bacteria (*Salmonella*, *Campylobacter*, *Listeria*, *Vibrio cholerae*, enterohaemorrhagic *Escherichia coli*), viruses (Norovirus, Hepatitis A), parasites (*Echinococcus* spp, *Taenia solium*, *Ascaris*, *Cryptosporidium*, *Entamoeba histolytica*, *Giardia*), prions, fungi, and chemical agents (naturally occurring toxins such as mycotoxins and phycotoxins, persistent organic pollutants such as dioxins and polychlorinated biphenyls, heavy metals such as lead, mercury and cadmium) (World Health Organization, 2020a, 2020b). Over the past few years, there has been a significant increase in the number of documented cases of foodborne diseases reported by various countries across the globe. Worldwide, the number is estimated to be 600 million cases of foodborne illnesses annually, resulting in 420,000 deaths, with about 30% of the deaths recorded in children (Food and Agriculture Organization, 2019). In the Asia-Pacific region, which comprises countries within South, East, Southeast Asia and Oceania, the number of foodborne illnesses continues to rise, with current estimates pegged at 275 million cases annually (Food and Agriculture Organization, 2019). The Asia-Pacific region has a population of about 4.6 billion people, accounting for about 60% of the global population (Worldometer, 2021). This large, growing population places a lot of demand on water, energy and food resources. Furthermore, although the region is rapidly becoming one of the largest economies globally, over 40% of the population live at or below the poverty line (ESCAP, 2020). The high poverty rate coupled with the growing population and other factors, including poor hygiene, lack of adequate food processing and preservation routines and poor food policy enforcement, can be considered the main drivers of foodborne disease outbreaks within the region (Food and Agriculture Organization, 2019).

A large proportion of foodborne disease outbreaks are not noticed immediately, traced to the consumption of a particular food item, or even recognised as a foodborne disease outbreak. Others go unreported and are not investigated. Furthermore, there are limitations in the capacity of many countries to undertake food surveillance due to inefficient food regulatory agencies and unavailability of reference laboratories and analytical tools/expertise to identify sources of foodborne disease outbreaks and conduct investigations (Wang, et al., 2016). The Rapid Alert System for Food and Feeds (RASFF), created in 1979, is a monitoring and notification tool developed by the European Commission to serve as a central database for collating food safety-related information for member states of the European Union (EU). It allows the food regulatory bodies of member countries to report and submit information and notifications about unsafe foods and unauthorised/illegal foods/food additives in circulation within the market, ensuring the rapid transmission and sharing of information regarding food safety risks as they occur in real time. This allows actions, including product recalls/withdrawals, to prevent adverse health and economic consequences to consumers within the European Union and beyond (European Commission, 2020a). RASFF features a consumer portal that is interactive and has been available to consumers since 2014. Its interactive online database can be used for customised searches of food safety-related information using criteria including Notification, Type, Hazard, Date, Product and Keywords. In addition to monitoring food safety, RASFF is also used to track food fraud, substitutions, and other economically motivated adulterations (EMAs), which may not be considered direct food hazards, but are still of public health and economic significance (Tähkääpää, Majjala,

Korkeala, & Nevas, 2015). The system operates on a 24-hour basis, allowing for a seamless flow of information and ensuring regulatory compliance. Thus, the RASFF serves the very important function of mitigating risks associated with foods and continuously ensuring public health safety (Kowalska & Manning, 2020).

The RASFF has recently been used in tracing the origin of food contaminations and in foodborne outbreak investigations such as the *Salmonella* Poona infections reported in France in 2019 and the multi-country outbreak of *Listeria monocytogenes* in cold-smoked fish products in five member EU states of Denmark, Estonia, Finland, France and Sweden (European Commission, 2020b). Similarly, there have been RASFF notifications from countries within the Asia-Pacific region, including Australia, Hong Kong, Japan and Nepal, amongst others (European Commission, 2020a). It is imperative that food exporters, especially those exporting from developing countries to the more developed countries in Europe, meet the food standards of the destination country. The presence of microbiological and chemical hazards in foods destined for Europe, when flagged by RASFF, can lead to import rejection with attendant economic losses (Somorin, Odeyemi, & Ateba, 2021). The European Food Safety Authority (EFSA) and the European Center for Disease Prevention and Control (ECDC) have identified *Salmonella enteritidis*, *Campylobacter* spp., shiga-toxin producing *Escherichia coli*, *Yersinia* spp. and *Listeria monocytogenes* as the major bacteria species implicated in human cases of foodborne illnesses (European Food Safety Authority, 2021) and are actively monitored for by the RASFF. These species have previously been identified in foods of Asian-Pacific origin (Food and Agricultural Organisation, 2018; Kaakoush, Castaño-Rodríguez, Mitchell, & Man, 2015; Rivas, Strydom, Paine, Wang, & Wright, 2021; Sodagari, Wang, Robertson, Habib, & Sahibzada, 2020; Sugiri, et al., 2014), highlighting the need for deliberate measures to eliminate these pathogens from foods, especially those destined for export to Europe.

Although systems such as the RASFF are in operation worldwide in both the public and private sector to assure food safety, there continue to be incidences of consumer foodborne illnesses that are directly related to lapses in food regulations, inadequate monitoring and enforcement or the deliberate attempts by Food Business Operators (FBOs) to increase profits through opportunistic malpractices (Tähtkääpää, et al., 2015). Furthermore, the Covid-19 pandemic, with its disruption of the food supply chain both domestically and internationally, has worsened the food security risks, especially in the Asia-Pacific region. The pandemic undermined food accessibility and availability, while the global lockdown of 2020 disrupted food production, distribution operations, logistics and trading (Kim, Kim, & Park, 2020). Also, the disruption in the food supply chain and food production and trade led to an increase in food spoilage rates due to longer wait times, thus increasing the risks of foodborne illnesses and economically motivated fraud (Kim, et al., 2020). Therefore, it is evident that continued vigilance is required to assure food safety, which continues to be a challenge globally, especially regarding the presence of microorganisms in foods. This article evaluates the microbiological safety of foods originating from the Asia-Pacific region by analysing incidences and trends of notifications and alerts on the RASFF database.

## 2. Materials and Methods

### 2.1. RASFF database

Six sections (Notification, Type, Hazard, Date, Product and Keywords), presented as fill-in forms, constitute the RASFF portal search page. In the "Notification" section, users can use four sub-sections to enter details related to a food item for which notification information is being sought. These sub-sections include 1) 'Reference' – a unique reference number associated with each notification; 2) 'Subject' – a field wherein specific subjects can be searched using the "AND","OR" Boolean operators; 3) 'Notified by' – a drop-down list of notifying countries (EU member states); and 4) 'Open alert' – a field where users can select for information that includes or excludes open alerts. The "Type" section consists of three subsections; 1) 'Type'- allows users to select their preference for information related to food, feed or food contact material; 2) 'Classification'- where users can select the notification classification; and 3) 'Basis'- where users can select the notification basis for which information is being sought. The 'Date' section allows users to select dates or periods of the year for which notification data is required. The 'Product' section consists of four subsections; 'Category' (a drop-down list of types of food included in the RASFF database); 'Country' (a drop-down list of all notifying and originating countries); 'action taken' (a drop-down list of action taken following notification); and 'flagged as' (with two sublevels – distribution and origin). The 'Keyword' section allows users to search by keywords.

### 2.2 Data collection

The dataset used for this study was obtained from the RASFF database as described by Yinka M. Somorin, Olumide A. Odeyemi, and Collins N. Ateba (2021) with slight modifications based on the three search criteria that focused on the product country of origin, product type and hazard category. In terms of the product's country of origin, the database was searched for information relating to all Asia-Pacific countries. Notification data was only available for imports from 20 exporting countries in the Asia-Pacific region: Afghanistan, Australia, Bangladesh, Cambodia, China, Hong Kong, India, Indonesia, Laos, Malaysia, Nepal, New Zealand, Pakistan, Philippines, Singapore, South Korea, Sri Lanka, Taiwan, Thailand and Vietnam. In terms of product type, the search was restricted to food only, excluding feed and food contact materials. In terms of hazard category, the focus was placed on "pathogenic microorganisms" included in the RASFF database. The initial notification date was not restricted to enable complete capturing of all the available notifications; however, considering the paucity of data preceding the year 2000, notification data preceding 2000 were subsequently excluded from the dataset. The dataset used therefore spans two decades.

### 2.2. Data analysis

The notifications obtained for the imports from 20 exporting countries in the Asia-Pacific region were exported as an Excel file. Data sorting, filtering, and formatting were completed using Microsoft Excel spreadsheets. Missing data instances were replaced with the term "not specified". Descriptive statistics were collated to determine the predominant pathogens associated with seafood originating from countries in the Asia-Pacific region (the most contaminated food category); period of lowest and highest notifications; number of notifying countries; and the country of origin. To determine trends in the yearly notifications, Pearson correlations of year versus number of notifications per sub-region and per food category were conducted using IBM® Statistical Package for the Social Sciences (SPSS®) software version 24. The data was split into distant years (2000-2014) versus most recent years (2015-2020) and assessed for trends in the different data timeframes (i.e., does the pathogen notification increase with years considering the entire period (2000-2020), most recent years (2015-2020) or distant years (2000-2014). A statistically significant increasing trend of notifications was

reported when the Pearson correlation statistics (r) were >0.5 with an associated p-value <0.05. A statistically significant decreasing trend of notifications was reported when the Pearson correlation statistics (r) were < -0.5 with an associated p-value <0.05. A map of originating Asia-Pacific countries was processed in Tableau, and Chord diagrams were generated using Rstudio, a free and open-source environment for the statistical language R (<https://www.r-project.org>). A Sankey Diagram was developed using Python to display notification flows and their quantities in proportion to one another, with the originating country designated as the source node, while the product type and action taken were designated as target nodes.


### 3. Results

#### 3.1. Notifications - originating and notifying countries.

A total of 1873 notifications from 2000 to 2020 were obtained, based on our search criteria, for food items imported from 20 countries in the exporting countries in the Asia-Pacific region: Afghanistan, Australia, Bangladesh, Cambodia, China, Hong Kong, India, Indonesia, Laos, Malaysia, Nepal, New Zealand, Pakistan, Philippines, Singapore, South Korea, Sri Lanka, Taiwan, Thailand and Vietnam (**Fig.1**).


#### 3.2 Notification categories

**Presented in Table 1** are three levels of notification categories associated with the RASFF database (i.e., Notification basis, Notification type and Action taken). was observed in this study. In more than half of the notifications (51%), the basis of notification was border control, with the consignment detained (**Table 1**). Only 1% of notifications were due to food poisoning and < 1% due to consumer complaints. The notification type was mainly border rejection (43%) while the most prevalent action taken was re-dispatching (17.6%) and destruction of the consignments (17.5%, **Table 1**). The highest number of notifications were for foods imported~~exported~~ from India (31% = 588/1873), followed by Thailand (23% = 428/1873, **Table 2**). The categories of foods categories originating from Asia-Pacific Region (n=1873) by country of origin from 2020 – 2020 are shown in **Fig. 2**. Together, foods imported~~exported~~ from these two countries constituted more than half (54%) of the notifications recorded on the RASFF database for the entire Asia-Pacific region from 2000-2020 (**Fig 3**), compared to ANZ (Australia and New Zealand), which had very low notifications (1.2% = 23/1873). Also, the prevalent actions taken associated with notifications from India and Thailand were re-dispatching and destruction of the consignments (**Fig. 4**).




**Fig. 1:** Map of originating country and number of notifications listed in the *RASFF* database

[N.B. Additional countries not on the map: Hong Kong (Notifications: 6, % of total notifications: 0.32%) and Singapore (Notifications: 3, % of total notifications: 0.16%)]




**Fig. 2** Food categories of RASFF notifications (n=1873) originating from Asia-Pacific Region (n=1873) by country of origin from 2020 - 2020. Meat includes meat and meat products (other than poultry), poultry meat and poultry meat products. Seafoods include bivalve molluscs and products thereof, cephalopods and products thereof, crustaceans and products thereof, fish and fish products. Other foods include prepared dishes and snacks, soups, broths, sauces and condiments, cereals and bakery products, cocoa and cocoa preparations, coffee and tea, confectionery, eggs and egg products, food additives and flavourings, milk and milk products and other food product / mixed dietetic foods, food supplements, fortified foods. ANZ= Australia and New Zealand, IDN = Indonesia, MYS=Malaysia. Other\_O = Other originating countries, i.e., Pakistan, Philippines, Hong Kong, Taiwan, Sri Lanka, Singapore, South Korea, Afghanistan, Chile, Nepal and Cambodia. The length of the arc on the circumference of the circle represents the number of notifications in each food category.



**Fig. 3.** Types of RASFF notifications (n=1873) originating from Asia-Pacific Region by country of origin from 2020 - 2020. Inform\_ATTN= Information for attention, Inform\_FLP = Information for follow-up. ANZ= Australia and New Zealand, IDN = Indonesia, MYS=Malaysia. Other\_O = Other originating countries, i.e., Pakistan, Philippines, Hong Kong, Taiwan, Sri Lanka, Singapore, South Korea, Afghanistan, Chile, Nepal and Cambodia. The length of the arc on the circumference of the circle represents the number of notifications in each food category.





**Fig. 4.** Sankey diagram of originating country versus food type and action taken.

N.B. The width of the arrows or lines show their magnitudes, so the bigger the arrow, the larger the flow (i.e., the number of pathogen notifications associated with that source-target node). Colours have been used to divide the diagram into different categories in a way that shows different source-target node connections.



298 **Table 1**

299 RASFF notifications of pathogenic microorganisms in seafood originating from the Asia-  
300 Pacific Region (n=1873) from 2020 – 2020 based on notification category.

Classification category	Notifications	
	Number	Percentage (%)
<b>Notification basis</b>		
border control - consignment detained	949	50.67%
official control on the market	349	18.63%
border control - consignment released	239	12.76%
not specified	216	11.53%
company's own check	70	3.74%
food poisoning	24	1.28%
border control - consignment under customs	17	0.91%
consumer complaint	8	0.43%
<b>Notification type</b>		
border rejection	810	43.25%
information	509	27.18%
information for attention	283	15.11%
alert	246	13.13%
information for follow-up	25	1.33%
<b>Action taken</b>		
re-dispatch	329	17.57%
destruction	327	17.46%
import not authorised	275	14.68%
official detention	130	6.94%
not specified	104	5.55%
withdrawal from the market	97	5.18%
informing recipient(s)	83	4.43%
informing authorities	76	4.06%
recall from consumers	67	3.58%
product recall or withdrawal	56	2.99%
no action taken	51	2.72%
no stock left	43	2.30%
prohibition to trade - sales ban	37	1.98%
re-dispatch or destruction	29	1.55%
return to consignor	27	1.44%
placed under customs seals	21	1.12%
physical/chemical treatment	21	1.12%
detained by operator	19	1.01%
reinforced checking	16	0.85%
withdrawal from recipient(s)	13	0.69%
public warning - press release	11	0.59%
Other actions	41	2.19%

301 Other actions: Screening sample, Product destruction or return after official permission,  
302 physical treatment - heat treatment, relabelling, Product recalled / Product to be destroyed,  
303 destination of the product identified, informing consignor and seizure.

304 **Table 2**

305 RASFF notifications of pathogenic microorganisms in foods originating from the Asia-Pacific Region (n=1873) from 2000 - 2020 based on  
306 notifying country.

Notifying country	Country of origin										Grand Total
	India	Thailand	Vietnam	Bangladesh	China	Malaysia	Indonesia	Laos	ANZ	Other_O	
United Kingdom	216	80	5	150	28	11	3	16	2	19	<b>528</b>
Italy	51	17	29	1	22	58	27	0	3	16	<b>221</b>
Netherlands	29	92	35	2	10	1	1	6	1	3	<b>179</b>
Norway	27	54	36	19	6	1	9	1	8	15	<b>168</b>
Germany	46	30	39	1	9	3	2	1	3	14	<b>145</b>
Finland	13	72	7	4	10	1	0	0	1	8	<b>115</b>
Spain	15	2	30	0	18	8	1	0	0	0	<b>74</b>
Greece	64	0	2	1	2	0	3	0	0	1	<b>73</b>
Sweden	6	26	9	1	10	0	1	0	1	7	<b>60</b>
Poland	48	0	6	0	2	0	0	0	0	0	<b>56</b>
France	16	5	5	8	5	2	7	0	3	6	<b>54</b>
Denmark	3	27	10	1	8	1	1	0	0	2	<b>53</b>
Portugal	1	0	26	0	0	2	1	0	0	1	<b>31</b>
Belgium	4	8	4	2	3	1	0	0	0	2	<b>24</b>
Other_N	49	15	6	1	7	0	8	0	1	6	<b>92</b>
<b>Grand Total</b>	<b>588</b>	<b>428</b>	<b>249</b>	<b>191</b>	<b>140</b>	<b>89</b>	<b>64</b>	<b>24</b>	<b>23</b>	<b>100</b>	<b>1873</b>

307

308 **Other\_N: Other notifying countries:** Cyprus, Austria, Slovenia, Ireland, Lithuania, Iceland, Romania, Switzerland, Bulgaria, Croatia, Estonia,  
309 Hungary, Latvia, Luxembourg, Slovakia, Sri Lanka

310


311 **ANZ=** Australia and New Zealand. **Other\_O: Other originating countries:** Cambodia, Pakistan, Phillippines, Hong Kong, Taiwan, Singapore,  
312 South Korea, Afghanistan, Nepal.

313

During the two decades studied, very few notifications were recorded for foods ~~imported~~~~exported~~ from Cambodia, Pakistan, Philippines, Hong Kong, Taiwan, Singapore, South Korea, Afghanistan, and Nepal (Table 2). Notifications were recorded by a total of 29 notifying countries. Among these, the highest number of notifications were from the United Kingdom (28% = 528/1873), Italy (12% = 221/1873) and the Netherlands (~10% = 179/1873), while the least number of notifications were reported from Cyprus, Austria, Slovenia, Ireland, Lithuania, Iceland, Romania, Switzerland, Bulgaria, Croatia, Estonia, Hungary, Latvia, Luxembourg, Slovakia and Sri Lanka (a total of only 92 notifications were reported in these 16 countries over the two decades studied, Table 2).

### 3.3. Trends in RASFF notifications of pathogenic microorganisms across sub-regions and food categories

Fig. 5 presents RASFF notifications of pathogenic microorganisms in food originating from the Asia-Pacific Region (n=1873) from 2000–2020, based on the notifying year. In the two decades considered, notifications were greatest in 2015 and 2011. Consistently over the years, food originating from South Asia was associated with the highest number of notifications while the least notifications were reported for Oceania (Fig. 5). Over the entire two decades of notifications data recorded in the RASFF database, there was no observable trend across sub-regions, with the notable exception of Oceania. Pearson correlation of year versus RASFF notifications of pathogenic microorganisms in food sub-regions indicates that notifications from Oceania have significantly reduced over the years ( $r=-0.68$ ,  $p=0.006$ ). In more recent years, however, (2015-2020), notifications from the sub-regions have reduced significantly, with the notable exception of East Asia (Table 3a). Across the food categories, there has been a statistically significant reduction in the number of notifications associated with seafoods over the years ( $r = -0.73$ ,  $p=0.0001$ , Table 3b). In contrast, there has been a statistically significant increase in the number of notifications associated with nuts, nut products and seeds over the years ( $r = 0.57$ ,  $p=0.007$ , Table 3b). Although a statistically significant increase in notifications for fruits and vegetables ( $r = 0.66$ ,  $p=0.008$ , Table 3b) was recorded between 2000-2014, more recent years (2015-2020) have been associated with reducing trends in the number of notifications associated with this food type ( $r = -0.85$ ,  $p=0.03$ , Table 3b).



**Fig. 5.** RASFF notifications of pathogenic microorganisms in food originating from Asia-Pacific Region (n=1873) from 2000 - 2020 based on notifying year.

348 **Table 3**

349 Pearson correlation of year versus RASFF notifications of pathogenic microorganisms in various (a) Asia-Pacific sub-regions and (b) food  
350 categories.

**a) Yearly notification versus sub-region**

Period	Statistics	East Asia	Oceania	South Asia	Southeast Asia	Grand Total
2000-2020	Pearson Correlation	.486*	-.676**	.452*	-0.188	0.353
	Sig. (2-tailed)	0.025	0.006	0.040	0.415	0.116
	N	21	15	21	21	21
2000-2014	Pearson Correlation	0.205	-.731*	.653**	0.025	.562*
	Sig. (2-tailed)	0.464	0.011	0.008	0.929	0.029
	N	15	11	15	15	15
2015-2020	Pearson Correlation	.821*	. <sup>b</sup>	-.873*	-.957**	-.918**
	Sig. (2-tailed)	0.045		0.023	0.003	0.010
	N	6	4	6	6	6

**b) Yearly notification versus food category**

Period	Statistics	nuts, nut products and seeds	seafoods	fruits and vegetables	herbs and spices	meat	others	Grand Total
2000-2020	Pearson Correlation	.567**	-.725**	0.387	0.004	0.354	0.419	0.294
	Sig. (2-tailed)	0.007	0.000	0.083	0.988	0.115	0.059	0.197
	N	21	21	21	19	21	21	21
2000-2014	Pearson Correlation	.655**	-.727**	.658**	0.187	-0.258	0.496	0.483
	Sig. (2-tailed)	0.008	0.002	0.008	0.541	0.353	0.060	0.068
	N	15	15	15	13	15	15	15
2015-2020	Pearson Correlation	-0.681	-0.709	-.849*	-0.332	-.878*	-0.427	-.915*
	Sig. (2-tailed)	0.136	0.115	0.032	0.520	0.021	0.399	0.011
	N	6	6	6	6	6	6	6

\*. Correlation is significant at the 0.05 level (2-tailed). \*\*. Correlation is significant at the 0.01 level (2-tailed). b- too few samples, blue cells indicate a downward trend, and red cells indicate an upward trend

#### 3.4. Pathogens associated with notifications

Although the total number of notifications was 1873, a higher total number of pathogens (n=2121) was recorded because of co-occurrence of pathogens in some food products (**Table 4**). Among the 29 notifying countries, more than a quarter (28%) of pathogen notifications for foods originating from the Asia-Pacific Region were reported by the United Kingdom (28%, i.e., 587/2121, see **Supplementary Table 1**). Cumulatively, the United Kingdom, Italy, Norway, the Netherlands, Germany and Finland accounted for more than 70% (i.e., 1556/2121) of pathogen notifications in the two-decade study period (**Supplementary Table 1**). The least number of pathogen notifications was observed for France, Poland, Portugal, Belgium, Cyprus, Austria, Slovenia, Ireland, Lithuania, Iceland, Romania, Switzerland, Bulgaria, Croatia, Estonia, Luxembourg, Hungary, Latvia and Slovakia, as these 19 countries accounted for less than 15% (i.e., 277/2121) of pathogen notifications in the two-decade study period (**Supplementary Table 1**).

Among the 2121 notifications of pathogenic microorganisms from 2000-2020, the following 14 genera consisting of *Salmonella*, *Vibrio*, *Bacillus*, *Escherichia*, *Norovirus*, *Clostridium*, *Campylobacter*, *Staphylococcus*, *Enterococcus*, *Hepatitis*, *Pseudomonas*, *Cronobacter*, and *Plesiomonas*, were observed in this study (**Table 4**). Among these, the least commonly isolated pathogen genera were observed to be *Cronobacter*, Fungi (moulds - no specific genus was identified), *Plesiomonas*, *Shigella*, *Yersinia* and *Streptococci* (**Table 4**). Conversely, *Salmonella* was the most predominant as approximately 7 out of every 10 pathogens isolated from ~~foods imported from exported foods from~~ the Asia-Pacific Region in the past two decades (74%, 1560/2121) were *Salmonella* species (**Table 4**). Foods exported from India had the highest pathogen notifications (642/2121, **Table 4**).

378 **Table 4**

379 RASFF notifications of pathogenic microorganisms in food originating from the Asia-Pacific Region (n=1873) from 2000 - 2020 based on  
380 originating country.

Pathogen genus	Country of origin										Grand Total (%)
	India	Thailand	Vietnam	Bangladesh	China	Malaysia	Indonesia	Laos	ANZ	Others	
<i>Salmonella</i>	551	411	182	165	69	30	38	30	23	61	1560(73.55%)
<i>Vibrio</i>	61	40	59	35	12	62	31	0	1	8	309(14.57%)
<i>Bacillus</i>	16	5	14	2	42	0	3	0	0	10	92(4.34%)
<i>Escherichia</i>	3	16	5	0	2	4	0	15	0	2	47(2.22%)
Norovirus	0	0	30	0	11	0	0	0	0	1	42(1.98%)
Not specified	6	2	5	0	4	0	0	0	0	11	28(1.32%)
<i>Clostridium</i>	3	5	2	0	12	0	0	0	0	0	22(1.04%)
<i>Campylobacter</i>	0	5	1	0	0	0	0	0	0	0	6(0.28%)
<i>Staphylococcus</i>	0	1	2	0	0	0	0	0	0	0	3(0.14%)
<i>Enterococcus</i>	1	0	0	0	0	0	1	0	0	0	2(0.09%)
Hepatitis	0	0	0	0	1	0	0	0	1	0	2(0.09%)
<i>Pseudomonas</i>	1	0	0	0	0	0	0	0	0	1	2(0.09%)
<i>Cronobacter</i>	0	0	0	0	1	0	0	0	0	0	1(0.05%)
Fungi	0	0	0	0	1	0	0	0	0	0	1(0.05%)
<i>Plesiomonas</i>	0	0	0	0	0	0	1	0	0	0	1(0.05%)
<i>Shigella</i>	0	1	0	0	0	0	0	0	0	0	1(0.05%)
<i>Yersinia</i>	0	1	0	0	0	0	0	0	0	0	1(0.05%)
<i>Streptococci</i>	0	0	0	0	1	0	0	0	0	0	1(0.05%)


381

382 **Other\_N: Other notifying countries:** Cyprus, Austria, Slovenia, Ireland, Lithuania, Iceland, Romania, Switzerland, Bulgaria, Croatia, Estonia,  
383 Hungary, Latvia, Luxembourg, Slovakia, Sri Lanka. **Other\_O: Other originating countries:** Australia, Cambodia, New Zealand, Pakistan,  
384 Philippines, Hong Kong, Taiwan, Singapore, South Korea, Afghanistan, Nepal.

385 The total number of pathogens (n=2119) exceeds the total number of notifications (n=1873) because of the co-occurrence of pathogens.

Among the pathogens identified in **imported/exported** foods originating from the Asia-Pacific Region, India also had the highest *Salmonella* notifications (551/1560). On the whole, more than 80% (i.e. 1309/1560) of *Salmonella* notifications were from foods **imported/exported** from India, Thailand, Vietnam and Bangladesh. Forty percent (i.e., 625/1560) of the time, pathogen identification did not exceed the genus level for RASFF notifications of *Salmonella* in food originating from the Asia-Pacific Region during the two-decade study period (**Table 5**). Hence in these instances, the species or serovar implicated in the *Salmonella* notification was not identified. In the remaining 935 instances of *Salmonella* notifications in food originating from the Asia-Pacific Region, a total of 178 serovars of *Salmonella enterica* were identified (**Table 5**). The top 10 most frequently isolated serovars in the RASFF *Salmonella* notifications are serovars Weltevreden, Enteritidis, Hvittingfoss, Agona, Stanley, Senftenberg, Augustenborg, Brunei, Typhimurium, Lexington, Newport and Mbandaka (**Table 5**).

Apart from *Salmonella*, *Vibrio* was the next most common, albeit at a reported frequency far lower than that of *Salmonella* (14.6% = 309/1562). The most commonly isolated *Vibrio* in food products **imported/exported** from the Asia-Pacific Region from 2000 – 2020 were *V. cholerae* (46.6%, i.e., 144/309) and *V. parahaemolyticus* (46%, i.e., 142/309) while the least isolated *Vibrio* were *V. fluvialis* and *V. mimicus* (0.32% each, i.e., 1/309) (**Fig. 6**).



**Fig. 6.** Species identified in 309 RASFF notifications of *Vibrio* in food originating from Asia-Pacific Region from 2000 - 2020.


Apart from hepatitis A and norovirus, viruses were generally not detected in the foods **imported/exported** from the Asia-Pacific region (**Table 4**). When Hepatitis A was reported in food items, both instances were in 2012 when frozen green mussels (*Perna canaliculus*) from



412 New Zealand and frozen strawberry cubes from China tested positive for the virus. The action  
 413 taken in each instance was different. In the case of contaminated frozen green mussels from  
 414 New Zealand, the notifying country (Italy) withdrew the product from the market. In  
 415 contaminated frozen strawberry cubes from China, the notifying country (Belgium) only  
 416 informed recipients. Norovirus detection in foods imported from the Asia-Pacific region was  
 417 reported 42 times in the two-decade study period (**Table 4**). Eleven of these notifications  
 418 were from raspberries, seaweed salad and strawberries ~~imported~~~~exported~~ by China and thirty  
 419 other instances were from other fruits ~~imported~~~~exported~~ by Vietnam between 2011 and 2019.  
 420 In 6 out of the 11 norovirus notifications for fruits and vegetables from China, notifications were  
 421 informed by the occurrence of ongoing food poisoning or consumer complaints.  
 422 There was no consistency in response to ~~N~~orovirus notifications as the action taken varied  
 423 between countries. For instance, actions taken included rejection at the border (Denmark and  
 424 Lithuania). Other actions included recall from consumers (Denmark), withdrawal from the  
 425 market (Netherlands, Germany), product recall or withdrawal (Sweden), withdrawal from  
 426 recipients or destruction (Germany), public warning - press release (Norway), and informing  
 427 recipients (Spain). The two instances when ~~h~~Hepatitis A was reported in food items were in  
 428 2012 when frozen green mussels (*Perna canaliculus*) from New Zealand and frozen  
 429 strawberry cubes from China tested positive to the virus. The action taken in each instance  
 430 was different. In the case of contaminated frozen green mussels from New Zealand, the  
 431 notifying country (Italy) withdrew the product from the market while in the case of contaminated  
 432 frozen strawberry cubes from China, the notifying country (Belgium) only informed recipient.  
 433 *Campylobacter* notifications in foods originating from the Asia Pacific region were made only  
 434 in six instances (between 2009 and 2012) for asparagus, corn, ground peppers and spring  
 435 onions from Thailand and Vietnam. Nearly half (46%, i.e., 42/92) of *Bacillus* notifications in  
 436 food items from the Asia Pacific were from China.

#### 437 438 3.4. Pathogens associated with specific product categories

439 **Fig. 7** presents RASFF notifications of pathogenic microorganisms in food originating from  
 440 Asia-Pacific Region (n=1873) from 2020 - 2020 based on product category. The most  
 441 important product categories with *Salmonella* contamination were fruits and vegetables, herbs  
 442 and spices, nuts, nut products and seeds, while the most important product category with  
 443 *Vibrio* contamination was crustaceans and related products (**Fig. 7**).  
 444



**Fig. 7** RASFF notifications of pathogenic microorganisms in food originating from the Asia-Pacific Region (n=1873) from 2020 - 2020 based on food type.

A further analysis of pathogen diversity associated with notifications (2000 - 2020) revealed that crustaceans and related products from India, Malaysia and Vietnam showed the highest number of pathogen species (n >60) (**Supplementary Table 2**). Most of the pathogen species in crustaceans and related products in these countries were associated with shrimps (Table 6). Sri Lanka and Pakistan presented with the least (n=7) pathogen species in crustaceans and related products ~~imported~~~~exported~~ from the Asia-Pacific Region (**Supplementary Table 2**). Among the Asia-Pacific countries, the highest counts of pathogen species in fruits and vegetables were observed among ~~import~~~~export~~ from India and Bangladesh (**Supplementary Table 3**). More than 95% (i.e., India=184/193, Bangladesh = 149/152) of pathogen species notifications for fruits and vegetables ~~imported~~~~exported~~ from India and Bangladesh were associated with betel leaves (**Supplementary Table 3**). All the 346 pathogen notifications in fruits and vegetables ~~imported~~~~exported~~ from India and Bangladesh were *Salmonella* serovars (**Supplementary Table 3**). Pathogen notifications for mushrooms ~~imported~~~~exported~~ from the Asia-Pacific countries were all due to *Salmonella*, with the notable exception of mushrooms ~~imported~~~~exported~~ from China, which had additional pathogen notifications due to *Bacillus cereus*, *Clostridium perfringens*, *Clostridium* spp, aerobic mesophiles and moulds. Among the Asia-Pacific countries, spinach imported from Thailand had the highest number of reported *Salmonella* serovars (*Salmonella* Ndolo, *Salmonella* Aberdeen, *Salmonella* Bovismorbificans, *Salmonella enterica*, *Salmonella group B*, *Salmonella group D*, *Salmonella* Hvittingfoss, *Salmonella* Mbandaka, *Salmonella* Newport - see **Supplementary Table 3**).

Among the nuts, nut products and seeds, sesame seeds were the main food item contaminated by *Salmonella* (261 notifications), and these accounted for 87% of total *Salmonella* notifications (261/299, **Supplementary Table 4**). *Salmonella*-contaminated sesame seeds originated from India (n=260; 99.6%) and Thailand (n=1; 0.4%) (**Supplementary Table 5**). Fifty serovars of *Salmonella enterica* were isolated from sesame seeds (**Supplementary Table 4**). Among the herbs and spices, basil, mint and coriander from Thailand had the highest number of pathogen notifications. For instance, basil was associated with at least 8 *Salmonella enterica* serovars and 65 notifications, coriander with at least 12 *Salmonella enterica* serovars and 24 notifications, and mint with at least 2 *Salmonella enterica* serovars and 25 notifications (**Supplementary Table 6**).

#### 4. Discussion

The presence of pathogenic microorganisms in food products, particularly in minimally processed and ready-to-eat foods, constitutes a significant public health risk to consumers. In this study, we analysed notifications associated with microbiological hazards (MH) in foods ~~imported~~~~exported~~ from the Asia Pacific region into the EU between 2000 and 2020. Studies of this nature can potentially encourage countries to implement policies that improve quality of exported food products.

Unlike a previous report (Somorin, et al., 2021) which noted a growing trend in notifications from food products ~~imported~~~~exported~~ from Africa, there was no overall trend in the number of notifications from the Asia Pacific region during the same study period. This difference could be attributed to the significant decrease in MH notifications from the South and Southeast Asia subregions between 2015 and 2020. Interestingly, the countries responsible for the highest number of notifications during the study period are located in the South and Southeast Asia subregions.

Of the 1873 notifications, most (n=1456/1873) of them concerned food originating from India, Thailand, Vietnam and Bangladesh. A contributory factor could be the volume of trade between these countries and the EU. In 2018/19, India, Thailand and Vietnam ranked 10th, 14th and 17th, respectively, in terms of for the export value of food products traded with the EU (European Commission, 2020a). However, this appears not to be the sole reason. Compared to other countries in the Asia-Pacific region, EU food imports from China and Indonesia are the largest, with a combined value of about 10 billion euros (European Commission, 2020a), but only contributed 11% of notifications observed during the study period. New Zealand is the 19th highest exporter of food to the EU but contributed the least number of notifications from the region.

A more likely explanation for our observations is the nature of the microbial hazard and implicated foods. The highest number of MH notifications from the Asia-Pacific region was recorded in 2015. Aflatoxin contamination of nuts, nut products and seeds imported from China and *Salmonella* contamination in fruits and vegetables and nuts, nut products and seeds imported from India produced the three highest MH notifications from all regions for 2015 (European Commission 2016). In addition, *Salmonella* contamination of herbs and spices, fruits and vegetables, nuts, nut products, and seeds were responsible for the highest number of notifications overall. These ~~products~~ were predominantly products ~~importeds~~ from India, Thailand, Vietnam and Bangladesh. Food safety remains a significant challenge in the Asia-Pacific region and a hurdle to expanding agricultural and food product exports (Pham & Dinh, 2020). Food safety challenges include inadequate sanitation infrastructure, poor food safety knowledge and practices, and insufficient implementation of national food safety management systems (Food and Agricultural Organisation, 2018; Jaffee, Henson, Unnevehr, Grace, & Cassou, 2018; Mangla, et al., 2021; Ortega, 2017).

The highest number of notifications was made by the UK (28%). This is understandable because the standard for food quality, in terms of compliance requirement, is generally high in the UK. With about half of UK's food produced locally (DEFRA, 2020), a recent survey showed that a majority (97.5%) of food manufacturers within the UK had an integrated food safety management system in place (Mensah & Julien, 2011). Only about 5% of UK's food are imported from Asia and Australasia (DEFRA, 2020), yet exporters wishing to supply the UK are expected to have monitoring systems that ensure compliance with retail (product quality) and legislative (due diligence) requirements in the UK (Dolan & Humphrey, 2000). Notifications made by the UK (28%) mainly concerned foods originating from India and Bangladesh. Food safety has been a major challenge in South Asia and Southeast Asia, with huge impacts on public health, economics, and international trade (World Health Organization, 2020a). Southeast Asia has the second highest annual burden of foodborne diseases among the six

World Health Organization (WHO) regions, with 150 million foodborne illnesses and 175,000 deaths in 2010 (World Health Organization, 2015). In contrast to South Asia and Southeast Asia, there were significantly fewer notifications of pathogenic microorganisms in foods imported from Australia and New Zealand (n=23/1873; 1.2%), and this could be due to the strong food control systems in Australia and New Zealand (Ghosh, 2014).

Data received through the RASFF are a vital source of information on emerging health risks associated with specific food products. Import border rejections were the most common notification type in this study, while the most common actions taken were product redispach, destruction and non-authorisation of imports. Besides constituting a significant revenue loss for exporting countries, and causing reduced consumer confidence, these actions may lead to more stringent responses from the EU to safeguard the health of consumers (Chaoniruthisai, Punnakitkashem, & Rajchamaha, 2018; Somorin, et al., 2021). For example, peppers (*Capsicum* spp. except sweet) from India and Pakistan were subjected to increased official controls in 2018 due to concerns about the levels of pesticide residues (European Commission, 2020b). Data from these official controls and RASFF notifications provided evidence of continuous non-compliance, and these food products were assessed as constituting a serious risk to human health. In 2020, the EU legislated that these products should now include an official certificate confirming that maximum pesticide levels have not been exceeded (European Commission, 2020b).

Increased official controls are the first stage of a continuum that can end in the suspension of imports of a specific product. A good example is the suspension of imports into the EU of foodstuff containing betel leaves from Bangladesh due to the presence of *Salmonella*. This suspension has been in place since 2014 as no action plan for managing this hazard has been provided (European Commission, 2014, 2020b). A national monitoring programme for betel leaves imported into the United Kingdom between 2011 and 2017 showed that *Salmonella* contamination remains a significant hazard in this product (McLauchlin, et al., 2019). Betel is an important cash crop in Bangladesh and the principal source of livelihood in many rural communities (Nath & Inoue, 2009; Ullah, Tani, Tsuchiya, Rahman, & Rahman, 2020). Import/Export bans can have devastating consequences on income generation and food security. The poor food safety record in South Asia and Southeast Asia is due to several factors including climatic factors, poor sanitary conditions, and food habits. For example, extreme weather conditions and natural disasters, such as flooding, have been frequently occurring in countries Southeast Asia (A. Chen, Giese, & Chen, 2020), and since flood waters are often contaminated with sewage, this leads to increased contamination risk in the food supply chain and exposure to faecal pathogens (Yu, et al., 2018). Consumption of street-foods is common in many Asian countries and it is the only affordable form of nutrition for most people in low-income groups. However, inadequate observation of food hygiene practices significantly increases the exposure of consumers to foodborne pathogens (Reddy, Ricart, & Cadman, 2020).

There was a significant ( $p<0.006$ ) downward trend in the numbers of pathogenic microorganism notifications from foods imported/exported from Oceania to the EU from 2000-2020, unlike South Asia and East Asia. In more recent years (2015-2020), East Asia was the only sub-region with a significantly increasing trend in RASFF notifications due to contamination by pathogenic microorganisms. When considered by food categories, RASFF notifications regarding "nut, nut products and seeds" significantly increased over the years from 2000 - 2020. However, notifications of pathogenic microorganisms in "fruits and vegetables" significantly reduced in recent years (2015 – 2020). To address the food safety concerns in the affected regions, it is important that appropriate and effective methods to minimise contamination of foods by pathogenic microorganisms are implemented across the value chains in the originating countries. For instance, a framework has been recently established in the WHO Southeast Asia region (including India, Thailand and Bangladesh) to strengthen food control systems across the value chain and protect consumer health, building

on previous regional multi-actor interventions (World Health Organization, 2020b). These actions could have contributed to the downward trend in notifications of pathogenic microorganisms from South Asia and Southeast Asia in recent years (2015-2020) (Table 3). It is however important to take these RASFF database trend results cautiously As argued in published literature (Kowalska & Manning, 2020) iterative changes in food law potentially impacts on the frequency of regulatory sampling associated with border and inland regulatory checks in a way that could affect the generalisability of the trends noted.

Border rejection was the most frequent type of notification received (42.3%) in this study. When food ~~import~~~~exports~~ do not meet the standards set by the EU, a variety of actions can be taken, which have huge economic implications for the originating countries. About half of the notifications from the Asia-Pacific region resulted in importation not being authorized, leading to destruction or re-dispatch of the contaminated food product. Frequent non-compliance of some food products with EU food safety requirements has led to increased official control at border control posts and the imposition of suspension of entry into the EU (European Commission, 2020b). Increased identity and physical checks at EU border control points is in place for betel leaves and sesame seeds from India, and sweet peppers from China due to *Salmonella* contamination (European Commission, 2020b). Similar official controls exist for black pepper from Brazil as well as sesame seeds from Ethiopia, Nigeria, Sudan and Uganda due to *Salmonella* contamination (European Commission, 2020a, 2020b; Somorin, et al., 2021). Continuous non-compliance has led to the suspension of importation of betel leaves from Bangladesh to the EU since 2014 due to *Salmonella* contamination (European Commission, 2015). A similar ban is in place for dried beans from Nigeria due to pesticide residues (European Commission, 2020a, 2020b).

The most frequently reported genera of pathogenic microorganisms in foods from the Asia-Pacific region were *Salmonella* (73.6%), *Vibrio* (14.6%), *Bacillus* (4.3%) and *Escherichia* (2.2%). *Salmonella* spp. are the most reported pathogens associated with foods imported into the EU from the Asia-Pacific region. This corroborates the report by Somorin et al., (2021) on products imported into the EU from Africa. The predominance of *Salmonella*, an enteric bacterium, could be attributed to unhygienic practices during food production processes and this is corroborated by a previous study that reported a significant association between the presence of *Salmonella* spp. and *E. coli* in imported edible leaves retailed in the England (McLauchlin, et al., 2018). Foods reported in *Salmonella* notifications were mainly "fruits and vegetables", "herbs and spices" and "nuts, nut products and seeds". India accounted for over 35% of the *Salmonella* notifications in this study and were mostly from "nuts, nut products and seeds", "fruits and vegetables" and seafoods. In our study, about 178 serovars of *Salmonella enterica* were reported in foods from Asia-Pacific from 2000-2020 with serovars Weltevrede, Enteritidis and Hvitittingfoss being the most frequently reported. These serovars are significant causes of diarrhoea, and sometimes, invasive infections in tropical low-income Asian countries (Bangtrakulnonth, et al., 2004; Kantama & Jayanetra, 1996; Makendi, et al., 2016) as well as in Europe (ECDC/EFSA, 2021; Emberland, et al., 2007).

Over 95% of notifications regarding "fruit/vegetables" were from India and Bangladesh. Several *Salmonella* serovars have been shown to contaminate ready-to-eat betel leaves in Asia and those imported into the UK. Fakruddin et al. (2017) reported that 77% of betel leaves collected in local markets in Bangladesh were contaminated with *Salmonella* spp., with some of them imported from India. *Salmonella* from betel leaves also harbour multidrug resistance genes (Singh, et al., 2006), which could be disseminated to other countries through international trade. The majority of the notifications regarding *Salmonella* in betel leaves were from the UK, affecting products originating from India and Bangladesh. With South Asians being the largest ethnic minority group in the UK, and accounting for about 5% of the UK population (Office of National Statistics, 2011), there is a ready demand for betel leaves and other indigenous foods. Analysis of betel leaf consignments at Border Inspection Posts in England between 2011 and 2017 showed that 57% had *Salmonella* contamination

(McLauchlin, et al., 2019). In a separate study by McLauchlin *et al.* (2018), 14% of betel leaves collected from retailers in England had unsatisfactory levels of *Salmonella* including *Salmonella* Bareilly, *Salmonella* Hvittingfoss, *Salmonella* Lichfield, and *Salmonella* Newport. The frequency and high contamination rates of *Salmonella* in betel leaves pose a public health concern for importing countries. While the UK is no longer part of the EU, it is expected to remain an important market for produce from South Asia and Southeast Asia and retain stringent food safety regulations.

*Salmonella* is a principal aetiological agent of foodborne disease and is the third leading cause of death due to diarrhoeal disease globally (Ferrari, et al., 2019; World Health Organization, 2019, 2020c). *Salmonella* is the most common cause of foodborne outbreaks in the EU and in 2018, was implicated in almost 30% of outbreaks in the region (EFSA/ECDC, 2019). Fresh produce, including fruits, vegetables, herbs, and spices ~~imported-exported~~ from the Asia-Pacific region, were the food products most associated with *Salmonella*. Fruit and vegetables are increasingly recognised as vehicles for transmitting pathogenic organisms and have been implicated in global foodborne outbreaks (Bisht, et al., 2021; Carstens, Salazar, & Darkoh, 2019; Wadamori, Gooneratne, & Hussain, 2017).

Contamination of fresh produce can occur at different stages of the supply chain. *Salmonella* is widespread in the environment and is commonly found in materials susceptible to faecal contamination (Machado-Moreira, Richards, Brennan, Abram, & Burgess, 2019). Contaminated soil and irrigation water are important sources of preharvest contamination. Unhygienic practices during harvest, processing and transport are the main contributors to post-harvest contamination (Dos Santos, et al., 2020; Park, et al., 2012). Fresh produce is usually consumed with minimal processing, making it challenging to eradicate pathogenic microorganisms via conventional methods like cooking. Several studies have noted unacceptable levels of *Salmonella* contamination in fresh produce available for retail sale in the Asia-Pacific region (Abatcha, Effarizah, & Rusul, 2018; Nguyen, et al., 2021; Vital, Dimasuay, Widmer, & Rivera, 2014). These reports are of concern and highlight the need for improved food safety management practices along the fresh produce supply chain. To our knowledge, no outbreaks linked to *Salmonella* in fresh produce from the Asia-Pacific region have been reported in the EU. However, it is not always possible to identify the potential source and vehicle of infection during an outbreak (EFSA/ECDC, 2018).

The most frequently reported *Vibrio* spp. in the notifications were *V. cholerae* (46.6%) and *V. parahaemolyticus* (46%) while *V. vulnificus*, *V. alginolyticus*, *V. fluvialis* and *V. mimicus* were less frequently isolated. The main food category contaminated by *Vibrio* was “crustaceans and related products” particularly shrimps and prawns. Precious studies have reported that most of the foods with *Vibrio* contamination originated from India, Malaysia and Vietnam. For instance, *V. parahaemolyticus* was reported in 76% of shrimps ~~imported-exported~~ from Thailand (Wong, Chen, Liu, & Liu, 1999) and 43.4% of shrimps in China (Xu, Cheng, Wu, Zhang, & Xie, 2016). A *V. parahaemolyticus* pandemic serotype O3:K6 originating from India have been detected in many Asian countries (Bag, et al., 1999) as well as in Europe (Martinez-Urtaza, et al., 2005). predominantly linked to crustaceans such as shrimps and prawns and related products. *Vibrio* spp. are important causes of diarrhoea and cholera associated with the consumption of raw or undercooked seafood and are frequently detected in seafood products from countries in the Asia-Pacific region (X. Chen, et al., 2018; Fu, et al., 2020; Kumar & Lalitha, 2013; Lee, et al., 2019; Nakaguchi, 2013). Seafood imported from Asia was associated with an outbreak in France in 1997 (Lemoine, Germanetto, & Giraud, 1999). Similarly, Briet et al. (2018) reported the presence of a carbapenemase-producing strain of *V. parahaemolyticus* isolated from shrimps imported into France from Vietnam. This study provides evidence that imported foods may be potential vehicles for the transmission of clinically relevant microorganisms.



The predominant *Bacillus* spp. reported in the notifications was *Bacillus cereus* and it was reported in cheese, soy cheese, tofu, garlic powder, ground pepper, pumpkin seeds, mushroom, sesame seeds, desiccated coconut, cinnamon powder, curry powder, ginger powder and ground cumin across East-, South- and Southeast Asia. Production and processing of these food products involves drying and/or grinding, where they could come in contact with soil harbouring *Bacillus* spores (Vilain, Luo, Hildreth, & Brözel, 2006). *B. cereus* has been previously reported in soybean products/tofu (Ananchaipattana, et al., 2012; Keisam, Tuikhar, Ahmed, & Jeyaram, 2019), spice and herbs (Banerjee & Sarkar, 2003) and cheese (Kumari & Sarkar, 2014).

Furthermore, foodborne viruses, such as Hepatitis A and Norovirus, were reported in green mussels and berries (strawberry and raspberry) mainly originating from Vietnam and China. Norovirus has been reported in whole fruits and vegetable leaves in France (EFSA/ECDC, 2021) and the UK (Cook, Williams, & D'Agostino, 2019) and the majority of foodborne viral outbreaks are associated with frozen fruits (Nasheri, Vester, & Petronella, 2019). A large outbreak in Germany in 2012 was associated with strawberries imported from China (Bernard, et al., 2014). Norovirus was associated with 22.5% of all outbreak-related illnesses in the EU in 2019, with contaminated shellfish and fish products playing a significant role (EFSA/ECDC, 2021). Although there were few notifications regarding *Campylobacter* and *Yersinia*, and no report of *Listeria monocytogenes* in foods from the Asia-Pacific region, foodborne diseases caused by these pathogens are among the most common in Europe (EFSA/ECDC, 2021), hence requiring continuous surveillance of imported foods from other parts of the world. The lack of consistency in the actions taken by different EU member states in response to notifications arising from the presence of norovirus and Hepatitis A is of interest. This suggests a lack in uniformity across Member States with respect to pathogens detected in food items. To safeguard public health, it may be necessary for stakeholders to collate a guidance action document with respect to specific actions that should be taken if a particular pathogen is detected in imported food products. This will ensure consistency across notifying countries. One limitation with the current study is the lack of comparison between the numbers of pathogen notifications and the total number of foods or food product category from each country that is tested for pathogens. This is partly because this data is not recorded in the RASSF database. It is hoped that these considerations will be included in future improvements in the RASSF database.

## 5. Conclusion

The analysis of RASFF notifications for foods from the Asia-Pacific region from 2000 – 2020 showed that foods from India, Thailand, Vietnam and Bangladesh had the most notifications due to pathogenic microorganisms. *Salmonella* was the most predominant foodborne pathogen reported in foods ~~imported~~<sup>exported</sup> from Asia-Pacific to the EU, with India accounting for over 35% of *Salmonella* notifications (fresh produce) and *Vibrio* spp. contamination of seafood products such as shrimps and prawns from India, Malaysia, and Vietnam. There were very few notifications regarding *Campylobacter* spp. and *Yersinia* spp., and no notification of *L. monocytogenes* was reported. Increasingly, food producers are connected to consumers via food supply chains that cross international borders. These value chains offer a stable supply of food that meet consumers varied needs and preferences and provide a source of income to producers (OECD, 2020). An improved understanding of the food safety challenges along these international supply chains, and capacity building to support implementation of good manufacturing practices and food management systems, is required to safeguard consumer health and sustain the economic development of food-producing nations. Similarly, improved hygienic practices should be implemented across the value chains of imported food products from the region to ensure continuous access to international markets and sustained income.



752

753 **CRedit authorship contribution statement**

754 **Ayokunle C. Dada:** Conceptualization, Methodology, Formal analysis, Writing - original draft,  
 755 Review and Editing, Writing - review & editing. **Yinka M. Somorin:** Writing - original draft,  
 756 Review and Editing, Writing - review & editing. **Collins N. Ateba:** Writing - original draft,  
 757 Review and Editing, Writing - review & editing. **Helen Onyeaka:** Review and Editing, Writing  
 758 - review & editing **Amarachukwu Anyogu:** Review and Editing, Writing - review & editing.  
 759 **Nor Azman Kasan:** Writing - original draft, Review and Editing, Writing - review & editing.  
 760 **Olumide A. Odeyemi:** Conceptualization, Methodology, Formal analysis, Writing - original  
 761 draft, Review and Editing, Writing - review & editing.

762

763

764

765

766

767

768

769

770

771 **References**

772

773 [Abatcha, M.G., Effarizah, M.E., Rusul, G. \(2018\). Prevalence, antimicrobial resistance,](#)  
 774 [resistance genes and class 1 integrons of \*Salmonella\* serovars in leafy vegetables,](#)  
 775 [chicken carcasses and related processing environments in Malaysian fresh food](#)  
 776 [markets. \*Food Control\*, 91, 170–180. <https://doi.org/10.1016/j.foodcont.2018.02.030>.](#)  
 777 [Ananchaipattana, C., Hosotani, Y., Kawasaki, S., Pongswat, S., Latiful, B., Isobe, S. and Inatsu,](#)  
 778 [Y. 2012; Bacterial contamination of soybean curd \(tofu\) sold in Thailand. \*Food Sci.\*  
 779 \[Technol. Res.\]\(#\) 18\(6\), 843–848](#)  
 780 [Bag, P. K., Nandi, S., Bhadra, R. K., Ramamurthy, T., Bhattacharya, S. K., Nishibuchi, M.,](#)  
 781 [Hamabata, T., Yamasaki, S., Takeda, Y. and Nair, G. B. 1999. Clonal diversity](#)  
 782 [among recently emerged strains of \*Vibrio parahaemolyticus\* O3:K6 associated with](#)  
 783 [pandemic spread. \*J Clin Microbiol\* 37\(7\): 2354–2357.](#)  
 784 [Balali, G.I., Yar, D.D., Dela, G.B.A., Adjoi Kusi, P. \(2020\). Microbial Contamination, an](#)  
 785 [Increasing Threat to the Consumption of Fresh Fruits and Vegetables in Today's](#)  
 786 [World. \*International Journal of Microbiology\*, 3029295,](#)  
 787 [https://doi.org/10.1155/2020/3029295](#)  
 788 [Banerjee, M. and Sarkar, P. K. 2003. Microbiological quality of some retail spices in India.](#)  
 789 [Food Research International 36: 469–474.](#)  
 790 [Bangtrakulnonth, A., Pornreongwong, S., Pulsrikarn, C., Sawanpanyalert, P., Hendriksen, R.](#)  
 791 [S., Wong, D. M. and Aarestrup, F. M. 2004. \*Salmonella\* Serovars from Humans and](#)  
 792 [Other Sources in Thailand, 1993–2002. \*Emerging Infectious Diseases\*, 10\(1\), 131–](#)  
 793 [136.](#)  
 794 [Bernard, H., Faber, M., Wilking, H., Haller, S., Höhle, M., Schielke, A., Ducomble, T., Sifczyk,](#)  
 795 [C., Merbecks, S. S., Fricke, C., Hamouda, O., Stark, K., Werber, D. a on behalf of](#)  
 796 [the Outbreak Investigation Team. 2014. Large multistate outbreak of norovirus](#)  
 797 [gastroenteritis associated with frozen strawberries, Germany, 2012. \*Euro Surveill.\*](#)  
 798 [19\(8\): 20719.](#)  
 799 [Bisht, A., Kamble, M.P., Choudhary, P., Chaturvedi, K., Kohli, G., Juneja, V.K., Sehgal, S.,](#)  
 800 [Taneja, N.K. \(2020\). A surveillance of foodborne disease outbreaks in India: 2009–](#)  
 801 [2018, \*Food Control\*, 121, 107630, <https://doi.org/10.1016/j.foodcont.2020.107630>.](#)

Borchers, A., Teuber, S. S., Keen, C. L., & Gershwin, M. E. (2010). Food safety. Clinical reviews in allergy & immunology, 39(2), 95-141.

Briet A., Helsen N., Delannoy S., Debuiche S., Brisabois A., Midelet G., Granier SA. (2018). NDM-1-producing *Vibrio parahaemolyticus* isolated from imported seafood. J Antimicrob Chemother. 73, 2578-2579. <https://doi.org/10.1093/jac/dky200>.

Carstens, C. K., Salazar, J. K., & Darkoh, C. (2019). Multistate Outbreaks of Foodborne Illness in the United States Associated With Fresh Produce From 2010 to 2017. Frontiers in Microbiology, 10, 2667. <https://doi.org/10.3389/fmicb.2019.02667>

Chaoniruthisai, P., Punnakitikashem, P., Rajchamaha, K. (2018). Challenges and difficulties in the implementation of a food safety management system in Thailand: A survey of BRC-certified food productions, Food Control, 93, 274-282, <https://doi.org/10.1016/j.foodcont.2018.06.004>.

Chen, A., Giese, M. and Chen, D. 2020. Flood impact on Mainland Southeast Asia between 1985 and 2018—The role of tropical cyclones. J Flood Risk Management 13: e12598.

Chen, X., Zhu, Q., Yu, F., Zhang, W., Wang, R., Ye, X., Jin, L., Liu, Y., Li, S., & Chen, Y. (2018). Serology, virulence and molecular characteristics of *Vibrio parahaemolyticus* isolated from seafood in Zhejiang province. PloS one, 13(10), e0204892. <https://doi.org/10.1371/journal.pone.0204892>

Commission, E. (2020a). The Rapid Alert System for Food and Feed Annual Report 2019. Retrieved from <https://op.europa.eu/en/publication-detail/-/publication/2c5c7729-0c31-11eb-bc07-01aa75ed71a1/language-en/format-PDF/source-174742448>

Commission, E. (2020b). RASFF Food and Feed Safety Alerts. Retrieved from [https://ec.europa.eu/food/safety/rasff\\_en](https://ec.europa.eu/food/safety/rasff_en)

Cook, N., Williams, L. and D'Agostino, M. 2019. Prevalence of norovirus in produce sold at retail in the United Kingdom. Food Microbiology 79: 85-89.

D'Ortenzio, E., Weill, F. X., Ragonneau, S., Lebon, J. A., Renault, P. and Pierre, V. 2008. First report of a *Salmonella enterica* serovar Weltevreden outbreak on Réunion Island, France, August 2007. Euro Surveill. 13(32): 18949.

ECDC and EFSA (European Centre for Disease Prevention and Control/European Food Safety Authority) (2018). Multi-country outbreak of *Salmonella* Agona possibly linked to ready-to-eat food—26 July 2018. Stockholm and Parma.

EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control), (2019). The European Union One Health 2018 Zoonoses Report. EFSA Journal, 17:5926. <https://doi.org/10.2903/j.efsa.2019.5926>

EFSA. (2019). The European Union one health 2018 zoonoses report. European Centre for Disease Prevention, Control. European Food Safety Authority Journal, 17(12), e05926.

Emberland, K. E., Ethelberg, S., Kuusi, M., Vold, L., Jensvoll, L., Lindstedt, B. A., Nygård, K., Kjelsø, C., Torpdahl, M., Sørensen, G., Jensen, T., Lukinmaa, S., Niskanen, T. and Kapperud, G. 2007. Outbreak of *Salmonella* Weltevreden infections in Norway, Denmark and Finland associated with alfalfa sprouts, July-October 2007. Euro Surveill. 12(48): 3321.

ESCAP. (2020). Economic and Social Survey of Asia and the Pacific 2020: Towards sustainable economies. Retrieved from <https://www.unescap.org/sites/default/files/publications/Economic%20and%20Social%20Survey%20of%20Asia%20and%20the%20Pacific%202020%20Towards%20Sustainable%20economies.pdf>

European Centre for Disease Prevention and Control (ECDC) and European Food Safety Authority (EFSA). 2021. Multi-country outbreak of *Salmonella* Enteritidis sequence type (ST)11 infections linked to poultry products in the EU/EEA and the United Kingdom—25 February 2021. Stockholm: ECDC/EFSA; 2021.

European Commission Health & Food Safety (2016). RASFF The Rapid Alert System for Food and Feed 2015 annual report. Available from:

<https://op.europa.eu/en/publication-detail/-/publication/98e04cfc-aa54-11e6-aab7-01aa75ed71a1/language-en/format-PDF/source-174743070> Accessed 19.02.2021

European Commission. (2020a). Monitoring EU Agri Food Trade: Developments in 2019. Available from: <https://ec.europa.eu/info/news/2019-record-year-eu-agri-food-trade-2020-mar-27-en#moreinfo> Accessed 12.02.2021.

European Commission, 2020. Commission implementing regulation (EU) 2020/625 of 7 May 2020. *Official Journal of the European Union* 144: 13-33

European Commission. (2014). Commission implementing decision of 13 February 2014 suspending temporarily imports from Bangladesh of foodstuffs containing or consisting of betel leaves ('Piper betle'). Available from: <https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32014D0088&from=DA> Accessed 25.02.2021.

European Commission. (2020b). Commission implementing regulation (EU) 2020/625 of 6 May 2020. *Official Journal of the European Union*, 63, 13.

European Commission. 2015. Commission implementing regulation (EU) 2015/943 of 18 June 2015. *Official Journal of the European Union* 154: 8-9

European Food Safety Authority (EFSA) and European Centre for Disease Prevention and Control (ECDC). 2021. The European Union One Health 2019 Zoonoses Report. *EFSA Journal* 19(2): 6406.

Fakruddin, M., Sultana, R., Hossain, M. N., Rahaman, M. M., Islam, M. K. and Ahmed, M. M. 2017. Occurrence of ingression of *Salmonella* spp. in Betel leaf (*Piper betle* L.). *International Journal of Food Contamination* 4: 6.

FAO. (2018). Shiga toxin-producing *Escherichia coli* (STEC) and food: attribution, characterization, and monitoring. Retrieved from <https://apps.who.int/iris/bitstream/handle/10665/272871/0789241514279-eng.pdf?ua=1>

FAO. (2019). Foodborne illnesses in Asia-Pacific 'needlessly' sicken 275 million annually and threaten trade. Retrieved from <http://www.fao.org/asiapacific/news/detail-events/en/c/1197008/#:~:text=In%20the%20Asia%20Pacific%20region,of%20Zero%20Hunger%20by%202030>.

Ferrari, R.G., Rosario, D.K.A., Cunha Neto, A., Mano, S.B., Figueiredo, E.E.S., Conte Junior, C.A. (2019). Worldwide Epidemiology of *Salmonella* Serovars in Animal-Based Foods: a Meta-analysis. *Applied and Environmental Microbiology*, 85, e00591-19; <https://doi.org/10.1128/AEM.00591-19>

Food and Agricultural Organisation of the United Nations (2018). Addressing the food safety challenges of the Asia-Pacific region. 34<sup>th</sup> Session of the FAO Regional Conference for Asia and the Pacific. Available from: <http://www.fao.org/3/MV819en/mv819en.pdf> Accessed 24.02.2021

Fu, H., Yu, P., Liang, W., Kan, B., Peng, X., & Chen, L. (2020). Virulence, Resistance, and Genomic Fingerprint Traits of *Vibrio cholerae* Isolated from 12 Species of Aquatic Products in Shanghai, China. *Microbial drug resistance* (Larchmont, N.Y.), 26(12), 1526-1539. <https://doi.org/10.1089/mdr.2020.0269>

Ghosh, D. 2014. Food safety regulations in Australia and New Zealand Food Standards. *J Sci Food Agric* 94: 1970-1973

Jaffee, S., Henson, S., Unnevehr, L., Grace, D., Cassou, Emilie. (2018). The Safe Food Imperative: Accelerating Progress in Low- and Middle-Income Countries. *Agriculture and Food Series*; Washington, DC: World Bank. © World Bank. License: CC-BY 3.0 IGO. Available from: <https://openknowledge.worldbank.org/handle/10986/30568> Accessed 10.02.2021.

*Journal of Cleaner Production*, 284, 124709, <https://doi.org/10.1016/j.jclepro.2020.124709>.

Kaakoush, N. O., Castañero-Rodríguez, N., Mitchell, H. M., & Man, S. M. (2015). Global epidemiology of *Campylobacter* infection. *Clinical microbiology reviews*, 28(3), 687-720.

Kantama, L. and Jayanetra, P. 1996. *Salmonella enteritidis* outbreak in Thailand: study by random amplified polymorphic DNA (RAPD) analysis. *Southeast Asian J Trop Med Public Health*. 27(1): 119-125.

Keisam, S., Tuikhar, N., Ahmed, G., and Jeyaram, K. 2019. Toxigenic and pathogenic potential of enteric bacterial pathogens prevalent in the traditional fermented foods marketed in the Northeast region of India. *International Journal of Food Microbiology* 296: 21-30.

Kim, K., Kim, S., & Park, C. Y. (2020). Food Security in Asia and the Pacific amid the COVID-19 Pandemic.

Kowalska, A., & Manning, L. (2020). Using the rapid alert system for food and feed: potential benefits and problems on data interpretation. *Critical Reviews in Food Science and Nutrition*, 1-14.

Kumar R, Lalitha KV. Prevalence and molecular characterization of *Vibrio cholerae* O1, non-O1 and non-O139 in tropical seafood in Cochin, India. (2013). *Foodborne Pathog Dis*. 10, 278-83. <https://doi.org/10.1089/fpd.2012.1310>.

Kumari, S. and Sarkar, P. K. 2014. Prevalence and characterization of *Bacillus cereus* group from various marketed dairy products in India. *Dairy Sci. & Technol*. 94: 483-497.

Lee Y, Choi Y, Lee S, Lee H, Kim S, Ha J, Lee J, Oh H, Kim Y, Yoon Y. Occurrence of pathogenic *Vibrio parahaemolyticus* in seafood distribution channels and their antibiotic resistance profiles in S. Korea. *Lett Appl Microbiol*. 2019 Feb;68(2):128-133. <https://doi.org/10.1111/lam.13099>.

Lemoine, T., Germanetto, P. and Giraud, P. 1999. Toxi-infection alimentaire à *V. parahaemolyticus*. *Bull Epidemiol Hebdo*. 10: 37-38.

Machado-Moreira B, Richards K, Brennan F, Abram F, Burgess CM. (2019). Microbial Contamination of Fresh Produce: What, Where, and How? *Comprehensive Reviews in Food Science and Food Safety*, 6, 1727-1750. <https://doi.org/10.1111/1541-4337.12487>.

Makendi C, Page AJ, Wren BW, Le Thi Phuong T, Clare S, et al. 2016. A phylogenetic and phenotypic analysis of *Salmonella enterica* Serovar Weltevreden, an emerging agent of diarrheal disease in tropical regions. *PLOS Neglected Tropical Diseases* 10(2): e0004446.

Mangla, S.K., Bhattacharya, A., Yadav, A.K., Sharma, Y.K., Ishizaka, A., Luthra, S., Chakraborty, R. (2021). A framework to assess the challenges to food safety initiatives in an emerging economy.

Martinez-Urtaza, J., Simental, L., Velasco, D., DePaola, A., Ishibashi, M., Nakaguchi, Y., Nishibuchi, M., Carrera-Flores, D., Rey-Alvarez, C. and Pousa, A. 2005. Pandemic *Vibrio parahaemolyticus* O3:K6, Europe. *Emerg Infect Dis*. 11(8): 1319-1320.

McLauchlin, J., Aird, H., Andrews, N., Chattaway, M., de Pinna, E., Elviss, N., Jørgensen, F., Larkin, L., Willis, C. (2019). Public health risks associated with *Salmonella* contamination of imported edible betel leaves: Analysis of results from England, 2011-2017. *International Journal of Food Microbiology*, 298, 1-10. <https://doi.org/10.1016/j.ijfoodmicro.2019.03.004>.

McLauchlin, J., Aird, H., Andrews, N., Chattaway, M., de Pinna, E., Elviss, N., Jørgensen, F., Larkin, L. and Willis, C. 2019. Public health risks associated with *Salmonella* contamination of imported edible betel leaves: Analysis of results from England, 2011-2017. *International Journal of Food Microbiology* 298: 1-10.

McLauchlin, J., Aird, H., Charlett, A., Chattaway, M., Elviss, N., Hartman, H., Jenkins, C., Jørgensen, F., Larkin, L., Sadler-Reeves, L. and Willis, C. 2018. Imported edible leaves collected at retail sale in England during 2017 with an emphasis on betel and curry leaves: Microbiological quality with respect to *Salmonella*, Shiga toxin-producing *E. coli* (STEC) and levels of *Escherichia coli*. *J. Appl. Microbiol.*, 125: 1175-1185.

Nakaguchi, Y. (2013). Contamination by *Vibrio parahaemolyticus* and Its Virulent Strains in Seafood Marketed in Thailand, Vietnam, Malaysia, and Indonesia. *Tropical Medicine and Health*, 71, 95–102. <https://doi.org/10.2149/tmh.2011-06>

Nasheri, N., Vester, A. and Petronella, N. 2019. Foodborne viral outbreaks associated with frozen produce. *Epidemiology and Infection* 147: e291.

Nath, T.K., Inoue, M. (2009). Sustainability Attributes of a Small-Scale Betel Leaf Agroforestry System: A Case Study in North-eastern Hill Forests of Bangladesh. *Small-scale Forestry*, 8, 289–304. <https://doi.org/10.1007/s11842-009-9084-4>.

Nguyen, T.K., Bui, H.T., Truong, T.A., Lam, D.N., Ikeuchi, S., Ly, L.K.T., Hara-Kudo, Y., Taniguchi, T., Hayashidani, H. (2021). Retail fresh vegetables as a potential source of *Salmonella* infection in the Mekong Delta, Vietnam. *Int J Food Microbiol.* 341:109049. <https://doi.org/10.1016/j.ijfoodmicro.2021.109049>

North China. *BMC Microbiology* 16: 32

Office of National Statistics, 2011. UK Census 2011. <https://www.ons.gov.uk/census/2011census>

Organisation for Economic Cooperation and Development (2020). Global value chains connect producers to consumers across the world Available from: <https://www.oecd.org/agriculture/topics/global-value-chains-and-agriculture/>

Ortega, D.L. and Tschirley, D.L. (2017). Demand for food safety in emerging and developing countries: A research agenda for Asia and Sub-Saharan Africa. *Journal of Agribusiness in Developing and Emerging Economies*, 7, 21–34. <https://doi.org/10.1108/JADEE-12-2014-0045>.

Park, S., Szonyi, B., Gautam, J., Nightingale, K., Anciso, J., Ivanek, R. (2012). Risk Factors for Microbial Contamination in Fruits and Vegetables at the Preharvest Level: A Systematic Review. *Journal of Food Protection*, 75, 2055–2081. <https://doi.org/10.4315/0362-028X-JFP-12-160>

Pham, H.V. and Dinh, T.L. (2020). The Vietnam's food control system: Achievements and remaining issues. *Food Control*, 108, 106862. <https://doi.org/10.1016/j.foodcont.2019.106862>.

Reddy, A. A., Ricart, S. and Cadman, T. 2020. Driving factors of food safety standards in India: learning from street food vendors' behaviour and attitude. *Food Security* 12: 1201–1217

Rezaei, M., & Liu, B. (2017). Food loss and waste in the food supply chain. *International Nut and Dried Fruit Council: Reus, Spain*, 26–27.

Rivas, L., Strydom, H., Paine, S., Wang, J., & Wright, J. (2021). Yersiniosis in New Zealand. *Pathogens*, 10(2), 191.

Singh, B. R., Singh, M., Singh, P., Babu, N., Chandra, M., Agarwal, R. K. 2006. Prevalence of multidrug-resistant *Salmonella* on ready-to-eat betel leaves (Paan) and in water used for soaking betel leaves in North Indian cities. *J Food Prot.* 69(2): 288–292

Sodagari, H. R., Wang, P., Robertson, I., Habib, I., & Sahibzada, S. (2020). Non-Typhoidal *Salmonella* at the Human-Food-of-Animal-Origin Interface in Australia. *Animals*, 10(7), 1192.

Somarin, Y. M., Odeyemi, O. A., & Ateba, C. N. (2021). *Salmonella* is the most common foodborne pathogen in African food exports to the European Union: Analysis of Rapid Alert System for Food and Feed (1999–2019). *Food Control*, 107849. doi:<https://doi.org/10.1016/j.foodcont.2020.107849>

Somarin, Y. M., Odeyemi, O. A., & Ateba, C. N. (2021). *Salmonella* is the most common foodborne pathogen in African food exports to the European Union: Analysis of the Rapid Alert System for Food and Feed (1999–2019). *Food Control*, 123, 107849.

Somarin, Y.M., Odeyemi, O.A. and Ateba, C.N. 2021. *Salmonella* is the most common foodborne pathogen in African food exports to the European Union: Analysis of Rapid Alert System for Food and Feed (1999–2019). *Food Control* 123: 107849.

- Sugiri, Y. D., Geelz, G., Meeyam, T., Baumann, M. P., Kloor, J., Chaisowwong, W., & Alter, T. (2014). Prevalence and antimicrobial susceptibility of *Listeria monocytogenes* on chicken carcasses in Bandung, Indonesia. *Journal of food protection*, 77(8), 1407–1410.
- Tähkääpää, S., Majjala, R., Korkeala, H., & Nevas, M. (2015). Patterns of food frauds and adulterations reported in the EU rapid alert system for food and feed and in Finland. *Food Control*, 47, 175–184.
- Ullah, A.S.M., Tani, M., Tsuchiya, J., Rahman, A.M., Rahman, Z.M. (2020). Impact of Betel Leaf Cultivation on the Protected Forest Area of Teknaf Peninsula, Bangladesh. *Small-scale Forestry*, 19, 335–355. <https://doi.org/10.1007/s11842-020-09441-w>
- Vilain, S., Luo, Y., Hildreth, M.B. and Brözel, V.S. 2006. Analysis of the life cycle of the soil saprophyte *Bacillus cereus* in liquid soil extract and in soil. *Applied and Environmental Microbiology* 72(7): 4970–4977
- Vital, P.G., Dimasuay, G.B., Widmer, K.W., Rivera, W.L. (2014). Microbiological Quality of Fresh Produce from Open Air Markets and Supermarkets in the Philippines. *The Scientific World Journal*, Article ID 219534, <https://doi.org/10.1155/2014/219534>
- Wadamori, Y., Gooneratne, R., Hussain, M.A. (2016). Outbreaks and factors influencing microbiological contamination of fresh produce. *Journal of the Science of Food and Agriculture*, 97, 1396–1403. <https://doi.org/10.1002/jsfa.8125>
- Wang, S., Weller, D., Falardeau, J., Strawn, L. K., Mardones, F. O., Adell, A. D., & Switt, A. I. M. (2016). Food safety trends: from globalization of whole genome sequencing to application of new tools to prevent foodborne diseases. *Trends in Food Science & Technology*, 57, 188–198.
- WHO. (2020). Food Safety. Retrieved from <https://www.who.int/news-room/fact-sheets/detail/food-safety>
- Wong, H. C., Chen, M. C., Liu, S. H. and Liu, D. P. 1999. Incidence of highly genetically diversified *Vibrio parahaemolyticus* in seafood imported from Asian countries. *International Journal of Food Microbiology* 52: 181–189
- World Health Organisation (2020) *Salmonella* non-typhoidal. Available from: [https://www.who.int/news-room/fact-sheets/detail/salmonella-\(non-typhoidal\)](https://www.who.int/news-room/fact-sheets/detail/salmonella-(non-typhoidal)) Accessed 25.02.2024
- World Health Organization, 2015. WHO estimates of the global burden of foodborne diseases. Foodborne Disease Burden Epidemiology Reference Group 2007–2015. World Health Organization, Geneva.
- World Health Organization, 2020. Framework for action on food safety in the WHO South-East Asia Region. New Delhi: World Health Organization, Regional Office for South-East Asia.
- Worldometer. (2021). Current World Population. Retrieved from <https://www.worldometers.info/world-population/#ref-1>
- Xu, X., Cheng, J., Wu, Q., Zhang, J. and Xie, T. 2016. Prevalence, characterization, and antibiotic susceptibility of *Vibrio parahaemolyticus* isolated from retail aquatic products in
- Yu, P., Zaleski, A., Li, Q., He, Y., Mapili, K., Pruden, A., Alvarez, P. J. J. and Stadler, L. B. 2018. Elevated levels of pathogenic indicator bacteria and antibiotic resistance genes after Hurricane Harvey's flooding in Houston. *Environ. Sci. Technol. Lett.* 5(8): 481–486



1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113

## References

- Abatcha, M. G., Effarizah, M. E., & Rusul, G. (2018). Prevalence, antimicrobial resistance, resistance genes and class 1 integrons of *Salmonella* serovars in leafy vegetables, chicken carcasses and related processing environments in Malaysian fresh food markets. *Food Control*, 91, 170-180.
- Ananchaipattana, C., Hosotani, Y., Kawasaki, S., Pongswat, S., Latiful, B. M., Isobe, S., & Inatsu, Y. (2012). Bacterial contamination of soybean curd (tofu) sold in Thailand. *Food Science and Technology Research*, 18(6), 843-848.
- Bag, P. K., Nandi, S., Bhadra, R. K., Ramamurthy, T., Bhattacharya, S., Nishibuchi, M., Hamabata, T., Yamasaki, S., Takeda, Y., & Nair, G. B. (1999). Clonal diversity among recently emerged strains of *Vibrio parahaemolyticus* O3: K6 associated with pandemic spread. *Journal of clinical microbiology*, 37(7), 2354-2357.
- Banerjee, M., & Sarkar, P. K. (2003). Microbiological quality of some retail spices in India. *Food Research International*, 36(5), 469-474.
- Bangtrakulnonth, A., Pornreongwong, S., Pulsrikarn, C., Sawanpanyalert, P., Hendriksen, R. S., Wong, D. M. L. F., & Aarestrup, F. M. (2004). *Salmonella* serovars from humans and other sources in Thailand, 1993–2002. *Emerging infectious diseases*, 10(1), 131.
- Bernard, H., Faber, M., Wilking, H., Haller, S., Höhle, M., Schielke, A., Ducomble, T., Sifczyk, C., Merbecks, S., & Fricke, G. (2014). Large multistate outbreak of norovirus gastroenteritis associated with frozen strawberries, Germany, 2012. *Eurosurveillance*, 19(8), 20719.
- Bisht, A., Kamble, M. P., Choudhary, P., Chaturvedi, K., Kohli, G., Juneja, V. K., Sehgal, S., & Taneja, N. K. (2021). A surveillance of food borne disease outbreaks in India: 2009–2018. *Food Control*, 121, 107630.
- Borchers, A., Teuber, S. S., Keen, C. L., & Gershwin, M. E. (2010). Food safety. *Clinical reviews in allergy & immunology*, 39(2), 95-141.
- Briet, A., Helsens, N., Delannoy, S., Debuiche, S., Brisabois, A., Midelet, G., & Granier, S. (2018). NDM-1-producing *Vibrio parahaemolyticus* isolated from imported seafood. *J Antimicrob Chemother* doi, 10.
- Carstens, C. K., Salazar, J. K., & Darkoh, C. (2019). Multistate outbreaks of foodborne illness in the United States associated with fresh produce from 2010 to 2017. *Frontiers in microbiology*, 10, 2667.
- Chaoniruthisai, P., Punnakitakashem, P., & Rajchamaha, K. (2018). Challenges and difficulties in the implementation of a food safety management system in Thailand: A survey of BRC certified food productions. *Food Control*, 93, 274-282.
- Chen, A., Giese, M., & Chen, D. (2020). Flood impact on Mainland Southeast Asia between 1985 and 2018—The role of tropical cyclones. *Journal of Flood Risk Management*, 13(2), e12598.
- Chen, X., Zhu, Q., Yu, F., Zhang, W., Wang, R., Ye, X., Jin, L., Liu, Y., Li, S., & Chen, Y. (2018). Serology, virulence and molecular characteristics of *Vibrio parahaemolyticus* isolated from seafood in Zhejiang province. *PloS one*, 13(10), e0204892.
- Cook, N., Williams, L., & D'Agostino, M. (2019). Prevalence of Norovirus in produce sold at retail in the United Kingdom. *Food microbiology*, 79, 85-89.
- DEFRA. (2020). Food Statistics in your pocket: Global and UK supply. Department for Environment Food & Rural Affairs.
- Dolan, C., & Humphrey, J. (2000). Governance and trade in fresh vegetables: the impact of UK supermarkets on the African horticulture industry. *Journal of development studies*, 37(2), 147-176.

Formatted: Justified

Formatted: Font: (Default) Arial

Field Code Changed

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

1114 [Dos Santos, S. F., Cardoso, R. d. C. V., Borges, Í. M. P., e Almeida, A. C., Andrade, E. S.,](#)  
1115 [Ferreira, I. O., & do Carmo Ramos, L. \(2020\). Post-harvest losses of fruits and](#)  
1116 [vegetables in supply centers in Salvador, Brazil: Analysis of determinants, volumes](#)  
1117 [and reduction strategies. \*Waste Management\*, 101, 161-170.](#)

1118 [ECDC/EFSA. \(2021\). Multi-country outbreak of Salmonella Enteritidis sequence type \(ST\)11](#)  
1119 [infections linked to poultry products in the EU/EEA and the United Kingdom – 25](#)  
1120 [February 2021. Stockholm: ECDC/EFSA; 2021. European Centre for Disease](#)  
1121 [Prevention and Control \(ECDC\) and European Food Safety Authority \(EFSA\).](#)

1122 [EFSA/ECDC. \(2019\). Scientific report on the European Union one health 2018. Zoonoses](#)  
1123 [report. European Centre for Disease Prevention and Control \(ECDC\) and European](#)  
1124 [Food Safety Authority \(EFSA\). \*EFSA J.\*, 17, 5926.](#)

1125 [EFSA/ECDC. \(2021\). The European Union One Health 2019 Zoonoses Report. European](#)  
1126 [Food Safety Authority \(EFSA\) and European Centre for Disease Prevention and](#)  
1127 [Control \(ECDC\). \*EFSA Journal\* 19, 6406.](#)

1128 [EFSA/ECDC. \(2018\). Multi-country outbreak of Salmonella Agona infections possibly linked](#)  
1129 [to ready-to-eat food. European Food Safety Authority \(EFSA\) and European Centre](#)  
1130 [for Disease Prevention and Control \(ECDC\). In: Wiley Online Library.](#)

1131 [Emberland, K., Ethelberg, S., Kuusi, M., Vold, L., Jensvoll, L., Lindstedt, B., Nygård, K., Kjelsø,](#)  
1132 [C., Torpdahl, M., & Sørensen, G. \(2007\). Outbreak of \*Salmonella\* Weltevreden](#)  
1133 [infections in Norway, Denmark and Finland associated with alfalfa sprouts, July-](#)  
1134 [October 2007. \*Weekly releases \(1997–2007\)\*, 12\(48\), 3321.](#)

1135 [European Commission. \(2014\). Commission implementing decision of 13 February 2014](#)  
1136 [suspending temporarily imports from Bangladesh of foodstuffs containing or](#)  
1137 [consisting of betel leaves \('Piper betle'\) Available from: \[https://eur-\]\(https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32014D0088&from=DA\)](#)  
1138 [lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32014D0088&from=DA](#)  
1139 [Accessed 25.02.2021.](#)

1140 [European Commission. \(2015\). Commission implementing regulation \(EU\) 2015/943 of 18](#)  
1141 [June 2015. \*Official Journal of the European Union\*\(154\), 8-9.](#)

1142 [European Commission \(2016\). RASFF — The Rapid Alert System for Food and Feed — 2015](#)  
1143 [annual report. Available from: \[https://op.europa.eu/en/publication-detail/-\]\(https://op.europa.eu/en/publication-detail/-/publication/98e04cfc-aa54-11e6-aab7-01aa75ed71a1/language-en/format-PDF/source-174743070\)](#)  
1144 [/publication/98e04cfc-aa54-11e6-aab7-01aa75ed71a1/language-en/format-](#)  
1145 [PDF/source-174743070](#) Accessed 19.02.2021.

1146 [European Commission. \(2020a\). Commission implementing regulation \(EU\) 2020/625 of 6](#)  
1147 [May 2020. \*Official Journal of the European Union\*\(63\), 13.](#)

1148 [European Commission. \(2020b\). Monitoring EU Agri-Food Trade: Developments in 2019.](#)  
1149 [Available from: \[https://ec.europa.eu/info/news/2019-record-year-eu-agri-food-trade-\]\(https://ec.europa.eu/info/news/2019-record-year-eu-agri-food-trade-2020-mar-27\_en#moreinfo\)](#)  
1150 [2020-mar-27\\_en#moreinfo](#) Accessed 12.02.2021.

1151 [European Food Safety Authority. \(2021\). The European Union One Health 2019 Zoonoses](#)  
1152 [Report. \*EFSA Journal\*, 19\(2\).](#)

1153 [Fakruddin, M., Sultana, R., Hossain, M. N., Rahaman, M. M., Islam, M. K., & Ahmed, M. M.](#)  
1154 [\(2017\). Occurrence of ingress of \*Salmonella\* spp. in Betel leaf \(Piper betle L.\).](#)  
1155 [International Journal of Food Contamination, 4\(1\), 1-10.](#)

1156 [Ferrari, R. G., Rosario, D. K., Cunha-Neto, A., Mano, S. B., Figueiredo, E. E., & Conte-Junior,](#)  
1157 [C. A. \(2019\). Worldwide epidemiology of \*Salmonella\* serovars in animal-based foods:](#)  
1158 [a meta-analysis. \*Applied and environmental microbiology\*, 85\(14\).](#)

1159 [Food and Agricultural Organisation. \(2018\). Addressing the food safety challenges of the Asia-](#)  
1160 [Pacific region. 34th Session of the FAO Regional Conference for Asia and the Pacific.](#)  
1161 [Available from: <http://www.fao.org/3/MV819en/mv819en.pdf> Accessed 24.02.2021.](#)

1162 [Food and Agriculture Organization. \(2019\). Foodborne illnesses in Asia-Pacific 'needlessly'](#)  
1163 [sicken 275 million annually and threaten trade. Retrieved from](#)  
1164 [http://www.fao.org/asiapacific/news/detail-](#)  
1165 [events/en/c/1197008/#:~:text=In%20the%20Asia%2DPacific%20region,of%20Zero%20Hunger%20by%202030,.](#)

Formatted: Font: (Default) Cambria Math

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Cambria Math

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Cambria Math

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial



1167 [Fu, H., Yu, P., Liang, W., Kan, B., Peng, X., & Chen, L. \(2020\). Virulence, Resistance, and](#)  
1168 [Genomic Fingerprint Traits of \*Vibrio cholerae\* Isolated from 12 Species of Aquatic](#)  
1169 [Products in Shanghai, China. \*Microbial Drug Resistance\*, 26\(12\), 1526-1539.](#)  
1170 [Ghosh, D. \(2014\). Food safety regulations in Australia and New Zealand food standards.](#)  
1171 [Journal of the Science of Food and Agriculture, 94\(10\), 1970-1973.](#)  
1172 [Jaffee, S., Henson, S., Unnevehr, L., Grace, D., & Cassou, E. \(2018\). \*The safe food\*](#)  
1173 [imperative: Accelerating progress in low-and middle-income countries: World Bank](#)  
1174 [Publications.](#)  
1175 [Kaakoush, N. O., Castaño-Rodríguez, N., Mitchell, H. M., & Man, S. M. \(2015\). Global](#)  
1176 [epidemiology of \*Campylobacter\* infection. \*Clinical microbiology reviews\*, 28\(3\), 687-](#)  
1177 [720.](#)  
1178 [Kantama, L., & Jayanetra, P. \(1996\). \*Salmonella enteritidis\* outbreak in Thailand: study by](#)  
1179 [random amplified polymorphic DNA \(RAPD\) analysis. \*The Southeast Asian journal of\*](#)  
1180 [tropical medicine and public health](#), 27(1), 119-125.  
1181 [Keisam, S., Tuikhar, N., Ahmed, G., & Jeyaram, K. \(2019\). Toxigenic and pathogenic potential](#)  
1182 [of enteric bacterial pathogens prevalent in the traditional fermented foods marketed in](#)  
1183 [the Northeast region of India. \*International journal of food microbiology\*, 296, 21-30.](#)  
1184 [Kim, K., Kim, S., & Park, C.-Y. \(2020\). Food Security in Asia and the Pacific amid the COVID-](#)  
1185 [19 Pandemic. 1-15.](#)  
1186 [Kowalska, A., & Manning, L. \(2020\). Using the rapid alert system for food and feed: potential](#)  
1187 [benefits and problems on data interpretation. \*Critical reviews in food science and\*](#)  
1188 [nutrition](#), 1-14.  
1189 [Kumar, R., & Lalitha, K. V. \(2013\). Prevalence and molecular characterization of \*Vibrio\*](#)  
1190 [cholerae O1, non-O1 and non-O139 in tropical seafood in Cochin, India. \*Foodborne\*](#)  
1191 [pathogens and disease](#), 10(3), 278-283.  
1192 [Kumari, S., & Sarkar, P. K. \(2014\). Prevalence and characterization of \*Bacillus cereus\* group](#)  
1193 [from various marketed dairy products in India. \*Dairy science & technology\*, 94\(5\), 483-](#)  
1194 [497.](#)  
1195 [Lee, Y., Choi, Y., Lee, S., Lee, H., Kim, S., Ha, J., Lee, J., Oh, H., Kim, Y., & Yoon, Y. \(2019\).](#)  
1196 [Occurrence of pathogenic \*Vibrio parahaemolyticus\* in seafood distribution channels](#)  
1197 [and their antibiotic resistance profiles in S. Korea. \*Letters in applied microbiology\*,](#)  
1198 [68\(2\), 128-133.](#)  
1199 [Lemoine, T., Germanetto, P., & Giraud, P. \(1999\). Toxi-infection alimentaire à \*V.\*](#)  
1200 [parahaemolyticus. \*Bull Epidémiol Hebdo\*, 10, 37-38.](#)  
1201 [Machado-Moreira, B., Richards, K., Brennan, F., Abram, F., & Burgess, C. M. \(2019\). Microbial](#)  
1202 [contamination of fresh produce: what, where, and how? \*Comprehensive reviews in\*](#)  
1203 [food science and food safety](#), 18(6), 1727-1750.  
1204 [Makendi, C., Page, A. J., Wren, B. W., Le Thi Phuong, T., Clare, S., Hale, C., Goulding, D.,](#)  
1205 [Klemm, E. J., Pickard, D., & Okoro, C. \(2016\). A phylogenetic and phenotypic analysis](#)  
1206 [of \*Salmonella enterica\* serovar Weltevreden, an emerging agent of diarrheal disease](#)  
1207 [in tropical regions. \*PLoS neglected tropical diseases\*, 10\(2\), e0004446.](#)  
1208 [Mangla, S. K., Bhattacharya, A., Yadav, A. K., Sharma, Y. K., Ishizaka, A., Luthra, S., &](#)  
1209 [Chakraborty, R. \(2021\). A framework to assess the challenges to food safety initiatives](#)  
1210 [in an emerging economy. \*Journal of Cleaner Production\*, 284, 124709.](#)  
1211 [Martinez-Urtaza, J., Simental, L., Velasco, D., DePaola, A., Ishibashi, M., Nakaguchi, Y.,](#)  
1212 [Nishibuchi, M., Carrera-Flores, D., Rey-Alvarez, C., & Pousa, A. \(2005\). Pandemic](#)  
1213 [vibrio parahaemolyticus O3: K6, Europe. \*Emerging infectious diseases\*, 11\(8\), 1319.](#)  
1214 [McLauchlin, J., Aird, H., Andrews, N., Chattaway, M., De Pinna, E., Elviss, N., Jørgensen, F.,](#)  
1215 [Larkin, L., & Willis, C. \(2019\). Public health risks associated with \*Salmonella\*](#)  
1216 [contamination of imported edible betel leaves: analysis of results from England, 2011–](#)  
1217 [2017. \*International journal of food microbiology\*, 298, 1-10.](#)  
1218 [McLauchlin, J., Aird, H., Charlett, A., Chattaway, M., Elviss, N., Hartman, H., Jenkins, C.,](#)  
1219 [Jørgensen, F., Larkin, L., & Sadler-Reeves, L. \(2018\). Imported edible leaves collected](#)  
1220 [at retail sale in England during 2017 with an emphasis on betel and curry leaves:](#)  
1221 [microbiological quality with respect to \*Salmonella\*, Shiga-toxin-producing \*E. coli\*](#)

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Cambria Math

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Cambria Math

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Cambria Math

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Cambria Math

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

(STEC) and levels of *Escherichia coli*, *Journal of applied microbiology*, 125(4), 1175-1185.

Mensah, L. D., & Julien, D. (2011). Implementation of food safety management systems in the UK. *Food Control*, 22(8), 1216-1225.

Nakaguchi, Y. (2013). Contamination with *Vibrio parahaemolyticus* and its virulent strains in seafood marketed in Thailand, Vietnam, Malaysia, and Indonesia from 2008 to 2011. *Tropical Medicine and Health*, 2011-2006.

Nasheri, N., Vester, A., & Petronella, N. (2019). Foodborne viral outbreaks associated with frozen produce. *Epidemiology & Infection*, 147.

Nath, T. K., & Inoue, M. (2009). Sustainability attributes of a small-scale betel leaf agroforestry system: a case study in north-eastern hill forests of Bangladesh. *Small-scale Forestry*, 8(3), 289-304.

Nguyen, T. K., Bui, H. T., Truong, T. A., Lam, D. N., Ikeuchi, S., Ly, L. K. T., Hara-Kudo, Y., Taniguchi, T., & Hayashidani, H. (2021). Retail fresh vegetables as a potential source of *Salmonella* infection in the Mekong Delta, Vietnam. *International journal of food microbiology*, 341, 109049.

Office of National Statistics. (2011). UK Census 2011. Available online: <https://www.ons.gov.uk/census/2011census>.

Ortega, D. L. a. T., D.L. (2017). Demand for food safety in emerging and developing countries: A research agenda for Asia and Sub-Saharan Africa. *Journal of Agribusiness in Developing and Emerging Economies*(7), 21-34.

Park, S., Szonyi, B., Gautam, R., Nightingale, K., Anciso, J., & Ivanek, R. (2012). Risk factors for microbial contamination in fruits and vegetables at the preharvest level: a systematic review. *Journal of food protection*, 75(11), 2055-2081.

Pham, H. V., & Dinh, T. L. (2020). The Vietnam's food control system: Achievements and remaining issues. *Food Control*, 108, 106862.

Reddy, A. A., Ricart, S., & Cadman, T. (2020). Driving factors of food safety standards in India: learning from street-food vendors' behaviour and attitude. *Food Security*, 1-17.

Rezaei, M., & Liu, B. (2017). Food loss and waste in the food supply chain. *International Nut and Dried Fruit Council: Reus, Spain*, 26-27.

Rivas, L., Strydom, H., Paine, S., Wang, J., & Wright, J. (2021). Yersiniosis in New Zealand. *Pathogens*, 10(2), 191.

Singh, B. R., Singh, M., Singh, P., Babu, N., Chandra, M., & Agarwal, R. K. (2006). Prevalence of multidrug-resistant *Salmonella* on ready-to-eat betel leaves (Paan) and in water used for soaking betel leaves in North Indian cities. *Journal of food protection*, 69(2), 288-292.

Sodagari, H. R., Wang, P., Robertson, I., Habib, I., & Sahibzada, S. (2020). Non-Typhoidal *Salmonella* at the Human-Food-of-Animal-Origin Interface in Australia. *Animals*, 10(7), 1192.

Somarin, Y. M., Odeyemi, O. A., & Ateba, C. N. (2021). *Salmonella* is the most common foodborne pathogen in African food exports to the European Union: Analysis of the Rapid Alert System for Food and Feed (1999–2019). *Food Control*, 123, 107849.

Sugiri, Y. D., Goelz, G., Meeyam, T., Baumann, M. P., Kleer, J., Chaisowwong, W., & Alter, T. (2014). Prevalence and antimicrobial susceptibility of *Listeria monocytogenes* on chicken carcasses in Bandung, Indonesia. *Journal of food protection*, 77(8), 1407-1410.

Tähtkääpää, S., Majjala, R., Korkeala, H., & Nevas, M. (2015). Patterns of food frauds and adulterations reported in the EU rapid alert system for food and feed and in Finland. *Food Control*, 47, 175-184.

Ullah, A. S., Tani, M., Tsuchiya, J., Rahman, A. M., & Rahman, Z. M. (2020). Impact of Betel Leaf Cultivation on the Protected Forest Area of Teknaf Peninsula, Bangladesh. *Small-scale Forestry*, 19(3), 335-355.

Vilain, S., Luo, Y., Hildreth, M. B., & Brözel, V. S. (2006). Analysis of the life cycle of the soil saprophyte *Bacillus cereus* in liquid soil extract and in soil. *Applied and environmental microbiology*, 72(7), 4970-4977.

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

1277 [Vital, P. G., Dimasuay, K. G. B., Widmer, K. W., & Rivera, W. L. \(2014\). Microbiological quality](#)  
1278 [of fresh produce from open air markets and supermarkets in the Philippines. \*The\*](#)  
1279 [Scientific World Journal, 2014.](#)

1280 [Wadamori, Y., Gooneratne, R., & Hussain, M. A. \(2017\). Outbreaks and factors influencing](#)  
1281 [microbiological contamination of fresh produce. \*Journal of the Science of Food and\*](#)  
1282 [Agriculture, 97\(5\), 1396-1403.](#)

1283 [Wang, S., Weller, D., Falardeau, J., Strawn, L. K., Mardones, F. O., Adell, A. D., & Switt, A. I.](#)  
1284 [M. \(2016\). Food safety trends: from globalization of whole genome sequencing to](#)  
1285 [application of new tools to prevent foodborne diseases. \*Trends in Food Science &\*](#)  
1286 [Technology, 57, 188-198.](#)

1287 [Wong, H.-C., Chen, M.-C., Liu, S.-H., & Liu, D.-P. \(1999\). Incidence of highly genetically](#)  
1288 [diversified \*Vibrio parahaemolyticus\* in seafood imported from Asian countries.](#)  
1289 [International journal of food microbiology, 52\(3\), 181-188.](#)

1290 [World Health Organization. \(2015\). WHO estimates of the global burden of foodborne](#)  
1291 [diseases: foodborne disease burden epidemiology reference group 2007-2015; World](#)  
1292 [Health Organization.](#)

1293 [World Health Organization. \(2019\). Shiga Toxin-producing Escherichia Coli \(STEC\) and Food:](#)  
1294 [Attribution Characterization and Monitoring, \(Vol. 19\): World Health Organization.](#)

1295 [World Health Organization. \(2020a\). Food Safety. Retrieved from \[https://www.who.int/news-\]\(https://www.who.int/news-room/fact-sheets/detail/food-safety\)](#)  
1296 [room/fact-sheets/detail/food-safety.](#)

1297 [World Health Organization. \(2020b\). Framework for action on food safety in the WHO South-](#)  
1298 [East Asia Region. New Delhi: World Health Organization, Regional Office for South-](#)  
1299 [East Asia.](#)

1300 [World Health Organization. \(2020c\). \*Salmonella\* – non-typhoidal. Available from:](#)  
1301 [https://www.who.int/news-room/fact-sheets/detail/salmonella-\(non-typhoidal\)](#)  
1302 [Accessed 25.02.2021.](#)

1303 [Worldometer. \(2021\). Current World Population. Retrieved from](#)  
1304 [https://www.worldometers.info/world-population/#ref-1.](#)

1305 [Xu, X., Cheng, J., Wu, Q., Zhang, J., & Xie, T. \(2016\). Prevalence, characterization, and](#)  
1306 [antibiotic susceptibility of \*Vibrio parahaemolyticus\* isolated from retail aquatic products](#)  
1307 [in North China. \*BMC microbiology\*, 16\(1\), 1-9.](#)

1308 [Yu, P., Zaleski, A., Li, Q., He, Y., Mapili, K., Pruden, A., Alvarez, P. J., & Stadler, L. B. \(2018\).](#)  
1309 [Elevated levels of pathogenic indicator bacteria and antibiotic resistance genes after](#)  
1310 [Hurricane Harvey's flooding in Houston. \*Environmental science & technology letters\*,](#)  
1311 [5\(8\), 481-486.](#)

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Not Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Not Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial

Formatted: Font: (Default) Arial, Italic

Formatted: Font: (Default) Arial

1313 **Supplementary Table 1**

1314 RASFF notifications of pathogenic microorganisms in food originating from the Asia-Pacific Region (n=1873) from 2020 - 2020 based on  
1315 notifying country.

Pathogen genus	Notifying country											Total (%)
	United Kingdom	Italy	Norway	Netherlands	Germany	Finland	Greece	Spain	Denmark	Sweden	Others	
<i>Salmonella</i>	515	95	102	172	137	122	80	47	30	44	216	1560(73.55%)
<i>Vibrio</i>	4	87	124	4	10	7	2	8	24	12	27	309(14.57%)
<i>Bacillus</i>	36	19	1	3	13	9	0	0	0	0	11	92(4.34%)
<i>Escherichia</i>	27	2	8	1	1	1	0	0	2	1	4	47(2.22%)
<i>Norovirus</i>	0	4	1	1	3	1	0	20	3	1	8	42(1.98%)
<i>Not specified</i>	2	11	0	1	1	0	0	0	1	5	7	28(1.32%)
<i>Clostridium</i>	2	18	0	0	1	0	0	0	0	0	1	22(1.04%)
<i>Campylobacter</i>	0	0	0	0	0	0	0	0	4	1	1	6(0.28%)
<i>Staphylococcus</i>	0	0	2	0	0	0	0	0	1	0	0	3(0.14%)
<i>Enterococcus</i>	0	1	0	0	0	0	0	0	1	0	0	2(0.09%)
<i>Hepatitis</i>	0	1	0	0	0	0	0	0	0	0	1	2(0.09%)
<i>Pseudomonas</i>	0	0	0	0	2	0	0	0	0	0	0	2(0.09%)
<i>Cronobacter</i>	0	0	0	1	0	0	0	0	0	0	0	1(0.05%)
Fungi	0	0	0	0	0	0	0	0	0	0	1	1(0.05%)
<i>Plesiomonas</i>	1	0	0	0	0	0	0	0	0	0	0	1(0.05%)
<i>Shigella</i>	0	0	0	0	0	0	0	0	1	0	0	1(0.05%)
<i>Yersinia</i>	0	1	0	0	0	0	0	0	0	0	0	1(0.05%)
<i>Streptococci</i>	0	0	0	0	0	1	0	0	0	0	0	1(0.05%)
<b>Total</b>	<b>587</b>	<b>239</b>	<b>238</b>	<b>183</b>	<b>168</b>	<b>141</b>	<b>82</b>	<b>75</b>	<b>67</b>	<b>64</b>	<b>277</b>	<b>2121</b>

1316 **Other\_N: Other notifying countries:** Cyprus, Austria, Slovenia, Ireland, Lithuania, Iceland, Romania, Switzerland, Bulgaria, Croatia, Estonia,  
1317 Hungary, Latvia, Luxembourg, Slovakia, Sri Lanka. **Other\_O: Other originating countries:** Australia, Cambodia, New Zealand, Pakistan,  
1318 Philippines, Hong Kong, Taiwan, Singapore, South Korea, Afghanistan, Nepal. The total number of pathogens (n=2121) exceeds the total  
1319 number of notifications (n=1873) because of the co-occurrence of pathogens.

1320 **Supplementary Table 2**

1321 Pathogens associated with notifications and their diversities in Crustaceans and related  
1322 products from Asia and Pacific (2000 - 2020).

Country of origin	Notifications (n)	Pathogens	No of pathogens	Crustaceans and related products	
Bangladesh	38	<i>Salmonella</i> Bareilly, <i>Salmonella</i> Brunei, <i>Salmonella</i> enterica, <i>Salmonella</i> spp, <i>Vibrio cholerae</i> , <i>Vibrio parahaemolyticus</i>	43	Shrimps	Formatted: Font: Not Italic
		<i>Salmonella</i> spp, <i>Vibrio cholerae</i> , <i>Vibrio parahaemolyticus</i>	5	Prawns	Formatted: Font: Not Italic
China	12	<i>Salmonella enterica</i> ser. Newport, <i>Vibrio mimicus</i> , <i>Vibrio parahaemolyticus</i> , <i>Vibrio cholerae</i>	9	Crayfish	Formatted: Font: Not Italic
		<i>Salmonella</i> spp	2	Crab	Formatted: Font: Not Italic
		<i>Vibrio parahaemolyticus</i>	1	Shrimps	
India	49	aerobic mesophiles, <i>Enterococcus</i> spp, <i>Pseudomonas</i> spp, <i>Salmonella enterica</i> , <i>Salmonella</i> spp, <i>Salmonella</i> Weltevreden, <i>Vibrio alginolyticus</i> , <i>Vibrio cholerae</i> , <i>Vibrio fluvialis</i> , <i>Vibrio parahaemolyticus</i> , <i>Vibrio vulnificus</i>	60	Shrimps	Formatted: Font: Not Italic
					Formatted: Font: Not Italic
		<i>Salmonella enterica</i> , <i>Salmonella</i> spp, <i>Vibrio cholerae</i> , <i>Vibrio parahaemolyticus</i>	7	Prawns	Formatted: Font: Not Italic
		<i>Vibrio cholerae</i>	1	Crab	
Indonesia	26	<i>Salmonella enterica</i> subsp. houtenae (VI), <i>Salmonella</i> Lexington, <i>Salmonella</i> spp, <i>Salmonella</i> Wandsworth, <i>Vibrio alginolyticus</i> , <i>Vibrio cholerae</i> , <i>Vibrio parahaemolyticus</i> , <i>Vibrio</i> spp, <i>Vibrio vulnificus</i> ,	21	Shrimps	Formatted: Font: Not Italic
					Formatted: Font: Not Italic
		<i>Plesiomonas shigelloides</i> , <i>Salmonella</i> spp, <i>Vibrio parahaemolyticus</i> , <i>Vibrio cholerae</i>	12	Prawns	Formatted: Font: Not Italic
Malaysia	65	<i>Salmonella enterica</i> , <i>Salmonella</i> Give, <i>Salmonella</i> spp, <i>Vibrio cholerae</i> , <i>Vibrio parahaemolyticus</i> , <i>Vibrio vulnificus</i> ,	48	Shrimps	Formatted: Font: Not Italic
		<i>Salmonella enterica</i> , <i>Salmonella</i> spp, <i>Vibrio cholerae</i> , <i>Vibrio parahaemolyticus</i>	17	Prawns	Formatted: Font: Not Italic
		<i>Vibrio parahaemolyticus</i>	1	shellfish	
Sri Lanka, Pakistan	4	<i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	4	Crab	
		<i>Vibrio cholerae</i> , aerobic mesophiles, <i>Salmonella</i> spp	3	Shrimps	Formatted: Font: Not Italic
Thailand	23	<i>Salmonella enterica</i> , <i>Vibrio cholerae</i> , <i>Vibrio parahaemolyticus</i> , <i>Vibrio vulnificus</i>	21	Shrimps	
		<i>Clostridium perfringens</i> , <i>Clostridium</i> sulphite reducer	3	Shrimp paste	Formatted: Font: Not Italic
		<i>Vibrio cholerae</i>	1	Crab	
		<i>Vibrio parahaemolyticus</i>	1	Prawns	

1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343

Country of origin	Notifications (n)	Pathogens	No of pathogens	Crustaceans and related products
Vietnam	48	<i>Escherichia coli</i> , <i>Salmonella enterica</i> , <i>Salmonella enterica</i> ser. Weltevreden, <i>Salmonella</i> Kentucky , <i>Salmonella</i> Newport, <i>Salmonella</i> Oranienburg, <i>Salmonella</i> spp, <i>Salmonella</i> Virchow , <i>Salmonella</i> Welshimeri, <i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i> , <i>Vibrio parahaemolyticus</i> , <i>Vibrio</i> spp, <i>Vibrio vulnificus</i>	55	Shrimps
		<i>Salmonella enterica</i> , <i>Salmonella</i> spp, <i>Vibrio cholerae</i> , <i>Vibrio parahaemolyticus</i>	6	Prawns
		<i>Clostridium</i> sulphite reducer	1	Shrimp paste
		<i>Salmonella enterica</i> ser. Hvittefoss	1	Crab
		<i>Vibrio parahaemolyticus</i>	1	Clams
		<i>Salmonella enterica</i>	1	Crustaceans

- Formatted: Font: Not Italic
- Formatted: Font: Not Italic
- Formatted: Font: Not Italic
- Formatted: Font: Not Italic
- Formatted: Font: Not Italic
- Formatted: Font: Not Italic
- Formatted: Font: Not Italic
- Formatted: Font: Not Italic
- Formatted: Font: Not Italic
- Formatted: Font: Not Italic
- Formatted: Font: Not Italic

1344 **Supplementary Table 3**

1345 Pathogens associated with notifications and their diversities in Fruits and Vegetables from

1346 Asia and Pacific (2000 - 2020).

Country of origin	No of notifications	Pathogen	Count of Pathogen species	Food	
Afghanistan	1	<i>Salmonella enterica</i> <del>ser-ser.</del> Agona	1	mulberry	Formatted
Australia	1	<i>Salmonella enterica</i> <del>ser-ser.</del> Chester	1	mushroom	Formatted
Bangladesh	151	<i>Salmonella enterica</i> , <i>Salmonella</i> Jerusalem, <i>Salmonella</i> spp, <i>Salmonella</i> Typhimurium, <i>Salmonella</i> Virchow	149	betel leaves	Formatted
Cambodia	6	<i>Salmonella enterica</i>	3	paan leaves	Formatted: Font: Italic
		<i>Salmonella</i> spp	2	Basil	Formatted: Font: Italic
		<i>Salmonella enterica</i> , <i>Salmonella enterica</i> <del>ser-ser.</del> Stanley	2	betel leaves	Formatted
		<i>Salmonella</i> Hvittingfoss, <i>Salmonella</i> Brunei, <i>Salmonella enteritidis</i> , <i>Salmonella</i> Typhimurium, <i>Salmonella</i> Weltevreden	5	vine leaves	Formatted
		<i>Salmonella</i> spp	1	mushroom	Formatted: Font: Italic
Chile	1	<i>Salmonella</i> spp	1	mushroom	Formatted: Font: Italic
China	39	<i>Salmonella</i> Enteritidis	1	beans	Formatted
				sprouts	
		<i>Escherichia coli</i> , <i>Salmonella</i> spp	2	Cep powde	Formatted: Font: Italic
		<i>Salmonella</i> Thompson	1	Chanterelle powder	Formatted
		<i>Bacillus cereus</i>	1	Cheese	Formatted: Font: Italic
		<i>Salmonella</i> Rissen	1	chlorella	Formatted
		<i>Salmonella</i> spp	1	goji berries	Formatted: Font: Italic
		<i>Bacillus cereus</i> , <i>Clostridium perfringens</i> , <i>Clostridium sporulated sulphite reducer</i> , <i>Clostridium</i> spp, <i>Salmonella</i> spp, aerobic mesophiles, moulds, <i>Salmonella enterica</i>	18	mushroom	Formatted: Font: Italic
		Norovirus, <del>Norovirus</del> Norovirus GII	5	raspberries	Formatted
		<del>Norovirus</del> Norovirus GI-GI & GII	2	seaweed salad	Formatted
		<i>Bacillus cereus</i>	1	soy cheese	Formatted: Font: Italic
		<i>Salmonella</i> Tennessee	1	soy protein	Formatted
		Hepatitis A, Norovirus, Norovirus, <del>GI-GI</del> & GII	5	strawberries	Formatted
		<i>Bacillus cereus</i>	2	tofu	Formatted: Font: Italic
Hong Kong	2	<i>Salmonella</i> Senftenberg	1	mushrooms	Formatted: Font: Italic
		<i>Bacillus cereus</i>	1	tofu	Formatted: Font: Italic
India	192	<i>Salmonella enterica</i> , <i>Salmonella</i> spp	184	betel leaves	Formatted: Font: Italic
		<i>Salmonella enterica</i>	2	curry leaves	Formatted: Font: Italic
		<i>Salmonella</i> spp	1	fennel seed	Formatted: Font: Italic
		<i>Salmonella</i> spp	1	moringa leaves	Formatted: Font: Italic
		<i>Salmonella agona</i> , <i>Salmonella</i> Westminster	2	onions	Formatted: Font: Italic
		<i>Salmonella infantis</i>	1	sorrel leaves	Formatted: Font: Italic
					Formatted: Font: Italic



Country of origin	No of notifications	Pathogen	Count of Pathogen species	Food	
Laos	1	<i>Escherichia coli</i> , <i>Salmonella Brunei</i> , <i>Salmonella Weltevreden</i>	3	vine leaves	Formatted: Font: Italic
Malaysia	15	<i>Salmonella enterica</i> <del>ser.</del> <i>ser.</i> <i>Aberdeen</i>	1	beans	Formatted: Font: Italic
		<i>Escherichia coli</i> , <i>Salmonella Colindale</i> , <i>Salmonella group D</i> , <i>Salmonella spp</i> , <i>Salmonella Weltevreden</i> , <i>Vibrio cholerae</i>	14	betel leaves	Formatted: Font: Italic
					Formatted: Font: Italic
Pakistan	1	<i>Salmonella Weltevreden</i> , <i>Vibrio cholerae</i>	2	Alfalfa	Formatted: Font: Italic
Sri Lanka	7	<i>Escherichia coli</i> , <i>Salmonella enterica</i> , <i>Salmonella enterica</i> <del>ser.</del> <i>ser.</i> <i>Weltevreden</i> , <i>Salmonella spp</i>	5	betel leaves	Formatted: Font: Italic
		<i>Salmonella Weltevreden</i> , <i>Vibrio cholerae</i>	2	centella	Formatted: Font: Italic
		<i>Salmonella poona</i> , <i>Salmonella Weltevreden</i> , <i>Vibrio cholerae</i>	3	Spinach	Formatted: Font: Italic
Taiwan	1	<i>Bacillus cereus</i>	1	tofu	Formatted: Font: Italic
Thailand	81	<i>Salmonella anatum</i> , <i>Escherichia coli</i> , <i>Salmonella Stanley</i>	3	magosa	Formatted: Font: Italic
		<i>Salmonella spp</i> , <i>Salmonella Virchow</i>	2	Acacia	Formatted: Font: Italic
		<i>Campylobacter spp</i> , <i>Salmonella Bareilly</i>	2	Asparagus	Formatted: Font: Italic
		<i>Salmonella spp</i> , <i>Salmonella Weltevreden</i> , <i>Vibrio cholerae</i>	3	Banana leaves	Formatted: Font: Italic
		<i>Escherichia coli</i> , <i>Salmonella enterica</i> , <i>Salmonella Jerusalem</i> , <i>Salmonella spp</i>	16	betel leaves	Formatted: Font: Italic
		<i>Bacillus cereus</i> , <i>Salmonella Weltevreden</i> , <i>Vibrio cholerae</i>	3	black fungus	Formatted: Font: Italic
		<i>Salmonella Thompson</i> , <i>Salmonella spp</i>	1	broccoli	Formatted: Font: Italic
		<i>Salmonella spp</i>	3	Cabbage, Cha leaves	Formatted: Font: Italic
		<i>Campylobacter spp</i> , <i>Salmonella enterica</i> , <i>Salmonella spp</i> , <i>Salmonella typhimurium</i> , <i>Salmonella Weltevreden</i> , <i>Salmonella Zanzibar</i> , <i>Shigella sonnei</i> , <i>Vibrio cholerae</i>	8	Corn	Formatted: Font: Italic
		<i>Salmonella spp</i>	2	fresh lime and banana, Ipomea aquatica	Formatted: Font: Italic
		<i>Salmonella Stanley</i>	1	kaffir lime leaves	Formatted: Font: Italic
		<i>Salmonella Hvittingfoss</i> , <i>Salmonella infantis</i> , <i>Salmonella Zanzibar</i>	3	lemon grass	Formatted: Font: Italic
		<i>Salmonella Hadar</i> , <i>Salmonella Lexington</i>	2	lime leaves	Formatted: Font: Italic
		<i>Salmonella enterica</i> , <i>Salmonella enterica</i> <del>ser.</del> <i>ser.</i> <i>Hvittingfoss</i> , <i>Salmonella enterica</i> <del>ser.</del> <i>ser.</i> <i>Newport</i> , <i>Salmonella spp</i> , <i>Salmonella Stanley</i>	6	morning glory	Formatted: Font: Italic
		<i>Salmonella Hvittingfoss</i> , <i>Salmonella spp</i>	2	mushroom	Formatted: Font: Italic
		<i>Salmonella enterica</i> <del>ser.</del> <i>ser.</i> <i>Hvittingfoss</i>	1	neem leaves	Formatted: Font: Italic



Country of origin	No of notifications	Pathogen	Count of Pathogen species	Food	
		<i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	2	okra	Formatted
		<i>Salmonella</i> spp	1	ong choi	Formatted: Font: Italic
		<i>Salmonella</i> bBraenderup	1	Other vegetables	Formatted
		<i>Salmonella enterica</i>	1	paan leaves	Formatted: Font: Italic
		<i>Salmonella</i> aAltona, <i>Salmonella</i> Stanley	2	pak peaw	Formatted
		<i>Salmonella</i> Brunei	1	pak wan	Formatted
		<i>Salmonella enterica</i> ser-ser. Chester	3	pandang	Formatted
		<i>Salmonella enterica</i> ser-ser. Rubislaw		leaf	Formatted
		<i>Salmonella enterica</i> subsp. Salamae			Formatted: Font: Italic
		<i>Salmonella</i> Ndolo, <i>Salmonella</i> Aberdeen	12	Spinach	Formatted: Font: Italic
		<i>Salmonella</i> bBovismorbificans, <i>Salmonella enterica</i> , <i>Salmonella</i> group B, <i>Salmonella</i> group D, <i>Salmonella</i> Hvittingfoss, <i>Salmonella</i> Mbandaka, <i>Salmonella</i> Newport			Formatted
		<i>Campylobacter</i> spp, <i>Salmonella enterica</i> , <i>Salmonella</i> Hvittingfoss, <i>Salmonella</i> spp	5	spring onions	Formatted
		<i>Salmonella</i> aAnatum	1	tamarind leaves	Formatted
		<i>Salmonella</i> aAugustenborg	1	watercress	Formatted
		<i>Salmonella</i> Saint Paul, <i>Salmonella</i> Stanley	2	wildbetal leafbush	Formatted
Vietnam	38	<i>Bacillus cereus</i> , <i>Salmonella</i> aAugustenborg, <i>Salmonella enterica</i> ser-ser. Stanley, <i>Salmonella</i> spp, <i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	8	black fungus	Formatted
		<i>Salmonella</i> Brunei	1	celery	Formatted
		<i>Salmonella</i> eEnterica	1	coriander	Formatted
		<i>Salmonella</i> group B	1	morels	Formatted
		<i>Salmonella</i> spp	1	morning glory	Formatted
		<i>Bacillus cereus</i> , <i>Salmonella</i> Thompson, <i>Clostridium perfringens</i> , <i>Salmonella enterica</i> , <i>Salmonella enterica</i> ser-ser. Derby, <i>Salmonella enterica</i> ser-ser. Panama, <i>Salmonella enterica</i> ser-ser. Stanley, <i>Salmonella enterica</i> ser-ser. Virchow, <i>Salmonella</i> group B, <i>Salmonella</i> Java, <i>Salmonella</i> Meleagridis, <i>Salmonella</i> Rissen, <i>Salmonella</i> spp, <i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	29	mushroom	Formatted
		<i>Salmonella</i> Stanley	1	okra	Formatted
		<i>Campylobacter</i> spp	1	peppers	Formatted
		<i>Salmonella</i> spp	1	pineapple	Formatted
		<i>Salmonella</i> spp	1	Spinach	Formatted

Country of origin	No of notifications	Pathogen	Count of Pathogen species	Food
		<i>Salmonella</i> Lexington, <i>Salmonella</i> Weltevreden	2	coriander, houttuynia and celery

Formatted

1347 **Supplementary Table 4**

1348 Pathogens associated with notifications and their diversities in nuts and related products

1349 from Asia and Pacific (2000 - 2020).

Country of origin	Notifications (n)	Pathogens	No of pathogens	Nuts and related products
China	7	<i>Salmonella</i> Newport, <i>Salmonella</i> Stanley	2	peanuts
		<i>Salmonella</i> spp.	1	pine nuts
		<i>Bacillus cereus</i>	1	pumpkin seeds
		<i>Bacillus cereus</i>	1	soy cheese
		<i>Salmonella</i> Senftenberg, <i>Salmonella</i> spp., <i>Salmonella</i> Tennessee	3	sunflower seeds
India	253	<i>Salmonella</i> spp.	1	amaranth
		<i>Salmonella enterica</i> ser. Matadi	1	cashew
		<i>Salmonella</i> <del>e</del> Elomrane, <i>Salmonella enterica</i> ser. Bareilly	2	nigella seed
		<i>Escherichia coli</i> , <i>Salmonella</i> spp.	2	Coconut
		<i>Bacillus cereus</i> , <i>Escherichia coli</i> , <i>Salmonella</i> <del>m</del> Mbandaka, <i>Salmonella</i> Montevideo, <i>Salmonella</i> spp., <i>Salmonella</i> <del>a</del> Agona, <i>Salmonella</i> Amsterdam, <i>Salmonella</i> <del>a</del> Anatum, <i>Salmonella</i> Bareilly, <i>Salmonella</i> <del>b</del> Binza, <i>Salmonella</i> <del>B</del> Braenderup, <i>Salmonella</i> Chittagong, <i>Salmonella</i> <del>d</del> Dallgow, <i>Salmonella</i> <del>d</del> Drypool, <i>Salmonella enterica</i> , <i>Salmonella enterica</i> ser. Agona, <i>Salmonella enterica</i> ser. Amsterdam, <i>Salmonella enterica</i> ser. Bareilly, <i>Salmonella enterica</i> ser. Braenderup, <i>Salmonella enterica</i> ser. Derby, <i>Salmonella enterica</i> ser. Hvittingfoss, <i>Salmonella enterica</i> ser. Isangi, <i>Salmonella enterica</i> ser. Kentucky, <i>Salmonella enterica</i> ser. Kisil, <i>Salmonella enterica</i> ser. Livingstone, <i>Salmonella enterica</i> ser. Londo, <i>Salmonella enterica</i> ser. Montevideo, <i>Salmonella enterica</i> ser. Orion, <i>Salmonella enterica</i> ser. Ouakam, <i>Salmonella enterica</i> ser. Pensacola, <i>Salmonella enterica</i> ser. Schwarzengrund, <i>Salmonella enterica</i> ser. Senftenberg, <i>Salmonella enterica</i> ser. Typhimurium, <i>Salmonella enterica</i> ser. Umbadah, <i>Salmonella</i> <del>E</del> enteritidis, <i>Salmonella</i> <del>e</del> Gaminara, <i>Salmonella</i> group C1, <i>Salmonella</i> Hvittingfoss	262	Sesame seeds

Formatted

Formatted

Formatted: Font: Italic

Formatted: Font: Italic

Formatted

Formatted: Font: Italic

Formatted

Formatted

Formatted

Formatted

Country of origin	Notifications (n)	Pathogens	No of pathogens	Nuts and related products
		<i>Salmonella</i> Java, <i>Salmonella</i> Kastrup, <i>Salmonella</i> Kentucky, <i>Salmonella</i> Kristianstad, <i>Salmonella</i> Livingstone, <i>Salmonella</i> London, <i>Salmonella</i> Matopeni, <i>Salmonella</i> Molade, <i>Salmonella</i> Newport, <i>Salmonella</i> Ngili, <i>Salmonella</i> Orion, <i>Salmonella</i> Richmond, <i>Salmonella</i> Ruiru, <i>Salmonella</i> Senftenberg, <i>Salmonella</i> spp, <i>Salmonella</i> Tennessee, <i>Salmonella</i> Tilburg, <i>Salmonella</i> Urbana		
Indonesia	6	<i>Bacillus cereus</i> , <i>Enterococcus</i> spp, <i>Salmonella</i> Brunei, <i>Salmonella enterica</i> , <i>Salmonella enterica</i> ser. Senftenberg, <i>Salmonella</i> spp, <i>Salmonella</i> Westhampton, <i>Salmonella</i> Bareilly,	12	Coconut, desiccated coconut
Malaysia	2	<i>Salmonella</i> spp	3	Coconut, desiccated coconut
Pakistan	3	<i>Salmonella enterica</i> , <i>Salmonella</i> Hadar,	2	pine nuts
		<i>Bacillus cereus</i>	2	Sesame seeds
Philippines	2	<i>Salmonella</i> Mbandaka, <i>Salmonella</i> Albany, <i>Salmonella</i> Typhimurium, <i>Salmonella</i> Westhampton, <i>Salmonella</i> spp	5	Coconut
Singapore	1	<i>Salmonella enterica</i> ser. Weltevreden	2	Coconut
Sri Lanka	4	<i>Salmonella enterica</i>	2	coconut flour
		<i>Salmonella</i> Enteritidis	1	desiccated coconut
		<i>Salmonella</i> spp	1	poppy seeds
Thailand	1	<i>Salmonella</i> Lexington	1	Sesame seeds
Vietnam	1	<i>Salmonella enterica</i>	1	Coconut

1350

1351

1352

1353

1354

1355

1356

1357

1358

1359

1360 **Supplementary Table 5**

1361 Yearly RASFF notifications of pathogenic microorganisms in nuts, nut products and seeds originating from Asia-Pacific Region (n=1873).

Nuts, nut products and seeds						Sesame seeds		
Year	India	Indonesia	Thailand	Others	Total	India	Thailand	Total
2000	0	0	0	1	1	0	0	0
2001	3	0	1	2	6	3	1	4
2002	0	0	0	2	2	0	0	0
2003	1	0	0	2	3	1	0	1
2004	6	1	0	0	7	6	0	6
2005	10	1	0	1	12	10	0	10
2006	1	1	0	0	2	1	0	1
2007	4	0	0	1	5	4	0	4
2008	11	0	0	2	13	11	0	11
2009	13	0	0	2	15	13	0	13
2010	8	0	0	1	9	8	0	8
2011	6	0	0	0	6	6	0	6
2012	8	2	0	2	12	8	0	8
2013	5	0	0	2	7	5	0	5
2014	24	2	0	1	27	24	0	24
2015	68	1	0	1	70	67	0	67
2016	19	0	0	0	19	19	0	19
2017	20	1	0	0	21	18	0	18
2018	24	0	0	3	27	23	0	23
2019	17	0	0	0	17	17	0	17
2020	18	0	0	0	18	16	0	16
Total	266	9	1	23	299	260	1	261

1362 Others: Malaysia, Pakistan, Phillipines, Singapore, Sri Lanka, China, Vietnam

1363 **Supplementary Table 6**

1364 Pathogens associated with notifications and their diversities in herbs and spices from Asia and  
1365 Pacific (2000 - 2020).

1366

Country of origin	Notifications (n)	Pathogens	No of pathogens	Herbs and spices
Bangladesh	2	<i>Bacillus cereus</i>	2	other spice
Cambodia	3	<i>Salmonella</i> spp.	1	basil
China	35	<i>Salmonella enterica</i> , <i>Salmonella</i> Stanley	2	mint
		<i>Bacillus cereus</i>	2	garlic powder
		<i>Salmonella</i> Aberdeen, <i>Salmonella</i> spp.	3	Ginger powder
		<i>Bacillus cereus</i> , <i>Salmonella</i> Aequatoria, <i>Salmonella enterica</i> , <i>Salmonella enterica</i> ser. Enteritidis, <i>Salmonella</i> spp., <i>Salmonella</i> Thompson	12	ground pepper
		<i>Salmonella</i> group B, <i>Salmonella</i> spp.	2	other spice
		<i>Salmonella</i> spp.	16	paprika
		<i>Escherichia coli</i> , <i>Salmonella</i> spp., sulphite reducing anaerobes	3	parsley
		<i>Salmonella</i> spp.	1	red ground pepper
		<i>Salmonella</i> spp.	1	turmeric powder
				seasoning mix
India	56	<i>Salmonella enterica</i>	1	
		<i>Salmonella</i> <del>P</del> Poona, <i>Salmonella</i> spp.	4	coriander
		<i>Salmonella</i> <del>E</del> Derby	1	cumin powder
		<i>Salmonella</i> spp.	1	cumin seed
		<i>Salmonella</i> <del>E</del> Enteritidis, <i>Salmonella</i> spp.	2	curry leaves
		<i>Bacillus cereus</i> , <i>Salmonella</i> <del>A</del> Agona, <i>Salmonella</i> <del>B</del> Braenderup, <i>Salmonella enterica</i> , <i>Salmonella</i> spp.	6	Curry powder
		<i>Escherichia coli</i> , <i>Salmonella</i> spp.	2	fenugreek leaves

Country of origin	Notifications (n)	Pathogens	No of pathogens	Herbs and spices	Formatted: Font: Italic
		<i>Bacillus cereus</i> , <i>Clostridium perfringens</i> , <i>Salmonella</i> Ohio	4	ginger powder	Formatted: Font: Italic
					Formatted: Font: Italic
					Formatted: Font: Italic
					Formatted: Font: Italic
					Formatted: Font: Italic
					Formatted: Font: Italic
					Formatted: Font: Italic
		<i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	2	ginseng powder	Formatted: Font: Italic
					Formatted: Font: Italic
		<i>Bacillus cereus</i> , <i>Salmonella</i> group C, <i>Salmonella</i> spp	3	ground cumin	Formatted: Font: Italic
					Formatted: Font: Italic
		<i>Salmonella</i> Agona, <i>Salmonella</i> Bareilly, <i>Salmonella</i> Cubana	3	ground cumin and ground coriander	Formatted: Font: Italic
					Formatted: Font: Italic
		<i>Salmonella</i> spp	1	ground nutmeg	Formatted: Font: Italic
		<i>Bacillus cereus</i> , <i>Bacillus pumilus</i> , <i>Bacillus subtilis</i> , <i>Clostridium perfringens</i> , <i>Salmonella</i> Bareilly, <i>Salmonella enterica</i> <i>Salmonella</i> Give, <i>Salmonella</i> Newport, <i>Salmonella</i> Richmond, <i>Salmonella</i> spp	16	ground pepper	Formatted: Font: Italic
					Formatted: Font: Italic
		<i>Salmonella</i> Richmond	1	Indian ginseng powder	Formatted: Font: Italic
					Formatted: Font: Not Italic
		<i>Salmonella</i> spp	1	moringa leaves	Formatted: Font: Italic
					Formatted: Font: Italic
		<i>Salmonella enterica</i>	3	onion powder	Formatted: Font: Italic
		<i>Salmonella</i> spp			Formatted: Font: Italic
		<i>Salmonella</i> Butantan	1	other herbs	Formatted: Font: Italic
		<i>Bacillus cereus</i> , <i>Clostridium perfringens</i> , <i>Salmonella</i> Amsterdam, <i>Salmonella</i> spp, <i>Salmonella enterica</i>	4	other spices	Formatted: Font: Italic
					Formatted: Font: Italic
			1	sesame seeds	Formatted: Font: Italic
		<i>Salmonella</i> Senftenberg, <i>Salmonella</i> Amsterdam	2	turmeric powder	Formatted: Font: Italic
					Formatted: Font: Italic
		<i>Salmonella enterica</i>	1	ground peppers	Formatted: Font: Italic
					Formatted: Font: Italic
		<i>Salmonella enterica</i> , <i>Salmonella enterica</i> ser. Agona, <i>Salmonella</i> Senftenberg, <i>Salmonella</i> Livingstone, <i>Salmonella</i> spp, <i>Salmonella</i> Virchow	7	turmeric powder	Formatted: Font: Italic
					Formatted: Font: Italic
					Formatted: Font: Italic

Country of origin	Notifications (n)	Pathogens	No of pathogens	Herbs and spices	
Indonesia	4	<i>Bacillus cereus</i>	1	cinnamon powder	Formatted: Font: Italic
		<i>Salmonella</i> Aberdeen, <i>Salmonella</i> Albany, <i>Salmonella</i> Mbandaka	3	ground pepper	Formatted
					Formatted: Font: Italic
Laos	23	<i>Escherichia coli</i> , <i>Salmonella</i> Lexington, <i>Salmonella</i> spp.	4	basil	Formatted
		<i>Salmonella</i> spp.	1	chives	Formatted: Font: Italic
		<i>Salmonella</i> spp.	1	coriander	Formatted
		<i>Escherichia coli</i> , <i>Salmonella</i> Brunei, <i>Salmonella</i> spp., <i>Salmonella</i> Weltevreden	8	mint	Formatted
		<i>Salmonella</i> spp.	1	parsley	Formatted
		<i>Escherichia coli</i> , <i>Salmonella</i> Brunei, <i>Salmonella</i> Heidelberg, <i>Salmonella</i> spp., <i>Salmonella</i> Thompson, <i>Salmonella</i> typhimurium	12	perilla	Formatted
		<i>Escherichia coli</i> , <i>Salmonella</i> Aegona, <i>Salmonella</i> Bareilly, <i>Salmonella</i> Hvittefoss, <i>Salmonella</i> Javiana	8	piper lolot	Formatted
		<i>Escherichia coli</i> , <i>Salmonella</i> Hvittefoss, <i>Salmonella</i> Meleagridis, <i>Salmonella</i> Rissen, <i>Salmonella</i> spp.	6	praew leaf	Formatted
		<i>Salmonella enterica</i>	1	ground min	Formatted: Font: Italic
		<i>Salmonella</i> Bareilly, <i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	3	Curry powd	Formatted
Malaysia	3	<i>Salmonella</i> spp.	1	parsley and betel leaves	Formatted: Font: Italic
		<i>Salmonella enterica ser. Java</i>	1	betel leave	Formatted
Nepal	1	<i>Salmonella</i> eEdinburgh	1	ground pepper	Formatted
		<i>Bacillus cereus</i> , <i>Salmonella</i> Hvittefoss	2	other spice	Formatted
Pakistan	3				
Singapore	1	<i>Salmonella enterica</i>	1	ground pepper	Formatted: Font: Italic
Sri Lanka	8	<i>Bacillus cereus</i>	1	cinnamon powder	Formatted: Font: Italic
		<i>Salmonella</i> wWaycross, <i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	3	curry leave	Formatted
		Moulds, <i>Salmonella</i> Aegona, <i>Salmonella</i> Java, <i>Salmonella</i> Pparatyphi B	4	ground pepper	Formatted
		<i>Escherichia coli</i> , <i>Salmonella</i> spp.	2	moringa leaves	Formatted
		<i>Salmonella</i> spp.	1	other herbs	Formatted

Country of origin	Notifications (n)	Pathogens	No of pathogens	Herbs and spices	
Thailand	158	<i>Salmonella</i> spp.	1	betel leave	Formatted
		<i>Salmonella</i> spp.	2	pennywort	Formatted
		<i>Salmonella</i> Stanley			Formatted: Font: Italic
		<i>Salmonella</i> spp.	1	phak-phae	Formatted
		<i>Salmonella</i> Hvittingfoss, <i>Escherichia coli</i> , <i>Salmonella enterica</i> , <i>Salmonella</i> Hadar	10	acacia	Formatted
		<i>Salmonella</i> spp, <i>Salmonella</i> Stanley, <i>Salmonella</i> Virchow, <i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>			Formatted
		<i>Salmonella</i> spp	1	amaranth greens	Formatted: Font: Italic
		<i>Salmonella</i> spp	1	Asiatic pennywort	Formatted: Font: Italic
		<i>Salmonella</i> Thompson, <i>Escherichia coli</i> , <i>Salmonella</i> spp, <i>Salmonella</i> Agona, <i>Salmonella</i> Amsterdam, <i>Salmonella</i> arizonae, <i>Salmonella</i> augustinborg, <i>Salmonella</i> Bareilly, <i>Salmonella</i> Brunei, <i>Salmonella</i> Dublin, <i>Salmonella enterica</i> , <i>Salmonella</i> Goverdhan, <i>Salmonella</i> Java, <i>Salmonella enterica</i> subsp. Salamae, <i>Salmonella</i> group D, <i>Salmonella</i> Hvittingfoss, <i>Salmonella</i> Javiana, <i>Salmonella</i> Lexington, <i>Salmonella</i> Rubislaw, <i>Salmonella</i> Saint Paul, <i>Salmonella</i> spp, <i>Salmonella</i> Stanley, <i>Salmonella</i> Thompson, <i>Salmonella</i> Weltevreden	65	basil	Formatted
		<i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	2	betel leave	Formatted
		<i>Salmonella</i> spp	1	Cha leaves	Formatted
		<i>Salmonella</i> Augustenborg	1	chives	Formatted
		<i>Salmonella</i> Anatum, <i>Salmonella</i> Augustenborg, <i>Salmonella</i> Brunei, <i>Salmonella</i> Corvallis, <i>Salmonella enterica</i> , <i>Salmonella enterica</i> ser. Hvittingfoss, <i>Salmonella enterica</i> ser. Lexington, <i>Salmonella</i> Hvittingfoss, <i>Salmonella</i> Jerusalem, <i>Salmonella</i> Newport, <i>Salmonella</i> Rissen, <i>Salmonella</i> Rubislaw, <i>Salmonella</i> Singapore, <i>Salmonella</i> spp, <i>Salmonella</i> Virchow, <i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	24	coriander	Formatted
		<i>Salmonella</i> Augustenborg	1	doksadao	Formatted
		<i>Salmonella</i> Mbandaka	1	edible flower	Formatted
		<i>Salmonella</i> Weltevreden	1	galingale	Formatted



Country of origin	Notifications (n)	Pathogens	No of pathogens	Herbs and spices	
		<i>Salmonella</i> Hvittingfoss	2	ground pepper	Formatted
		<i>Salmonella</i> spp.			Formatted: Font: Italic
		<i>Salmonella</i> spp.	1	guichai leaves	Formatted
		<i>Salmonella</i> spp.	2	horopa leaves	Formatted
		<i>Escherichia coli</i> , <i>Salmonella</i> bBraenderup	2	horseradish	Formatted
		<i>Salmonella</i> spp.	1	kayang leaves	Formatted
		<i>Salmonella</i> spp.	1	lalo leaves	Formatted
		<i>Salmonella</i> group C, <i>Salmonella</i> spp., <i>Salmonella</i> Wandsworth	3	lemon grass	Formatted
		<i>Salmonella</i> Lexington	1	lime leaves	Formatted
		<i>Salmonella</i> spp.	1	magaso sadao	Formatted
		<i>Salmonella</i> spp.	1	malabar nightshade	Formatted
		<i>Salmonella enterica</i>	1	mimosa	Formatted: Font: Italic
		<i>Escherichia coli</i> , <i>Salmonella</i> Javiana, <i>Salmonella</i> spp., <i>Salmonella</i> Stanley, <i>Salmonella</i> Zanzibar, <i>Salmonella</i> Weltevreden, <i>Salmonella</i> eChester, <i>Salmonella enterica</i> , <i>Salmonella</i> Heidelberg, <i>Salmonella</i> Hvittingfoss, <i>Salmonella</i> Javiana, <i>Salmonella</i> Ppoona, <i>Salmonella</i> spp., <i>Salmonella</i> Stanley, <i>Salmonella</i> Thompson, <i>Salmonella</i> Zanzibar	25	mint	Formatted
		<i>Salmonella</i> GIVE, <i>Salmonella</i> Kkedougou, <i>Salmonella</i> Typhimurium	3	mint and basil	Formatted
		<i>Salmonella enterica</i>	3	mint leaves	Formatted: Font: Italic
		<i>Salmonella</i> Rissen			Formatted: Font: Italic
		<i>Salmonella</i> Virchow			Formatted: Font: Italic
		<i>Salmonella</i> Bleadon, <i>Salmonella</i> Hvittingfoss, <i>Salmonella</i> Kentucky, <i>Salmonella</i> Mount Pleasant, <i>Salmonella</i> Ramat-gan, <i>Salmonella</i> Rissen, <i>Salmonella</i> spp.	8	morning glory	Formatted: Font: Italic
		<i>Salmonella</i> Hvittingfoss, <i>Salmonella</i> Javiana, <i>Salmonella</i> Stanley, <i>Salmonella</i> Weltevreden, <i>Salmonella</i> Zanzibar	5	other herbs	Formatted
		<i>Salmonella</i> spp.	1	pakdum	Formatted
		<i>Salmonella enterica</i>	2	pandang leaves	Formatted: Font: Italic
		<i>Salmonella</i> augustinborg, <i>Salmonella</i> Newport, <i>Salmonella</i> spp.	3	parsley	Formatted: Font: Italic
		<i>Salmonella</i> Virchow	1	perilla	Formatted
		<i>Salmonella</i> spp.	3	praew leaves	Formatted

Country of origin	Notifications (n)	Pathogens	No of pathogens	Herbs and spices	
		<i>Salmonella enterica</i>	1	rice paddy herb	Formatted: Font: Italic
		<i>Salmonella</i> Saint Paul, <i>Salmonella</i> spp, <i>Salmonella</i> Stanley	3	Spinach	Formatted
		<i>Salmonella</i> Hvittingfoss	1	wildbetal le...	Formatted: Font: Italic
		<i>Salmonella</i> <del>a</del> Augustenborg, <i>Salmonella</i> Paratyphi B (variant Java monophasic variant 4,5,12)	2	Basil and ri paddy herb	Formatted
		<i>Salmonella</i> <del>a</del> Augustenborg, <i>Salmonella</i> Lexington	2	basiland lin leaves	Formatted
		<i>Salmonella</i> spp	2	pennywort	Formatted: Font: Italic
		<i>Escherichia coli</i> , <i>Salmonella</i> Javiana, <i>Salmonella</i> spp, <i>Salmonella</i> Virchow	8	basil	Formatted
		<i>Salmonella enterica</i>	1	celery	Formatted: Font: Italic
		<i>Salmonella enterica</i> ser. Virchow	1	chilli	Formatted
		<i>Salmonella</i> spp	1	coriander	Formatted
Vietnam	49	<i>Salmonella</i> spp	1	eryngo	Formatted
		<i>Salmonella</i> spp	2	green herb	Formatted
		<i>Salmonella enterica</i> , <i>Salmonella</i> Lexington, <i>Salmonella</i> Orion, <i>Salmonella</i> spp, <i>Salmonella</i> Weltevreden, <i>Vibrio cholerae</i>	13	ground pepper	Formatted
		<i>Salmonella</i> Stanley	1	hing choi	Formatted
		<i>Salmonella</i> spp	1	houuttunya leaves	Formatted
		<i>Salmonella</i> spp	1	houuttunya leaves	Formatted
		<i>Salmonella</i> spp	2	kinh gioi leaves	Formatted
		<i>Salmonella</i> spp	4	la lot leaves	Formatted
		<i>Salmonella enterica</i> , <i>Salmonella</i> spp	3	mint	Formatted
		<i>Salmonella</i> spp	1	mong toi leaves	Formatted
		<i>Salmonella</i> spp	1	morning gl	Formatted
		<i>Salmonella</i> Abony, <i>Salmonella</i> Dublin, <i>Salmonella</i> Orientalis, <i>Salmonella</i> Virchow, <i>Salmonella</i> Weltevreden, <i>Salmonella</i> spp	7	other herbs	Formatted
		<i>Salmonella</i> spp	3	parsley	Formatted
		<i>Salmonella</i> spp	1	perilla	Formatted
		<i>Salmonella</i> spp	1	Piper sarmentosum	Formatted

Country of origin	Notifications (n)	Pathogens	No of pathogens	Herbs and spices
		<i>Salmonella</i> spp.	1	rau ram leaves
		<i>Salmonella</i> Javiana	1	rice paddy herb

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Bold

- Abatcha, M. G., Effarizah, M. E., & Rusul, G. (2018). Prevalence, antimicrobial resistance, resistance genes and class 1 integrons of *Salmonella* serovars in leafy vegetables, chicken carcasses and related processing environments in Malaysian fresh food markets. *Food Control*, *91*, 170–180.
- Ananchaipattana, C., Hosotani, Y., Kawasaki, S., Pongswat, S., Latiful, B. M., Isobe, S., & Inatsu, Y. (2012). Bacterial contamination of soybean curd (tofu) sold in Thailand. *Food Science and Technology Research*, *19*(6), 843–848.
- Bag, P. K., Nandi, S., Bhadra, R. K., Ramamurthy, T., Bhattacharya, S., Nishibuchi, M., Hamabata, T., Yamasaki, S., Takeda, Y., & Nair, G. B. (1999). Clonal diversity among recently emerged strains of *Vibrio parahaemolyticus* O3: K6 associated with pandemic spread. *Journal of clinical microbiology*, *37*(7), 2354–2357.
- Banerjee, M., & Sarkar, P. K. (2003). Microbiological quality of some retail spices in India. *Food Research International*, *36*(5), 469–474.
- Bangtrakulnonth, A., Pornreongwong, S., Pulsrikarn, C., Sawanpanyalert, P., Hendriksen, R. S., Wong, D. M. L. F., & Aarestrup, F. M. (2004). *Salmonella* serovars from humans and other sources in Thailand, 1993–2002. *Emerging infectious diseases*, *10*(1), 131.
- Bernard, H., Faber, M., Wilking, H., Haller, S., Höhle, M., Schielke, A., Ducomble, T., Sifczyk, C., Merbecks, S., & Fricke, G. (2014). Large multistate outbreak of norovirus gastroenteritis associated with frozen strawberries, Germany, 2012. *Eurosurveillance*, *19*(8), 20719.
- Bisht, A., Kamble, M. P., Choudhary, P., Chaturvedi, K., Kohli, G., Juneja, V. K., Sehgal, S., & Taneja, N. K. (2021). A surveillance of food borne disease outbreaks in India: 2009–2018. *Food Control*, *121*, 107630.
- Borchers, A., Teuber, S. S., Keen, C. L., & Gershwin, M. E. (2010). Food safety. *Clinical reviews in allergy & immunology*, *39*(2), 95–141.
- Briet, A., Helsen, N., Delannoy, S., Debuiche, S., Brisabois, A., Midelet, G., & Granier, S. (2018). NDM-1-producing *Vibrio parahaemolyticus* isolated from imported seafood. *J Antimicrob Chemother* doi, 10.
- Carstens, C. K., Salazar, J. K., & Darkoh, C. (2019). Multistate outbreaks of foodborne illness in the United States associated with fresh produce from 2010 to 2017. *Frontiers in microbiology*, *10*, 2667.
- Chaoniruthisai, P., Punnakitikashem, P., & Rajchamaha, K. (2018). Challenges and difficulties in the implementation of a food safety management system in Thailand: A survey of BRC certified food productions. *Food Control*, *93*, 274–282.
- Chen, A., Giese, M., & Chen, D. (2020). Flood impact on Mainland Southeast Asia between 1985 and 2018—The role of tropical cyclones. *Journal of Flood Risk Management*, *13*(2), e12598.
- Chen, X., Zhu, Q., Yu, F., Zhang, W., Wang, R., Ye, X., Jin, L., Liu, Y., Li, S., & Chen, Y. (2018). Serology, virulence and molecular characteristics of *Vibrio parahaemolyticus* isolated from seafood in Zhejiang province. *PloS one*, *13*(10), e0204892.

1408 Cook, N., Williams, L., & D'Agostino, M. (2019). Prevalence of Norovirus in produce sold at retail in the  
 1409 United Kingdom. *Food microbiology*, 79, 85–89.

1410 DEFRA. (2020). Food Statistics in your pocket: Global and UK supply. Department for Environment  
 1411 Food & Rural Affairs.

1412 Dolan, C., & Humphrey, J. (2000). Governance and trade in fresh vegetables: the impact of UK  
 1413 supermarkets on the African horticulture industry. *Journal of development studies*, 37(2), 147–  
 1414 176.

1415 Dos Santos, S. F., Cardoso, R. d. C. V., Borges, Í. M. P., e Almeida, A. C., Andrade, E. S., Ferreira, I. O., &  
 1416 do Carmo Ramos, L. (2020). Post-harvest losses of fruits and vegetables in supply centers in  
 1417 Salvador, Brazil: Analysis of determinants, volumes and reduction strategies. *Waste*  
 1418 *Management*, 101, 161–170.

1419 ECDC/EFSA. (2021). Multi-country outbreak of *Salmonella* Enteritidis sequence type (ST)11 infections  
 1420 linked to poultry products in the EU/EEA and the United Kingdom — 25 February 2021.  
 1421 Stockholm: ECDC/EFSA; 2021. European Centre for Disease Prevention and Control (ECDC) and  
 1422 European Food Safety Authority (EFSA).

1423 EFSA/ECDC. (2019). Scientific report on the European Union one health 2018. Zoonoses report.  
 1424 European Centre for Disease Prevention and Control (ECDC) and European Food Safety  
 1425 Authority (EFSA). *EFSA J.*, 17, 5926.

1426 EFSA/ECDC. (2021). The European Union One Health 2019 Zoonoses Report. European Food Safety  
 1427 Authority (EFSA) and European Centre for Disease Prevention and Control (ECDC). *EFSA*  
 1428 *Journal* 19, 6406.

1429 EFSA/ECDC. (2018). Multi-country outbreak of *Salmonella* Agona infections possibly linked to ready-  
 1430 to-eat food. European Food Safety Authority (EFSA) and European Centre for Disease  
 1431 Prevention Control (ECDC). In: Wiley Online Library.

1432 Emberland, K., Ethelberg, S., Kuusi, M., Vold, L., Jensvoll, L., Lindstedt, B., Nygård, K., Kjelsø, C.,  
 1433 Torpdahl, M., & Sørensen, G. (2007). Outbreak of *Salmonella* Weltevreden infections in  
 1434 Norway, Denmark and Finland associated with alfalfa sprouts, July–October 2007. *Weekly*  
 1435 *reports (1997–2007)*, 12(48), 3321.

1436 European Commission. (2014). Commission implementing decision of 13 February 2014 suspending  
 1437 temporarily imports from Bangladesh of foodstuffs containing or consisting of betel leaves  
 1438 ('Piper betle'). Available from: [https://eur-lex.europa.eu/legal-](https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32014D0088&from=DA)  
 1439 [content/EN/TXT/PDF/?uri=CELEX:32014D0088&from=DA](https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32014D0088&from=DA) Accessed 25.02.2021.

1440 European Commission. (2015). Commission implementing regulation (EU) 2015/943 of 18 June 2015.  
 1441 *Official Journal of the European Union*(154), 8–9.

1442 European Commission (2016). RASFF — The Rapid Alert System for Food and Feed — 2015 annual  
 1443 report. Available from: [https://op.europa.eu/en/publication-detail/-/publication/98e04cfc-](https://op.europa.eu/en/publication-detail/-/publication/98e04cfc-aa54-11e6-aab7-01aa75ed71a1/language-en/format-PDF/source-174743070)  
 1444 [aa54-11e6-aab7-01aa75ed71a1/language-en/format-PDF/source-174743070](https://op.europa.eu/en/publication-detail/-/publication/98e04cfc-aa54-11e6-aab7-01aa75ed71a1/language-en/format-PDF/source-174743070) Accessed  
 1445 19.02.2021.

1446 European Commission. (2020a). Commission implementing regulation (EU) 2020/625 of 6 May 2020.  
 1447 *Official Journal of the European Union*(63), 13.

1448 European Commission. (2020b). Monitoring EU Agri-Food Trade: Developments in 2019. Available  
 1449 from: [https://ec.europa.eu/info/news/2019-record-year-eu-agri-food-trade-2020-mar-](https://ec.europa.eu/info/news/2019-record-year-eu-agri-food-trade-2020-mar-27-en#moreinfo)  
 1450 [27-en#moreinfo](https://ec.europa.eu/info/news/2019-record-year-eu-agri-food-trade-2020-mar-27-en#moreinfo) Accessed 12.02.2021.

1451 European Food Safety Authority. (2021). The European Union One Health 2019 Zoonoses Report. *EFSA*  
 1452 *Journal*, 19(2).

1453 Fakruddin, M., Sultana, R., Hossain, M. N., Rahaman, M. M., Islam, M. K., & Ahmed, M. M. (2017).  
 1454 Occurrence of ingress of *Salmonella* spp. in Betel leaf (*Piper betle* L.). *International Journal*  
 1455 *of Food Contamination*, 4(1), 1–10.

1456 Ferrari, R. G., Rosario, D. K., Cunha Neto, A., Mano, S. B., Figueiredo, E. E., & Conte Junior, C. A. (2019).  
 1457 Worldwide epidemiology of *Salmonella* serovars in animal-based foods: a meta-analysis.  
 1458 *Applied and environmental microbiology*, 85(14).

- Food and Agricultural Organisation. (2018). Addressing the food safety challenges of the Asia-Pacific region. 34th Session of the FAO Regional Conference for Asia and the Pacific. Available from: <http://www.fao.org/3/MV819en/mv819en.pdf> Accessed 24.02.2021.
- Food and Agriculture Organization. (2019). Foodborne illnesses in Asia-Pacific 'needlessly' sicken 275 million annually and threaten trade. Retrieved from <http://www.fao.org/asiapacific/news/detail-events/en/c/1197008/#:~:text=In%20the%20Asia%2DPacific%20region,of%20zero%20Hunger%20by%202030.>
- Fu, H., Yu, P., Liang, W., Kan, B., Peng, X., & Chen, L. (2020). Virulence, Resistance, and Genomic Fingerprint Traits of *Vibrio cholerae* Isolated from 12 Species of Aquatic Products in Shanghai, China. *Microbial Drug Resistance*, 26(12), 1526–1539.
- Ghosh, D. (2014). Food safety regulations in Australia and New Zealand food standards. *Journal of the Science of Food and Agriculture*, 94(10), 1970–1973.
- Jaffee, S., Henson, S., Unnevehr, L., Grace, D., & Cassou, E. (2018). *The safe food imperative: Accelerating progress in low and middle income countries*. World Bank Publications.
- Kaakoush, N. O., Castaño-Rodríguez, N., Mitchell, H. M., & Man, S. M. (2015). Global epidemiology of *Campylobacter* infection. *Clinical microbiology reviews*, 28(3), 687–720.
- Kantama, L., & Jayanetra, P. (1996). Salmonella enteritidis outbreak in Thailand: study by random amplified polymorphic DNA (RAPD) analysis. *The Southeast Asian journal of tropical medicine and public health*, 27(1), 119–125.
- Keisam, S., Tuikhar, N., Ahmed, G., & Jeyaram, K. (2019). Toxigenic and pathogenic potential of enteric bacterial pathogens prevalent in the traditional fermented foods marketed in the Northeast region of India. *International journal of food microbiology*, 296, 21–30.
- Kim, K., Kim, S., & Park, C. Y. (2020). Food Security in Asia and the Pacific amid the COVID-19 Pandemic. 1–15.
- Kowalska, A., & Manning, L. (2020). Using the rapid alert system for food and feed: potential benefits and problems on data interpretation. *Critical reviews in food science and nutrition*, 1–14.
- Kumar, R., & Lalitha, K. V. (2013). Prevalence and molecular characterization of *Vibrio cholerae* O1, non-O1 and non-O139 in tropical seafood in Cochin, India. *Foodborne pathogens and disease*, 10(3), 278–283.
- Kumari, S., & Sarkar, P. K. (2014). Prevalence and characterization of *Bacillus cereus* group from various marketed dairy products in India. *Dairy science & technology*, 94(5), 483–497.
- Lee, Y., Choi, Y., Lee, S., Lee, H., Kim, S., Ha, J., Lee, J., Oh, H., Kim, Y., & Yoon, Y. (2019). Occurrence of pathogenic *Vibrio parahaemolyticus* in seafood distribution channels and their antibiotic resistance profiles in S. Korea. *Letters in applied microbiology*, 68(2), 128–133.
- Lemoine, T., Germanetto, P., & Giraud, P. (1999). Toxi-infection alimentaire à *V. parahaemolyticus*. *Bull Epidemiol Hebdo*, 10, 37–38.
- Machado-Moreira, B., Richards, K., Brennan, F., Abram, F., & Burgess, C. M. (2019). Microbial contamination of fresh produce: what, where, and how? *Comprehensive reviews in food science and food safety*, 18(6), 1727–1750.
- Makendi, C., Page, A. J., Wren, B. W., Le Thi Phuong, T., Clare, S., Hale, C., Goulding, D., Klemm, E. J., Pickard, D., & Okoro, C. (2016). A phylogenetic and phenotypic analysis of *Salmonella enterica* serovar Weltevreden, an emerging agent of diarrheal disease in tropical regions. *PLoS neglected tropical diseases*, 10(2), e0004446.
- Mangla, S. K., Bhattacharya, A., Yadav, A. K., Sharma, Y. K., Ishizaka, A., Luthra, S., & Chakraborty, R. (2021). A framework to assess the challenges to food safety initiatives in an emerging economy. *Journal of Cleaner Production*, 284, 124709.
- Martinez-Urtaza, J., Simental, L., Velasco, D., DePaola, A., Ishibashi, M., Nakaguchi, Y., Nishibuchi, M., Carrera-Flores, D., Rey-Alvarez, C., & Pousa, A. (2005). Pandemic *vibrio parahaemolyticus* O3:K6, Europe. *Emerging infectious diseases*, 11(8), 1319.

McLauchlin, J., Aird, H., Andrews, N., Chattaway, M., De Pinna, E., Elviss, N., Jørgensen, F., Larkin, L., & Willis, C. (2019). Public health risks associated with Salmonella contamination of imported edible betel leaves: analysis of results from England, 2011–2017. *International journal of food microbiology*, 298, 1–10.

McLauchlin, J., Aird, H., Charlett, A., Chattaway, M., Elviss, N., Hartman, H., Jenkins, C., Jørgensen, F., Larkin, L., & Sadler-Reeves, L. (2018). Imported edible leaves collected at retail sale in England during 2017 with an emphasis on betel and curry leaves: microbiological quality with respect to Salmonella, Shiga toxin-producing E. coli (STEC) and levels of Escherichia coli. *Journal of applied microbiology*, 125(4), 1175–1185.

Mensah, L. D., & Julien, D. (2011). Implementation of food safety management systems in the UK. *Food Control*, 22(8), 1216–1225.

Nakaguchi, Y. (2013). Contamination with *Vibrio parahaemolyticus* and its virulent strains in seafood marketed in Thailand, Vietnam, Malaysia, and Indonesia from 2008 to 2011. *Tropical Medicine and Health*, 2011–2006.

Nasheri, N., Vester, A., & Petronella, N. (2019). Foodborne viral outbreaks associated with frozen produce. *Epidemiology & Infection*, 147.

Nath, T. K., & Inoue, M. (2009). Sustainability attributes of a small-scale betel leaf agroforestry system: a case study in north-eastern hill forests of Bangladesh. *Small-scale Forestry*, 8(3), 289–304.

Nguyen, T. K., Bui, H. T., Truong, T. A., Lam, D. N., Ikeuchi, S., Ly, L. K. T., Hara-Kudo, Y., Taniguchi, T., & Hayashidani, H. (2021). Retail fresh vegetables as a potential source of Salmonella infection in the Mekong Delta, Vietnam. *International journal of food microbiology*, 341, 109049.

Office of National Statistics. (2011). UK Census 2011. Available online: <https://www.ons.gov.uk/census/2011census>

Ortega, D. L. a. T., D. L. (2017). Demand for food safety in emerging and developing countries: A research agenda for Asia and Sub-Saharan Africa. *Journal of Agribusiness in Developing and Emerging Economies*(7), 21–34.

Park, S., Szonyi, B., Gautam, R., Nightingale, K., Anciso, J., & Ivanek, R. (2012). Risk factors for microbial contamination in fruits and vegetables at the preharvest level: a systematic review. *Journal of food protection*, 75(11), 2055–2081.

Pham, H. V., & Dinh, T. L. (2020). The Vietnam's food control system: Achievements and remaining issues. *Food Control*, 108, 106862.

Reddy, A. A., Ricart, S., & Cadman, T. (2020). Driving factors of food safety standards in India: learning from street food vendors' behaviour and attitude. *Food Security*, 1–17.

Rezaei, M., & Liu, B. (2017). Food loss and waste in the food supply chain. *International Nut and Dried Fruit Council: Reus, Spain*, 26–27.

Rivas, L., Strydom, H., Paine, S., Wang, J., & Wright, J. (2021). Versiniosis in New Zealand. *Pathogens*, 10(2), 191.

Singh, B. R., Singh, M., Singh, P., Babu, N., Chandra, M., & Agarwal, R. K. (2006). Prevalence of multidrug-resistant Salmonella on ready-to-eat betel leaves (Paan) and in water used for soaking betel leaves in North Indian cities. *Journal of food protection*, 69(2), 288–292.

Sodagari, H. R., Wang, P., Robertson, I., Habib, I., & Sahibzada, S. (2020). Non-Typhoidal Salmonella at the Human-Food-of Animal-Origin Interface in Australia. *Animals*, 10(7), 1192.

Somorin, Y. M., Odeyemi, O. A., & Ateba, C. N. (2021). Salmonella is the most common foodborne pathogen in African food exports to the European Union: Analysis of the Rapid Alert System for Food and Feed (1999–2019). *Food Control*, 123, 107849.

Sugiri, Y. D., Goelz, G., Meeyam, T., Baumann, M. P., Kleer, J., Chaisowwong, W., & Alter, T. (2014). Prevalence and antimicrobial susceptibility of *Listeria monocytogenes* on chicken carcasses in Bandung, Indonesia. *Journal of food protection*, 77(8), 1407–1410.

Tähhkäpää, S., Maijala, R., Korkeala, H., & Nevas, M. (2015). Patterns of food frauds and adulterations reported in the EU rapid alert system for food and feed and in Finland. *Food Control*, 47, 175–184.

- Ullah, A. S., Tani, M., Tsuchiya, J., Rahman, A. M., & Rahman, Z. M. (2020). Impact of Betel Leaf Cultivation on the Protected Forest Area of Teknaf Peninsula, Bangladesh. *Small-scale Forestry*, 19(3), 335-355.
- Vilain, S., Luo, Y., Hildreth, M. B., & Brözel, V. S. (2006). Analysis of the life cycle of the soil saprophyte *Bacillus cereus* in liquid soil extract and in soil. *Applied and environmental microbiology*, 72(7), 4970-4977.
- Vital, P. G., Dimasuay, K. G. B., Widmer, K. W., & Rivera, W. L. (2014). Microbiological quality of fresh produce from open air markets and supermarkets in the Philippines. *The Scientific World Journal*, 2014.
- Wadamori, Y., Gooneratne, R., & Hussain, M. A. (2017). Outbreaks and factors influencing microbiological contamination of fresh produce. *Journal of the Science of Food and Agriculture*, 97(5), 1396-1403.
- Wang, S., Weller, D., Falardeau, J., Strawn, L. K., Mardones, F. O., Adell, A. D., & Switt, A. I. M. (2016). Food safety trends: from globalization of whole genome sequencing to application of new tools to prevent foodborne diseases. *Trends in Food Science & Technology*, 57, 188-198.
- Wong, H.-C., Chen, M.-C., Liu, S.-H., & Liu, D.-P. (1999). Incidence of highly genetically diversified *Vibrio parahaemolyticus* in seafood imported from Asian countries. *International journal of food microbiology*, 52(3), 181-188.
- World Health Organization. (2015). *WHO estimates of the global burden of foodborne diseases: foodborne disease burden epidemiology reference group 2007-2015*. World Health Organization.
- World Health Organization. (2019). *Shiga Toxin-producing Escherichia Coli (STEC) and Food: Attribution Characterization and Monitoring* (Vol. 19). World Health Organization.
- World Health Organization. (2020a). Food Safety. Retrieved from <https://www.who.int/news-room/fact-sheets/detail/food-safety>.
- World Health Organization. (2020b). Framework for action on food safety in the WHO South-East Asia Region. New Delhi: World Health Organization, Regional Office for South-East Asia.
- World Health Organization. (2020c). Salmonella – non-typhoidal. Available from: [https://www.who.int/news-room/fact-sheets/detail/salmonella-\(non-typhoidal\)](https://www.who.int/news-room/fact-sheets/detail/salmonella-(non-typhoidal)) Accessed 25.02.2021.
- Worldometer. (2021). Current World Population. Retrieved from <https://www.worldometers.info/world-population/#ref-1>.
- Xu, X., Cheng, J., Wu, Q., Zhang, J., & Xie, T. (2016). Prevalence, characterization, and antibiotic susceptibility of *Vibrio parahaemolyticus* isolated from retail aquatic products in North China. *BMC microbiology*, 16(1), 1-9.
- Yu, P., Zaleski, A., Li, Q., He, Y., Mapili, K., Pruden, A., Alvarez, P. J., & Stadler, L. B. (2018). Elevated levels of pathogenic indicator bacteria and antibiotic resistance genes after Hurricane Harvey's flooding in Houston. *Environmental science & technology letters*, 5(8), 481-486.

**Declaration of competing interest**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.



**CRedit authorship contribution statement**

**Ayokunle C. Dada:** Conceptualization, Methodology, Formal analysis, Writing - original draft, Review and Editing, Writing - review & editing. **Yinka M. Somorin:** Writing - original draft, Review and Editing, Writing - review & editing. **Collins N. Ateba:** Writing - original draft, Review and Editing, Writing - review & editing. **Helen Onyeaka:** Review and Editing, Writing - review & editing. **Amarachukwu Anyogu:** Review and Editing, Writing - review & editing. **Nor Azman Kasan:** Writing - original draft, Review and Editing, Writing - review & editing. **Olumide A. Odeyemi:** Conceptualization, Methodology, Formal analysis, Writing - original draft, Review and Editing, Writing - review & editing.