



## **UWL REPOSITORY**

**repository.uwl.ac.uk**

Microorganisms and food safety risks associated with indigenous fermented foods from Africa

Anyogu, Amarachukwu ORCID logoORCID: <https://orcid.org/0000-0001-9652-7728>, Olukorede, Ayomide, Anumudu, Christian, Onyeaka, Helen, Areo, Esther, Adewale, Obadina, Odimba, Joyce N. and Nwaiwu, Ogueri (2021) Microorganisms and food safety risks associated with indigenous fermented foods from Africa. Food Control, 129. p. 108227. ISSN 0956-7135

<http://dx.doi.org/10.1016/j.foodcont.2021.108227>

This is the Accepted Version of the final output.

UWL repository link: <https://repository.uwl.ac.uk/id/eprint/8244/>

**Alternative formats:** If you require this document in an alternative format, please contact: [open.research@uwl.ac.uk](mailto:open.research@uwl.ac.uk)

**Copyright:** Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy:** If you believe that this document breaches copyright, please contact us at [open.research@uwl.ac.uk](mailto:open.research@uwl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

**Microorganisms and food safety risks associated with indigenous fermented foods from  
Africa**

Amarachukwu Anyogu<sup>a\*</sup>, Ayomide Olukorede<sup>b</sup>, Christian Anumudu<sup>c</sup>, Helen Onyeaka<sup>c</sup>, Esther Areo<sup>d</sup>,  
Obadina Adewale<sup>d</sup>, Joyce N. Odimba<sup>e</sup>, Ogueri Nwaiwu<sup>f</sup>

**\*Corresponding author:** Amarachukwu Anyogu, PhD, Applied Biotechnology Research Group,  
College of Liberal Arts and Sciences, University of Westminster, 115 New Cavendish Street, London  
W1W 6UW, United Kingdom.

**Email:** amara.anyogu@gmail.com

## Abstract

Indigenous fermented foods (IFFs) have a long history in Africa and are embedded in cultural norms and practices. Typically, these foods are produced at small or household scale using indigenous processing technologies. In addition, limited knowledge of good manufacturing and handling practices can lead to production under unhygienic conditions. This results in variations in the quality and safety attributes of IFFs, as spoilage and pathogenic bacteria can be introduced at any stage of the value chain. These foods have an important role in the African diet and can contribute to food security by increasing the availability of cheap, nutritious food and supporting livelihoods. However, the presence of foodborne pathogens and antibiotic-resistant bacteria in IFFs may constitute a health risk to consumers. Therefore, this review presents an overview of the microorganisms associated with IFFs from Africa, focusing on microbial food safety hazards. African indigenous fermented foods offer a vast genetic potential of undiscovered strains that possess valuable technical characteristics. However, IFFs may also serve as vehicles of pathogenic and antibiotic-resistant bacteria and genetic determinants. Significant research and data gaps exist regarding the microbiological safety of these food products, which warrant urgent attention. We propose practical solutions for improving the safety of African IFFs requiring action and collaboration from all stakeholders, including researchers, producers, governmental regulatory bodies, and consumers.

**Keywords:** Africa; antibiotic resistance; fermented foods; food safety; lactic acid bacteria; pathogens

## 1. Introduction

Indigenous fermented foods (IFFs) have a long history in Africa and are embedded in the cultural norms and practices. They make a valuable contribution to the continent's effort in achieving its sustainable development goals focused on food security, poverty alleviation and gender equality (Franz et al., 2014). Fermentation adds value, improving the organoleptic characteristics, variety, nutritional quality, digestibility, and safety of foods (FAO, 2010; Ganzle, 2020). As a low-cost technology, indigenous fermentations improves access to local, inexpensive, nutritious food. In addition, IFFs contribute to the livelihoods of many, especially women, through income generation via small-scale enterprise (Misihairabgwi & Cheikyoussef, 2017; Rolle & Satin, 2002). Fermentation is a processing technique where desirable changes occur in a food product due to the metabolic activities of microorganisms (Caplice and Fitzgerald, 1999). There has been increasing interest from researchers to understand the diversity and technological properties of the microbial community in IFFs from Africa. This knowledge is required for producing local, well-characterised starter cultures to improve the quality and safety of IFF products (Akabanda et al., 2013; Anyogu et al., 2014; Bayili et al., 2019). Despite these efforts, IFF production in Africa largely remains a cottage-level technology, mainly reliant on spontaneous fermentation processes or backslopping, which may introduce spoilage or pathogenic organisms (Benkerroum and Tamime, 2004). Bacteria of public health interest have been isolated from IFFs, and some investigators have noted concerns about the safety of these foods (Ahaotu et al., 2013; Omemu et al., 2014; Samet-Bali et al., 2016; Walsh et al., 2017). Data from the World Health Organisation (WHO) estimate that diarrhoeal diseases account for 70% of mortalities associated with foodborne disease in the African region (WHO, 2015). Also, antibiotic resistance (AR) has emerged as one of the utmost global public health concerns. Fermented food products have diverse microbial ecosystems, yet their ability to serve as vehicles for transferring AR microorganisms and genes remains unclear.

Reviews discussing IFFs from Africa have mainly focused on the microbial diversity of fermenting and technologically important bacteria (Achi and Ukwuru, 2013; Jans et al., 2017; Parkouda et al., 2009; Tamang et al., 2016). Recently, Nwaiwu et al. (2020) reviewed spoilage and pathogenic microbiota associated with indigenous fermented beverages but only concentrated on one country. Paudyal et al. (2017) analysed the prevalence of foodborne pathogens in foods from selected African countries; however, their focus was on raw and ready-to-eat foods and not IFFs specifically. A comprehensive overview of the microbiological safety of IFFs from Africa is lacking. The purpose of this review is to summarise current knowledge on the microbiology of IFFs with a focus on pathogenic and AR bacteria. It also discusses approaches to improve the safety of these foods and highlights data gaps that could be explored in further research.

## **2. Indigenous African fermented food products**

Fermentation has long been used as a preservation technology for extending the shelf life of various substrates. The characteristic flavour, texture and colour of IFFs due to the metabolic activities of fermenting microorganisms has guaranteed their widespread acceptability by consumers, thereby establishing a major role for IFFs in the African diet (Mokoena et al., 2016). Olasupo et al. (2010) classified IFFs from Africa into five major groups based on their raw materials. These include (1) fermented non-alcoholic cereals, (2) alcoholic beverages, (3) fermented animal proteins, (4) starchy root crops, and (5) fermented vegetable proteins.

Cereal grains comprising maize, sorghum, millet, and wheat are important staple crops in Africa, accounting for as much as 50% of the total daily calorific consumption (OECD/FAO, 2016). These cereals are common starting materials for lactic acid-fermented beverages and porridges known by different names such as *ogi* in Nigeria (Oguntoyinbo et al., 2011), *togwa* in Tanzania (Mugula et al., 2003), *koko* in Ghana (Lei and Jakobsen, 2004) and *hussuwa* in Sudan (Yousif et al., 2010).

Fermented non-alcoholic cereal-based products have an important role in the diet as complementary foods for infants or breakfast meals (Byakiya et al., 2019; Soro-Yao et al., 2014). The extensive use of cereal-based complementary foods makes them an attractive target in efforts

towards combating infant malnutrition in Africa. For example, *koko*, made from fermented corn dough, is the most commonly consumed complementary food in Ghana. However, it has been noted to be inadequate in meeting dietary protein and micronutrient needs (Suri et al., 2014). To improve the nutritional profile of *koko*, a legume-based supplement was developed recently, with preliminary studies suggesting good acceptability by consumers (Tano-Debrah et al., 2019).

Fermented alcoholic beverages are consumed across the continent. The majority of these are produced from cereals, e.g., sorghum, millet, and maize. These include *dolo*, *burukutu*, or *pito* in West Africa (Sawodogo-Lingani et al., 2007; Onyenekwe et al., 2015 ), *borde* in Ethiopia (Abegaz, 2007), and *sesotho* in South Africa (Cason et al., 2020). Fruit fermentation for alcohol production is uncommon. However, plantain and banana can be fermented to produce *agadagidi* (Sanni and Lonner, 1993). Palm wine is a popular alcoholic drink in many West African countries. Palm wine is obtained from the fermentation of sap obtained from palm tree species such as *Elaeis guineensis*, *Raphia hookeri*, *Borassus aethiopum*, and *Borassus akeassii*. The production of ethanol, lactic acid and acetic acids by yeasts, lactic acid bacteria (LAB) and acetic acid bacteria (AAB) are the most significant activities contributing to the distinctive organoleptic characteristics and stability of the product (Amoa-Awua et al., 2007; Karamoko et al., 2012; Nwaiwu & Itumoh, 2017). The gradual accumulation of ethanol at the early stages of fermentation correlates with the increase in the AAB population, as AAB can use ethanol as a carbon source (Amoa-Awua et al., 2007). Organic acids produced by LAB & AAB contribute to the sour taste, aroma and colour, although AAB are often considered as spoilage organisms. The low pH and alcohol content control the growth of undesirable bacteria, including *Enterobacteriaceae* (Djeni et al., 2020; Ouoba et al., 2012).

Cassava (*Manihot esculenta*, Crantz) ranks third after rice and maize as a source of calories in tropical countries and is the most consumed starchy root crop in Africa. Total annual cassava consumption has tripled from 20 million tonnes in 1970 to just over 60 million tonnes in 2013 (FAOSTAT, 2020a; Szyniszweska et al., 2020). Fermentation practices involve either submerged fermentation for the production of *lafun* and *chikwangue* (Miambi et al, 2003; Padonou et al., 2009)

or solid-state to produce *attieke* and *garri* (Ahaotu et al., 2017; Djeni et al., 2015). Fermentation is essential for preventing the rapid post-harvest deterioration of cassava tubers. It is also an important processing technique for decreasing the amount of cyanogenic glucosides (CGs), which occur naturally in cassava. These CGs can be hydrolysed to produce hydrogen cyanide when the plant tissue is damaged, but it should be noted that the concentration of CGs can be cultivar specific (Abiodun et al., 2020). Cyanide can inhibit cellular respiration by binding to cytochrome oxidase in the electron transport chain, and high exposure to cyanide can lead to severe illness and, in some cases, death (Akintowa et al., 1994; Kimaryo et al., 2000). Consequently, the WHO has recommended a safe limit of 10 mg cyanide/kg of cassava (FAO/WHO, 1995). Although the breakdown of cyanide has been attributed to the activity of endogenous enzymes, Obilie et al. (2004) observed that the highest loss of cyanogenic glucosides, from 72.4 mg cyanide/kg to below detectable limits, occurred during fermentation compared to other processing stages. These results are corroborated by Kivunde et al. (2000) during controlled fermentations using LAB as a starter where the total cyanide content reduced from 176.3 mg/kg in raw cassava to 8.2 mg/kg in *kivunde*. Their observation of higher cyanide content in spontaneously fermented and backslopped samples was attributed to  $\beta$ -glycosidase activity of the starter strains, suggesting an additional advantage of controlled fermentations in guaranteeing the safety of fermented cassava products. Another demerit of cassava is its low nutritional content, specifically protein, vitamins and minerals and studies on fermented cassava products biofortified with protein (Ahaotu et al., 2017) and Vitamin A (Abiodun et al., 2020) have recently been reported.

In Africa, milk has a long historical connection with pastoral communities, e.g., the Berbers, Fulani and Maasai ethnic groups in North, West and East Africa. Milk is a significant source of nutrients, and total milk production in Africa was 49 million tonnes in 2019 (FAOStat, 2020). Fermentation is an essential processing technology for extending the shelf life of milk, a highly perishable product. African fermented dairy products comprise of naturally fermented milks, e.g., *gariss* and *mabisi*; yoghurt-based products, e.g., *kindirmo* and *amasi*, and cheeses, e.g., *wara*, *jben* and *klilia*

(Abdelgadir et al., 2008; Mourad & Bettache, 2015; Moonga et al., 2020; Nyambane et al., 2014; Omemu et al., 2014; Osvik et al., 2013; Sudi et al., 2011) are widely consumed across Africa. Despite consumer preferences for fresh meat and seafood, the combination of a warm, tropical climate and variable access to refrigeration facilities, particularly in rural areas, means that meat and seafood products are usually available in processed form. Smoking, salting, and drying are common preservative strategies for meat and seafood products in Africa (El-Sheika et al., 2014). Compared to other parts of the world, meat and fish fermentation is not widespread across Africa. However, fermented meats, e.g., *sujuk* and *boubanita*, are consumed in different North African countries (Benkerroum, 2013). Fermented fish products serve as both flavouring agents and a source of protein (Anihouvi et al., 2007) and can be produced from a wide variety of fish species. A significant amount of protein intake in the African diet comes from plant sources. The alkaline fermentation of proteinaceous oil seeds such as locust bean, melon, sesame, and castor oil have widespread use as flavour enhancers and meat substitutes (Ahaotu et al., 2013; Ezeokoli et al., 2016). These seeds are often inedible in their natural state as they contain anti-nutrients such as indigestible oligosaccharides and phytates, which are metabolised during fermentation (Parkouda et al., 2009). Plant leaves also serve as substrates for fermentation. *Kawal* and *ntoba mbodi* are produced from the fermentation of the leaves of *Cassia obtusifolia* and *Manihot esculenta*, respectively (Dirar, 1985; Voudibio Mbozo et al., 2017).

### **3. Fermenting and technologically important microbiota of African indigenous fermented foods**

Accurate identification and characterisation of the microbiota of IFFs is a critical first step for the selection of multi-functional starter cultures. Starter cultures are required for the controlled and large-scale production of IFFs with improved quality and safety attributes (Ahaotu et al., 2017; Edema and Sanni, 2008; Soro-Yao et al., 2014). The last four decades have seen an increase in research studies aimed at elucidating the microbial consortia present in IFFs from Africa. These studies have provided data on the vast diversity of microbial species, including lactic acid bacteria

(LAB), *Bacillus*, and yeasts that dominate the fermentation of cereals, legumes, dairy, and root crops (Table 1).

Lactic acid bacteria dominate the fermentation of cereals, dairy products, and starchy root tubers.

Of the LAB, species belonging to *Lactobacillus*, *Lactococcus*, *Weissella*, *Leuconostoc*, *Streptococcus*, and *Enterococcus* are frequently isolated (Diaz et al., 2019; Oguntinyinbo et al., 2011).

Previously, the beneficial effects of fermented foods were focused mainly on preservation and desirable organoleptic characteristics. Nowadays, a better understanding of the microorganisms involved in fermentation processes has drawn more attention to the various health benefits associated with IFFs. Lactic acid bacteria are among the most studied microorganisms in this regard, given their interesting technological characteristics. During fermentation, LAB metabolise sugars to produce a wide range of organic acids, including lactic, acetic and propionic acid, which produce a low pH environment that is detrimental to the growth and survival of spoilage and pathogenic organisms (Caplice & Fitzgerald, 1999).

Other antimicrobial metabolites associated with LAB metabolism include ethanol, hydrogen peroxide, diacetyl, and bacteriocins showcasing the diversity of LAB in utilising different metabolic pathways and substrates (Oguntinyinbo and Narbard, 2015; Mokoena et al., 2016; Soro-Yao et al., 2014). As LAB species are important members of the gut microbiome, their contribution to human health as probiotics has been extensively studied. Positive actions have been reported to include restoration of gut microbiota after antimicrobial therapy, vitamin production, and stimulating the immune system (Markowiak and Slizewska, 2017). A role for IFFs from Africa as delivery vehicles for probiotics is receiving significant attention (Achi & Ukwuru, 2015; Mokoena et al., 2016). Recently, two cereal-based fermented foods, *obushera* and *kwete*, have been produced using probiotic strain *Lactobacillus rhamnosus* yoba (Mukisa et al., 2019; Wacoo et al., 2019). Although the probiotic strain is not of African origin, the inherent presence of LAB species in IFFs from Africa suggests that IFFs may be sources of novel probiotic strains and warrants further studies.

The presence of *Enterococcus* spp., notably *Enterococcus faecium* and *Enterococcus faecalis*, in food systems is complex due to their increasing medical relevance as aetiological agents of nosocomial infections and their ability to disseminate antibiotic resistance determinants, particularly vancomycin (Oladipo et al., 2013; Oguntinyinbo and Okesuo, 2013). However, *Enterococcus* spp. are usually observed in IFFs, particularly in fermented dairy products where they are considered as contaminants (Jans et al., 2017). Although not recognised on the Qualified Presumption of Safety list (EFSA Panel on Biological Hazards, 2021), the positive technological contributions of enterococci include the production of extracellular polysaccharides, which influence the texture of fermented dairy products, and flavour and aroma development (Jans et al., 2017; Obioha et al., 2021). Some strains also produce bacteriocins, and their consideration for probiotics is receiving more research attention (Hanchi et al., 2018).

*Saccharomyces cerevisiae* is the most commonly isolated fungal species from African IFFs and has an essential role in the alcoholic fermentation of cereals and palm sap. Other dominant genera include *Candida*, *Pichia* and *Kluyveromyces* (Table 1). In addition to fermenting the sugars available in the natural substrates to alcohol and carbon dioxide, fungi produce flavour and aroma compounds such as organic acids, esters and carbonyls and contribute to textural changes via pectinases and cellulases (Cason et al., 2020; Johansen, 2019). Yeasts may also contribute to organic acid production by LAB species present during the fermentation (Ferreira and Mendes-Faia, 2020). Fungi, therefore, play a significant role in the final organoleptic characteristics of alcoholic IFFs over and above alcohol production.

*Bacillus* spp. dominate the fermentation of protein-rich legumes and seeds. An important aspect of the production process of alkaline fermented condiments such as *iru*, *soumbala* and *bikalga* produced from these protein-rich substrates is the long cooking time, and this heating process may select for spore formers that are more heat resistant (Ouoba et al., 2007; Parkouda et al., 2009). In addition, the degradation of proteins during fermentation by *Bacillus* spp., most significantly, *B. subtilis*, *B. pumilus*, and *B. licheniformis*, leads to the accumulation of peptides and ammonia. This

leads to an increase in pH, which also favours the proliferation of *Bacillus* spp. (Ouoba et al., 2008). *Bacillus* spp. are also frequently isolated from fermented cassava products (Anyogu et al., 2014; Assanvo et al., 2017; Padonou et al., 2009). Their ability to produce enzymes that hydrolyse cassava tissue has been reported to be responsible for the textural changes that occur during fermentation (Amoa-Awua & Jakobsen, 1995).

Molecular based techniques including fingerprinting-based methods, e.g., repetitive element palindromic (rep)-, intergenic transcribed spacing (ITS)-PCR in combination with the sequencing of ribosomal RNA (16S and 26S) or other housekeeping genes, e.g. *rpoA*, *rpoB*, *gyrA* and *pheS* are now routinely used to identify and characterise microorganisms in IFFs (Oguntinyinbo and Okesuo, 2012; Owusu-Kwarteng et al., 2012; Tadesse et al., 2019). These have provided new insights compared to earlier studies, which relied only on phenotypic identification (Abegaz, 2007; Nyambane et al., 2014), which do not always provide sufficient information. For example, molecular studies on submerged cassava fermentation have highlighted the dominance of *Weissella confusa* (Anyogu et al., 2014; Padonou et al., 2009), a microorganism that had not been previously reported in studies relying on phenotypic methods (Omafuvbe et al., 2007; Oyewole & Odunfa, 1988). *Weissella* spp. are also now frequently identified in other IFFs, e.g. cereals (Angelov et al., 2017; Oguntinyinbo et al., 2011) and dairy products (Akabanda et al., 2013). The extensive species diversity revealed by the use of genotyping techniques has been reported for other IFFs (Achi, 2005; Aderigbigbe et al., 2011; Nwaiwu and Itumoh, 2017; Ouoba et al., 2012) and has also uncovered new microbial species (Ouoba et al., 2015, a, b).

More recently, culture-independent methods such as PCR-DGGE and next-generation sequencing techniques are used to investigate the metagenomics associated with IFF production (Cason et al., 2020; Djeni et al., 2020; Walsh et al., 2017). An advantage of this approach is that it provides detailed information about the microbial community associated with different fermentation stages without the need for isolation. This results in a less biased microbial profile compared to culture-dependent methods as IFFs may contain uncultivable species (Bigot et al., 2015). Accurate data of

the microbial community involved during IFF production eliminates chance isolates that can be recovered on agar from the screening process for potential starter cultures. These more recent reports have confirmed that the microbes present in IFFs are more diverse than earlier reported. While investigating the microbial community of *soy-daddawa*, Ezeokoli et al. (2018) identified *Exiguobacterium* spp. for the first time. Diaz et al. (2019) observed *Zymomonas* spp. for the first time in fermented cereal, dairy, cassava and locust bean products. Although the contribution of these organisms to the fermentation process requires further study, their identification reveals new insights into IFFs. The factors that influence the composition of the microbial community has been noted to include fermentation conditions, pH changes, and geographical location (Houngbedji et al., 2018; Moonga et al., 2020). However, metagenomic studies have revealed that geographical location is not a consistent factor (Cason et al., 2020; Parker et al., 2018). Understanding the microbes involved in African IFF production will help in the design of starter cultures. However, culture-independent methods rely solely on DNA, so do not allow for the direct selection of microbial starters. Depending on the method used, DNA may be extracted from live and dead cells (Mukisa et al., 2012; Diaz et al., 2019). These limitations may be overcome by using both culture-dependent and independent methods (Adewunmi et al., 2012; Schoustra et al., 2013).

Despite these recent developments, there remain significant knowledge gaps concerning the microbiota of IFFs from Africa. Our review of the published research in this area highlights that further studies are required, particularly for alcoholic beverages, fermented meat and fish products (Gagaoua & Boudechicha, 2018; Djeni et al., 2020). These gaps limit the transition from household to large-scale controlled production of IFFs with consistent quality and safety attributes.

#### **4. Microbial hazards in indigenous fermented foods**

Globally, food safety remains a significant challenge. The World Health Organisation (WHO) estimates that as many as 1 in 10 people fall ill, and more than 120,000 children under 5 die each year after consuming unsafe food. Africa bears a high burden of the global incidence of foodborne

illness with an estimated annual morbidity of 90 million (WHO, 2015; WHO 2017b). Microbial hazards, including foodborne pathogens and their toxins, are primary aetiological agents of foodborne disease (FBD) and a growing public health issue.

Fermented foods are generally considered safe. Fermenting organisms, especially LAB, produce a range of antimicrobial compounds, e.g., organic acids, ethanol, bacteriocins and hydrogen peroxide, which are antagonistic to the growth and survival of foodborne pathogens (Adinsi et al., 2017; Devuyt and Vandamme, 1996; Mpofu et al., 2016). Cason et al. (2020) reported the decline of pathogenic and spoilage organisms during cereal fermentation for *sesotho* production. Similar observations were made by Karamoko et al. (2012), who noted a 4 – 7 log reduction in faecal coliform counts within a 24 h period in fermenting palm wine. When investigating the microbiological quality of milk products in Tanzania, Schoder et al. (2013) detected *Salmonella* and *Escherichia coli* in raw milk but not in fermented milk. In addition, Oguntinyinbo and Narbad (2015) isolated bacteriocin producing *Lactobacillus plantarum* strains from *kunu* and *ogi* that showed antimicrobial activity against *Salmonella enterica*.

However, indigenous practices for food production are often based on spontaneous fermentation, i.e., chance inoculation or the use of backslopping where utensils from a previous fermentation are reused (Caplice & Fitzgerald, 1999). Limited knowledge and utilisation of Hazard Analysis and Critical Control Points (HACCP) and good manufacturing processes (GMP) by farmers, food producers and handlers can lead to production and processing occurring under unhygienic conditions. These factors lead to variation in the microbial profile of IFFs, and consequently, the presence of spoilage and pathogenic bacteria in these foods cannot be ruled out (Oguntinyinbo, 2014; Olasupo et al., 2016).

Despite the lack of surveillance of foodborne infections in many African countries, several studies have investigated the prevalence of major foodborne pathogens in IFFs from Africa. Reports on microbiological hazards associated with IFFs from Africa are presented in Table 2. Microbes of public health significance, including *Bacillus cereus*, *Staphylococcus aureus*, and *Listeria monocytogenes*,

have been reported in IFFs from Africa. The introduction of pathogenic organisms can occur at different stages of the value chain via raw materials, the processing environment, and food handlers. An evaluation of the microbiological quality of water used for processing, fermenting broth, and the fermented cassava product, *lafun*, identified microbial hazards such as coliforms, including *Salmonella* spp. and *Staphylococcus* spp. (Lateef & Ojo, 2015). Potentially pathogenic bacteria have been found in utensils used for the fermentation process (Gran et al., 2002; Jans et al., 2017); however, only a few studies investigate the complete production chain to identify the source of contamination (Ademola et al., 2018; Thorsen et al., 2015). Adedeji et al. (2017) reported similarities in the microbial profiles of potentially pathogenic bacteria isolated from two fermented condiments from the same producer at the retail level. It is important to note that these bacteria were not present in the raw materials, suggesting unhygienic processing and handling. This underscores the need for further research to identify contamination sources to support the management of food safety hazards.

The microbiological safety of fermented vegetable proteins usually dominated by *Bacillus* spp. and *Staphylococcus* spp. requires consideration, as acid production, a potent antimicrobial attribute in lactic fermented IFFs, is not present (Ahaotu et al., 2013; Ouoba et al., 2019). *Bacillus cereus* and *Staphylococcus aureus* can produce toxins in food during their growth and have been identified in these products (Oranusi et al., 2015). *Bacillus* spp. are the dominant organisms involved in the fermentation of oil bean seeds; however, the presence of *Bacillus cereus* is routinely reported (Ahaotu et al., 2013; Ouoba et al., 2008b; Parkouda et al., 2009; Thorsen et al., 2015). Ahaotu et al. (2013) isolated *Bacillus cereus* capable of producing enterotoxins in *ugba* under fermentation conditions. A similar observation was made by Ouoba et al. (2008b) when investigating *B. cereus* involved in locust bean fermentation for soumbala production. However, these are heat-labile toxins that should be denatured with adequate cooking. Thorsen et al. (2015) detected the heat-stable, emetic type toxin-producing *B. cereus* strains in fermented baobab seeds, which is of concern.

The metabolic activities of some microorganisms involved in the fermentation process provide antagonistic conditions to the growth and survival of foodborne pathogens. However, inappropriate handling and the use of unsanitary packaging material can introduce microbial hazards post-processing (Adinsi et al., 2017; Mpofu et al., 2016; Schoder et al., 2013). The occurrence of potentially pathogenic bacteria at the retail level across all food categories highlights the potential risks to public health associated with IFFS (Odom et al., 2012; Owusu-Kwarteng et al., 2018). Of particular concern is the isolation of pathogenic bacteria from cereal-based and dairy fermented products, some of which are used as weaning or complementary foods (Adekoya et al., 2019; Samet-Bali et al., 2016). The presence of these organisms in ready-to-eat products suggests that IFFs may serve as vehicles of pathogenic bacteria. Therefore, the safety of these foods should not be taken for granted or assumed. The use of next-generation sequencing methods has provided more insight in identifying microbial hazards in IFFs, highlighting a role for metagenomic approaches as food safety tools (Walsh et al., (2017).

## **5. Antibiotic resistance of microbes from indigenous fermented foods from Africa**

The availability of antibiotics represented a landmark achievement in medicine and led to a significant decrease in mortality and morbidity from infectious disease agents (Spellberg, 2014). However, in recent times, an increasing number of pathogenic bacteria, including those that cause foodborne diseases, have become resistant to treatment with the antibiotic drugs currently available (WHO, 2017b). This scenario, termed antibiotic resistance (AR), has become one of the greatest threats to global public health and food security (McEwen & Collingon, 2018). It has been estimated that if left unchecked, the number of deaths attributed to AR each year could rise to 10 million by 2050 (O'Neill, 2016).

Antibiotic resistance is increasingly recognised as a food safety issue. The consumption of food contaminated with AR foodborne pathogens such as *Salmonella* could lead to treatment failure. Commensal bacteria such as *Escherichia coli* and *Enterococcus* spp. in food may transfer AR genes to human pathogens (WHO, 2017b). Drivers of AR include the overuse of antimicrobial drugs in human

and veterinary medicine. In addition to treatment, antimicrobials are used in agriculture and aquaculture as prophylaxis and in some parts of the world as growth promoters (Nhung et al., 2017; Van Boeckel et al., 2014). These factors lead to the spread of AR bacteria and AR genes (ARG) in the environment.

A comprehensive evaluation of the safety of IFFs in Africa should consider their role as vehicles of both pathogenic and AR bacteria. However, in Africa, the contribution of the food chain to the overall burden of AR is mostly unknown as there are little to no surveillance systems for monitoring antibiotic use in agriculture and food products (Founou et al., 2016; Oloso et al., 2018). For the most part, antibiotic use remains unregulated in Africa (Oguntinyinbo & Okesuo, 2012; Van et al., 2020).

Reports of AR bacteria associated with some African IFFs are presented in Table 3. Some of these studies have documented multiple drug-resistant (MDR)- bacteria in technologically relevant, indicator and potentially pathogenic organisms. Phenotypic resistance to more than three classes of antibiotics was observed in *Escherichia coli* O157:H7 and *Shigella* spp. isolated from cheese products available for retail sale in Egypt (Ahmed and Shimamoto, 2015 a,b). Fowoyo and Ogunbawo (2017) isolated 255 coagulase-negative staphylococci (CoNS) from fermented dairy, cereal and oilseed products in Nigeria. Of the total isolates, 27% exhibited MDR-phenotypes. Oguntinyinbo and Okesuo (2012) observed MDR *Enterococcus* spp. in *wara*, a fermented cheese. Ouoba et al., (2019) also identified MDR-*Staphylococcus* spp. in fermented vegetable products. However, other reports showed AR to one or two classes or overall sensitivity to the antibiotics tested. Owusu-Kwarteng et al., (2017) noted that *Bacillus cereus* isolated from fermented dairy products in Ghana only showed resistance to beta-lactams. *Enterococcus* spp. in IF products in Tunisia were shown to be sensitive to beta-lactams, aminoglycosides and macrolides (Rehaim et al., 2016).

Vancomycin is a critically important antibiotic, often used as a last resort treatment (WHO, 2019). Vancomycin resistance has been noted in isolates from African IFFs (Awopetu et al., 2016; Oguntinyinbo & Okesuo, 2012; Ouoba et al., 2008a). However, Rehaim et al., (2016) reported that *Enterococcus* spp. isolated from Tunisian fermented food products were susceptible to vancomycin.

A similar observation was also noted by Owusu-Kwarteng et al., (2017, 2018) when investigating *B. cereus* and *L. monocytogenes* in fermented dairy products.

It is important to note that most bacterial species have intrinsic and induced resistance mechanisms to specific antibiotic drugs (Reygaert, 2018). From a food safety perspective, the ability of foodborne bacteria to transfer resistance traits via mobile genetic elements to other bacteria is a primary concern. There are some reports of AR genetic determinants associated with African IFFs. Ahmed and Shimamoto (2015a) screened shiga toxin-producing *Escherichia coli* for antibiotic-resistant genes (ARG) and identified beta-lactamase encoding genes in all (n=5) isolates. Two of these isolates possessed plasmid-mediated quinolone resistance genes. Conjugation experiments undertaken by Ouoba et al., (2019) showed that CoNS from fermented condiments produced transconjugants with increased resistance to erythromycin and tetracycline. Unfortunately, most investigations of AR in African IFFs only consider phenotypic resistance (Adimpong et al., 2012; Awopetu et al., 2016; Eruteya and Eze, 2017). For this review, we found only one study (Aka et al., 2020) where ARGs in IFFs from Africa were investigated using whole-genome sequencing.

The prevalence of AR phenotypes in bacteria isolates from retail level IFFs in Africa and the high levels of resistance observed in some strains provide some evidence that these foods may serve as a reservoir for AR and is of public health significance. There remain considerable research and data gaps in this area, underscoring the need for large scale and long-term surveillance studies coordinated at national and international levels.

## **6. Discussion & Recommendations**

According to the United Nations, 50% of global population growth between now and 2050 is anticipated to occur in Africa (UN, 2020). However, Africa currently has the 2<sup>nd</sup> highest number of undernourished people globally, and about 20% of the population on the continent is already considered to be food insecure (FAO et al., 2020). The contribution of IFFs to combating food insecurity via increased food availability, improved nutrition and income generation is well established (Franz et al., 2014; Okafor, 1992; Rolle and Satin, 2002; Setta et al., 2020).

The global market for fermented products is predicted to surpass \$20 billion by 2022 (Sivamaruthi et al., 2019). This demand is fuelled in part by the increasing popularity of some IFFs, such as *kombucha* and *kefir*, in the international market as consumers become more aware of their beneficial effects (Soni et al., 2014). Besides meeting domestic demand from an increasingly urban populace, African IFFs could also become a source of foreign revenue via exports for a growing diaspora community and beyond. This requires improving the value chains that deliver IFFs with consistent quality and safety attributes to local and international markets using modern, industrial processes, including well-characterised starter cultures.

More work remains to be done towards achieving large scale production of IFFs from Africa. However, there are some success stories. For example, the production of the widely consumed sorghum-based fermented beer *umqombothi* has been industrialised in South Africa. Cereal-based porridges *ogi* and *uji*, and the alcoholic beverage palm wine are also now produced commercially (Adebo, 2020; Nwaiwu & Itumoh, 2017). Significant advances have been made in identifying dominant strains in IFFs and characterising technological aspects which make them suitable for use in controlled fermentations (Aderigbigbe et al., 2011; Ahaotu et al., 2013; Aka et al., 2020; Houngbédji et al., 2018; Moodley et al., 2019; Oguntinyinbo & Narbad, 2015; Sawadogo-Lingani et al., 2008). However, transferring these technologies to producers presents some challenges, including the stability, activity, and viability of these cultures (Benkerroum & Tamime, 2004; Benkerroum, 2013; Rolle and Satin, 2002). Although several studies have identified and characterised potential starters for use in soumbala fermentation, most processors still utilise spontaneous fermentation processes (Ouoba et al., 2004; Compaore et al., 2020; Parkouda et al., 2009). To overcome some of these obstacles, an alternative approach used by some researchers is the utilisation of commercially available starter cultures, as has been reported for the production of *obushera* using *Lact. rhamnosus* yoba (Mukisa et al., 2019) and *pito* using *Lact. delbrueckii* and *Sacch. cerevisiae* (Djameh et al., 2019). However, this may not be a sustainable solution due to cost implications.

Selecting starter cultures from the autochthonous community is often recommended as these cultures are considered more adapted to the fermentation parameters of the IFF and may show wider metabolic diversity required for achieving desirable properties and functional characteristics of the product (Ashaolu, 2019; Casquete et al., 2011). However, there is a lack of data showing if autochthonous fermenting organisms possess unique biochemical properties compared to similar strains obtained from other sources, which could be further explored. There is also a need for more studies that evaluate the application of starter cultures as monocultures or in combination for IFF production outside the laboratory environment (Compaore et al., 2020; Kimaryo et al., 2000; Mukisa et al., 2016). Beyond technology transfer, more progress is required to improve raw materials, conduct laboratory, and pilot plant production before scaling up to industrial production (Okafor et al., 1992; Benkerroum, 2013).

Food fermentations are complex microbial ecosystems (De Filippis et al., 2017). The use of metagenomics in studying African IFFs has provided more insight into the microbiota and succession dynamics of fermenting microorganisms (Cason et al., 2020; Diaz et al., 2019). However, only a limited number of IFFs have been investigated using this approach, and future research efforts should be directed here. Metagenomic data can be combined with outputs from other 'omics' technologies such as metabolomics and transcriptomics to develop a more comprehensive understanding of the relationships between the microorganisms present in the food, their metabolic interactions, and what these contribute to the fermentation process (De Filippis et al., 2017; Kergourlay et al., 2015). This will allow for a more informed starter selection process for improved fermentation processes.

The prevalence of foodborne disease remains severe in many African countries. Factors contributing to this include food preparation with contaminated water, poor hygiene, inadequate storage facilities, food safety knowledge, and insufficient food safety legislation and implementation (Belli et al., 2013). Currently, IFFs are mainly marketed via the informal economy in open markets, street vending and household producer/seller, therefore outside the scope of official health regulatory

standards where these exist. The reviewed studies suggest that contamination of IFFs mainly occurs post-processing. Similar to our observations, a meta-analysis of the prevalence of foodborne pathogens in ready-to-eat food from seven African countries showed that *Enterobacteriaceae*, *Escherichia coli*, *Salmonella*, *Staphylococcus aureus*, and *Listeria monocytogenes* were the most frequently reported organisms (Paudyal et al. 2017).

Given the potential for the introduction of food safety hazards at each production stage, the design and implementation of quality assurance and management systems, including GMP and HACCP in commercial food production, is recommended or incorporated into legislation as an effective strategy for improving food safety (Kafetzopoulos, 2013; Ropkins & Beck, 2000). There are now several reports on the development of HACCP systems for IFF production (Asagbra et al., 1998; Fasoyiro et al., 2010; Lateef & Ojo, 2015; Oguntoyinbo, 2012; ). Some studies have demonstrated the effective use of these approaches in improving the microbiological quality and safety attributes of some products such as *lafun* (Obadina et al., 2008) and *kenkey* (Amoa-Awua et al., 2007). However, there is a need for more data evaluating the application of HACCP systems in the commercial production of IFFs and on studies that report on producers and handlers' food safety knowledge, attitudes, and practices. Byakiya et al., (2019) noted that processors of *Obushera*, a fermented beverage widely consumed in Uganda, showed good food safety knowledge but observed poor hygienic practices. Amoa-Awua et al., (2007) reported significant improvements in the safety characteristics of *kenkey* after the implementation of a HACCP and GMP system. However, they noted that the level of formal education of processors was a significant hurdle when applying quality assurance systems in IFF production. After observing poor manufacturing and hygienic processes during the production of *ice kenkey* (Atter et al., 2015), researchers developed a simplified manual to train *ice kenkey* processors in GMP & HACCP principles. These studies highlight the need for more extension work, including training on basic food safety and hygiene, GMP, and HACCP, for food producers and handlers.

Assessing health risks to consumers from IAFFs requires more and better-quality data to underpin quantitative exposure assessments. There is a scarcity of research studies that focus on the microbiological safety of African IFFs. In some cases, the observation of pathogenic or indicator bacteria are incidental, occurring where investigation of the fermented product's microbial community is the main objective (Anyogu et al., 2014; Oranusi et al., 2015; Parkouda et al., 2010). This means that beyond identification, important information such as microbial load, virulence factors, or antimicrobial resistance determinants are not investigated and recorded. Another constraint in the quality of data collected is study design. Some reports do not include the number of samples collected to support estimating prevalence or utilised convenience sampling, which may not be representative. Many reports rely on conventional techniques for identifying foodborne pathogens, which may be misleading or do not provide sufficient information, e.g., species identification. While the presence of potential pathogens in a food product is a cause for concern, health risks associated with consumption must consider any national, regional, or international safety standards, such as the Codex Alimentarius. Except for a few exceptions (Byakika et al., 2019; Kouame et al., 2013; Gran et al., 2002), these standards are often not referred to indicate the microbiological quality of the food being studied. Antibiotic resistance of microorganisms has not been extensively studied in African IFFs. A significant AR data gap also exists in African clinical settings. In a systematic analysis, Tadesse et al., (2017) reported that AR data was not available for as many as 40% of African countries. The analysis concluded that resistance to commonly prescribed antibiotics was prevalent, and the quality of microbiological data is of serious concern. In the absence of rigorous government surveillance and regulation for IFF production in many parts of Africa, more studies will need to be carried out by the scientific community to raise the necessary awareness required and monitor prevalence trends, especially of acquired AR. Many of the technologies required to investigate IFFs and produce the innovation needed in the sector remain inaccessible to many researchers in low and middle-income countries (LMICs) due to

resource constraints in technical know-how and infrastructure. This limits opportunities to publish in more impactful journals, reducing the visibility of the outputs obtained, despite their importance to the scientific community. There is a need for more initiatives that support sustainable international collaborative efforts that bring together scientists in LMICs and High-income countries to share expertise and develop equitable capacity building research activities. An example of this is the ENRECA/DANIDA project, “Capability Building for Research and Quality Assurance in Traditional Food Processing in West Africa”, which has supported several successful collaborative research efforts in value-added processing of IFFs in some West African countries.

Research innovation must translate into improvements in IFF processing technology. This requires significant investment in stakeholder management between policymakers, scientists, and producers. Public sector funding could target knowledge transfer partnerships between research institutions and small-medium- enterprise (SME) producers, which must benefit both partners. Research projects can be designed in collaboration with SMEs, focusing on the real-world problems encountered by producers (Moodley et al., 2019). This model can contribute towards the buy-in of producers, required to drive change along the food processing value chain.

Weak enforcement of food safety regulations poses risks to public safety. Additionally, surveillance systems that should provide accurate and reliable data on the burden of foodborne illness in IFFs are often inadequate. This means that the number of illnesses or outbreaks associated with IFFs may go unreported. In 2012, an outbreak of botulism in Canada was linked to an African fermented fish product, fesikh, which led to a voluntary withdrawal from sale by the manufacturer (Walton et al., 2014). Many public health agencies across Africa have limited access to the required infrastructure to gather epidemiological evidence to support this type of timely action.

‘One Health’ is a transdisciplinary approach introduced by the WHO as a framework to be used by relevant stakeholders in developing and implementing strategies to safeguard public health (WHO, 2017c). The African Center for Disease Control (Africa CDC) has recently published a ‘One Health’ framework for managing zoonotic infections (Africa CDC, 2020). However, food safety and

antimicrobial resistance are also important priority areas for public health, requiring urgent attention in the African region. At a national level, policymakers also need to empower regulatory agencies with the required legal frameworks and infrastructure to develop and enforce food safety standards.

## **7. Conclusion**

Indigenous fermented foods have great potential in combating food insecurity in Africa and harbour a vast genetic potential of valuable undiscovered strains. To achieve the goal of improving and scaling up fermentation technology, the use of advanced molecular biology tools, including whole-genome sequencing, is required to accurately identify the microbial community in IFFs, both beneficial and harmful. A comprehensive understanding of the microbial community of IFFs could identify biomarkers for assessing the quality and safety attributes of these foods, including technological characteristics, virulence factors and antibiotic resistance.

The presence of pathogenic & AR bacteria in ready to eat IFFs constitute a risk to public health. Harmful bacteria may enter the food chain via the raw material, inadequate fermentation to lower the pH sufficiently in lactic fermented foods or post-processing contamination. The production of antimicrobial compounds by fermenting organisms may be insufficient to eradicate pathogenic organisms in the final product, and they are not substitute for GMP. Hygiene improvement in handling raw food, increased surveillance, and uniform protocols for sampling and identification is suggested to help evolve a common approach to studies of indigenous foods.

Any strategies to guarantee IFFs free of microbial hazards require considerable investment and collaboration from relevant stakeholders – consumers, producers, industry, policymakers, and scientists.

## **Acknowledgements**

The authors are grateful for the support of the Nigerian Applied Microbiologists network in developing this collaborative effort.

## **Funding**

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

#### **Author's contributions**

Amarachukwu Anyogu: Conceptualisation, Data Collection, Formal Analysis, Writing - Original draft preparation, Project Administration. Ogueri Nwaiwu: Conceptualisation, Data Collection, Writing – Original draft preparation, Writing – Review and Editing, Supervision. Ayomide Olukorede: Data Collection, Writing – Original draft preparation. Writing – Review and Editing. Christian Anumudu: Data Collection, Writing – Original draft preparation. Helen Onyeaka: Data Collection, Writing – Review and editing. Esther Areo: Data Collection, Writing – Original draft preparation. Obadina Adewale: Writing – Review and editing, Supervision. All authors read and approved the final manuscript.

**Declarations of interest:** None

#### **References**

- Abd-Allah, S. M. S. (2011). Bacterial-flora of Egyptian salted *Mugil cephalus* fish (fessiekh) PCR-identification. *Assiut Veterinary Medical Journal*, 57, 116-137.
- Abdelgadir, W., Nielsen, D.S., Hamad, S., & Jakobsen, M. (2008). A traditional Sudanese fermented camel's milk product, Gariss, as a habitat of *Streptococcus infantarius* subsp. *infantarius*. *International Journal of Food Microbiology*, 127, 215–219. <https://doi.org/10.1016/j.ijfoodmicro.2008.07.008>.
- Abegaz, K., (2007). Isolation, characterization and identification of lactic acid bacteria involved in traditional fermentation of borde, an Ethiopian cereal beverage. *African Journal of Biotechnology*, 6, 1469-1478.
- Abiodun, O.A., Ayano, B., Amanyunose, A.A. (2020). Effect of fermentation periods and storage on the chemical and physicochemical properties of biofortified cassava gari. *Journal of Food Processing and Preservation*, 44, e14958, <https://doi.org/10.1111/jfpp.14958>
- Abriouel, H., Ben Omar, N., López, R.L., Martínez-Cañamero, M., Keleke, S., & Gálvez, A. (2006). Culture-independent analysis of the microbial composition of the African traditional fermented foods poto poto and dégué by using three different DNA extraction methods. *International Journal of Food Microbiology*, 111, 228-233. <https://doi.org/10.1016/j.ijfoodmicro.2006.06.006>.
- Achi, O. K. (2005). Traditional fermented protein condiments in Nigeria. *African Journal of Biotechnology*, 4, 1612 – 1621.
- Achi, O.K., & Akubor, P.I. (2000). Microbiological characterization of yam fermentation for 'Elubo' (yam flour) production. *World Journal of Microbiology and Biotechnology*, 16, 3 – 7.

<https://doi.org/10.1023/A:1008980912708>.

Achi, O.K., & Ukwuru, U. (2015). Cereal-based fermented foods of Africa as functional foods. *International Journal of Microbiology and Application*, 2, 71 - 83.

Adebayo-Oyetoro, A.O., Oyewole, O.B., Obadina, A.O., Omemu, M.A. (2013). Microbiological Safety Assessment of Fermented Cassava Flour “Lafun” Available in Ogun and Oyo States of Nigeria. *International Journal of Food Science*, 845324. <https://doi.org/10.1155/2013/845324>.

Adebo, O.A. (2020). African Sorghum-Based Fermented Foods: Past, Current and Future Prospects. *Nutrients*, 12, 1111. <https://doi.org/10.3390/nu12041111>.

Adediji, B.S., Ezeokoli, O.T., Ezekiel, C.N., Obadina, A.O., Somorin, Y.M., Sulyok, M., Adeleke, R.A., Warth, B., Nwangburuka, C.C., Omemu, A.M., Oyewole, O.B., & Krska, R. (2017). Bacterial species and mycotoxin contamination associated with locust bean, melon and their fermented products in south-western Nigeria. *International Journal of Food Microbiology*, 258, 73-80. <https://doi.org/10.1016/j.ijfoodmicro.2017.07.014>.

Adekoya, I., Obadina, A., Olorunfemi, M., Akande, O., Landschoot, S., De Saeger, S., & Njobeh, P. (2019). Occurrence of bacteria and endotoxins in fermented foods and beverages from Nigeria and South Africa. *International Journal of Food Microbiology*, 305:108251. <https://doi.org/10.1016/j.ijfoodmicro.2019.108251>.

Ademola, O.M., Adeyemi, T.E., Ezeokoli, O.T., Ayeni, K.I., Obadina, A.O., Somorin, Y.M., Omemu, A.M., Adeleke, R.A., Nwangburuka, C.C., Oluwafemi, F., Oyewole, O.B. & Ezekiel C.N. (2018). Phylogenetic analyses of bacteria associated with the processing of iru and ogiri condiments. *Letters in Applied Microbiology*, 67, 354-362. <https://doi.org/10.1111/lam.13040>.

Aderibigbe E. Y., Visessanguan, W., Sumpavapol, P., & Kongtong, K. (2011). Sourcing starter cultures for *Parkia biglobosa* fermentation: Phylogenic grouping of *Bacillus* species from commercial ‘iru’ samples. *International Journal of Biotechnology and Molecular Biology Research*, 2, 121-127. <https://doi.org/10.5897/IJBMBR.9000036>.

Adewunmi, G.A., Oguntoyinbo, F.A., Keisam, S., Romi, W., & Jeyaram, K. (2013). Combination of culture-independent and culture-dependent molecular methods for the determination of bacterial community of iru, a fermented *Parkia biglobosa* seeds. *Frontiers in Microbiology*, 3, 436. <https://doi.org/10.3389/fmicb.2012.00436>.

Adimpong, D.B., Nielsen, D.S., Sørensen, K.I., Derkx, P.M.F., & Jespersen, L. (2012). Genotypic characterization and safety assessment of lactic acid bacteria from indigenous African fermented food products. *BMC Microbiology*, 12, 75. <https://doi.org/10.1186/1471-2180-12-75>

Adinsi, L., Mestres, C., Akissoé, N., Vieira-Dalodé, G., Anihouvi, V., Durand, N., Hounhouigan, D.J. (2017). Comprehensive quality and potential hazards of gowe, a malted and fermented cereal beverage from West Africa. A diagnostic for a future re-engineering. *Food Control*, 82, 18-25. <https://doi.org/10.1016/j.foodcont.2017.06.019>.

Africa Centre for Disease Control. (2020). Framework for One Health Practice in National Public Health Institutes. Zoonotic Disease Prevention and Control. Available from: <https://africacdc.org/download/framework-for-one-health-practice-in-national-public-health-institutes/> Accessed 24.12.2020

- Ahaotu, I., Anyogu, A., Njoku, O.H., Odu, N.N., Sutherland, J.P., & Ouoba, L.I.I. (2013). Molecular identification and safety of *Bacillus* species involved in the fermentation of African oil beans (*Pentaclethra macrophylla* Benth) for production of Ugba. *International Journal of Food Microbiology*, 162, 95-104. <https://doi.org/10.1016/j.ijfoodmicro.2013.01.001>.
- Ahaotu, N.N., Anyogu, A., Obioha, P., Aririatu, L., Ibekwe, V.I., Oranusi, S., Sutherland, J.P., & Ouoba, L.I.I. (2017). Influence of soy fortification on microbial diversity during cassava fermentation and subsequent physicochemical characteristics of garri. *Food Microbiology*, 66, 165 – 172. <https://doi.org/10.1016/j.fm.2017.04.019>.
- Ahmed, A.M., & Shimamoto, T. (2015a). Molecular analysis of multidrug resistance in Shiga toxin-producing *Escherichia coli* O157:H7 isolated from meat and dairy products. *International Journal of Food Microbiology*, 193, 68-73. <https://doi.org/10.1016/j.ijfoodmicro.2014.10.014>
- Ahmed, A.M., & Shimamoto, T. (2015b). Molecular characterization of multidrug-resistant *Shigella* spp. of food origin. *International Journal of Food Microbiology*, 194:78-82. <https://doi.org/10.1016/j.ijfoodmicro.2014.11.013>
- Aka, S., Dridi, B., Bolotin, A., Yapo, E.A., Koussemon-Camara, M., Bonfoh, B., & Renault, P. (2020). Characterization of lactic acid bacteria isolated from a traditional Ivoirian beer process to develop starter cultures for safe sorghum-based beverages. *International Journal of Food Microbiology*, 322,108547. <https://doi.org/10.1016/j.ijfoodmicro.2020.108547>
- Akabanda, F., Owusu-Kwarteng, J., Tano-Debrah, K., Glover, R.L., Nielsen, D.S., & Jespersen, L. (2013). Taxonomic and molecular characterisation of lactic acid bacteria and yeasts in nunu, a Ghanaian fermented milk product. *Food Microbiology*, 34, 277 – 283. <https://doi.org/10.1016/j.fm.2012.09.025>.
- Akintonwa, A., Tunwashe, O. & Onifade, A. (1994). Fatal and non-fatal acute poisoning attributed to cassava-based meal. *Acta Horticulturae*, 375, 285-288.
- Alo, M. N., Eze, U. A., & Eda, N. E. (2012). Microbiological qualities of burukutu produced from a mixture of sorghum and millet. *American Journal of Food and Nutrition*, 2, 96-102. <https://doi.org/10.5251/ajfn.2012.2.4.96.102>.
- Amoa-Awua, W.K., Appoh, F.E., & Jakobsen, M. (1996). Lactic acid fermentation of cassava dough into agbelima. *International Journal of Food Microbiology*, 31, 87-98. [https://doi.org/10.1016/0168-1605\(96\)00967-1](https://doi.org/10.1016/0168-1605(96)00967-1).
- Amoa-Awua, W.K.A., Jakobsen M. (1995). The role of *Bacillus* species in the fermentation of cassava. *Journal of Applied Bacteriology*, 79, 250-256. <https://doi.org/10.1111/j.1365-2672.2006.03074.x>
- Amoa-Awua, W.K.A., Ngunjiri, P., Anlobe, J., Kpodo, K., Halm, M., Hayford, A.E. & Jakobsen, M. (2007). The effect of applying GMP and HACCP to traditional food processing at a semi-commercial kenkey production plant in Ghana. *Food Control*, 18, 1449-1457. <https://doi.org/10.1016/j.foodcont.2006.10.009>.
- Amoa-Awua, W.K., Sampson, E., & Tano-Debrah, K. (2007). Growth of yeasts, lactic and acetic acid bacteria in palm wine during tapping and fermentation from felled oil palm (*Elaeis guineensis*) in Ghana. *Journal of Applied Microbiology*, 102, 599-606. doi: 10.1111/j.1365-2672.2006.03074.x.

- Amosun, E.A., Agbato, A.O., Daodu, O.B., & Ojo, O.E. (2017). Evaluation of Bacteria and Antibiotic Resistance Profiling of Wara (White Soft Cheese) in Oyo-State, South-West, Nigeria. *Nigerian Veterinary Journal*, 38, 319-330.
- Angelov, A.I., Petrova, G., Angelov, A.D., Stefanova, P., Bokossa, I.Y., Tchekessi, C.K.C., Marco, M.L., & Gotcheva, V. (2017). Molecular Identification of Yeasts and Lactic Acid Bacteria Involved in the Production of Beninese Fermented Food Degue. *The Open Biotechnology Journal*, 11, 94-104. <http://dx.doi.org/10.2174/1874070701711010094>.
- Anihouvi, V., Ayernor, G. S., Hounhouigan, J. D., & Sakyi-Dawson, E. (2006). Quality characteristics of Lanhoun: A traditional processed, fermented fish product in the Republic of Benin. *African Journal of Food, Agriculture, Nutrition and Development*, 6, 1-15. <https://doi.org/10.4314/ajfand.v6i1.19173>.
- Anyogu, A., Awamaria, B., Sutherland, J.P. & Ouoba, L.I.I. (2014). Molecular characterisation and antimicrobial activity of bacteria associated with submerged lactic acid cassava fermentation. *Food Control*, 39, 119-127. <https://doi.org/10.1016/j.foodcont.2013.11.005>.
- Asagbra, A. Y., Odunfa, S. A., Amoa-Awua, W. A., & Jakobsen, M. (1998). HACCP system for traditional African fermented foods: Soy-ogi. Taastrup: WAITRO.
- Assanvo, J.B., Agbo, G.N., Coulin, P., Monsan, V., Heuberger, C., Kati-Coulibaly, S & Farah, Z. (2017). Influence of microbiological and chemical quality of traditional starter made from cassava on “attiéké” produced from four cassava varieties. *Food Control*, 78, 286-296. <https://doi.org/10.1016/j.foodcont.2017.02.066>.
- Atter, A., Ofori, H., Anyebuno, G.A., Amoo-Gyasi, M., Amoa-Awua, W.K.A. (2015). Safety of a street vended traditional maize beverage, ice-kenkey, in Ghana. *Food Control*, 55, 200-205. <https://doi.org/10.1016/j.foodcont.2015.02.043>.
- Awopetu, S.A., Adelowo, O.O., Asaolu, O., Ogunlade, B.O., & Afolabi, O.B. (2016). Antimicrobial susceptibility testing on isolated gram positive and gram negative bacteria from local unpasteurized milk (Fura de nunu). *International Journal of Medicine Research*, 1, 5 – 8.
- Ashaolu, T.J. (2019). A review on selection of fermentative microorganisms for functional foods and beverages: the production and future perspectives. *International Journal of Food Science and Technology*, 54, 2511 – 2519. <https://doi.org/10.1111/ijfs.14181>
- Azokpota, P., Hounhouigan, D.J., & Nago, M.C. (2006). Microbiological and chemical changes during the fermentation of African locust bean (*Parkia biglobosa*) to produce afitin, iru and sonru, three traditional condiments produced in Benin. *International Journal of Food Microbiology*, Volume 107, 304-309. <https://doi.org/10.1016/j.ijfoodmicro.2005.10.026>.
- Bayili, G.R., Johansen, P., Nielsen, D.S., Sawadogo-Lingani, H., Ouedraogo, G.A., Diawara, B., & Jespersen, L. (2019). Identification of the predominant microbiota during the production of lait caillé, a spontaneously fermented milk product made in Burkina Faso. *World Journal of Microbiology and Biotechnology*, 35, Article number: 100. <https://doi.org/10.1007/s11274-019-2672-3>.
- Belli, P., Cantafora, A.F.A., Stella, A., Barbieri, S., Crimella, C. (2013). Microbiological survey of milk and dairy products from a small scale dairy processing unit in Maroua (Cameroon). *Food Control*, 32, 366-370. <https://doi.org/10.1016/j.foodcont.2012.12.021>.

- Benkerroum, N. (2013). Traditional Fermented Foods of North African Countries: Technology and Food Safety Challenges with regard to Microbiological Risks. *Comprehensive Reviews in Food Science and Food Safety*, 12, 54 – 89. <https://doi.org/10.1111/j.1541-4337.2012.00215.x>.
- Benkerroum, N., Bouhlal, Y., El Attar, A., & Marhaben, A. (2004). Occurrence of Shiga Toxin-Producing *Escherichia coli* O157 in Selected Dairy and Meat Products Marketed in the City of Rabat, Morocco. *Journal of Food Protection*, 67, 1234–1237. <https://doi.org/10.4315/0362-028X-67.6.1234>.
- Benkerroum, N., & Tamime, A.Y. (2004). Technology transfer of some Moroccan traditional dairy products (lben, jben and smen) to small industrial scale. *Food Microbiology*, 21, 399–413. <https://doi.org/10.1016/j.fm.2003.08.006>.
- Bessam, H.B., Missouri, M., & Kridech, S. (2016). Bacterial Ecology of the “Kaddid”, Typical Dried Meat of the North Africa, During Its Traditional Fermentation. *Journal of Food and Nutrition Sciences*, 4, 70–77. <https://doi.org/10.11648/j.jfns.20160403.15>.
- Bigot, C., Meile, J.-C., Remize, F., & Strub, C. (2015). Applications of Metagenomics to Fermented Foods. In D. Monet & R.C. Ray (Eds.), *Fermented Foods, Part I, Biochemistry and Biotechnology*, (p 333 – 346), CRC Press.
- Birmeta, G., Bakeeva, A., & Passoth, V. (2019). Yeasts and bacteria associated with kocho; an Ethiopian fermented food produced from enset (*Ensete ventricosum*). *Antonie Van Leeuwenhoek*, 112, 651–659. <https://doi.org/10.1007/s10482-018-1192-8>.
- Byakika, S., Mukisa, I.M., Byaruhanga, Y.B., Male, D., & Muyanja, C. (2019). Influence of food safety knowledge, attitudes and practices of processors on microbiological quality of commercially produced traditional fermented cereal beverages, a case of Obushera in Kampala. *Food Control*, 100, 212–219. <https://doi.org/10.1016/j.foodcont.2019.01.024>.
- Caplice, E., & Fitzgerald, G.F. (1999) Food fermentations: role of microorganisms in food production and preservation. *International Journal of Food Microbiology*, 50, 131– 149. [https://doi.org/10.1016/S0168-1605\(99\)00082-3](https://doi.org/10.1016/S0168-1605(99)00082-3).
- Casquete, R., Benito, M.J., Martín, A., Ruiz-Moyano, S., Hernández, A. & Córdoba, M.G. (2011). Effect of autochthonous starter cultures in the production of “salchichón”, a traditional Iberian dry-fermented sausage, with different ripening processes. *LWT - Food Science and Technology*, 44, 1562–1571. <https://doi.org/10.1016/j.lwt.2011.01.028>.
- Cason, E.D., Mahlomaholo, B.J., Taole, M. M., Abong, G. O., Vermeulen, J-G., de Smidt, O., Vermeulen, M., Steyn, L., Valverde, A., & Viljoen, B. (2020). Bacterial and fungal dynamics during the fermentation process of Sesotho, a traditional beer of Southern Africa. *Frontiers in Microbiology*, 11, <https://doi.org/10.3389/fmicb.2020.01451>.
- Cissé, H., Muandze-Nzambe, J. U., Somda, N. S., Sawadogo, A., Drabo, S. M., Tapsoba, F., Zongo, C., Traoré, Y., & Savadogo, A. (2019). Assessment of safety and quality of fermented milk of camels, cows, and goats sold and consumed in five localities of Burkina Faso. *Veterinary world*, 12, 295–304. <https://doi.org/10.14202/vetworld.2019.295-304>.
- Compaoré, C.S., Tapsoba, F.W., Parkouda, C., Tamboura, D., Traoré, E.M.A., Diawara, B., Savadogo, A., Jespersen, L. & Sawadogo-Lingani, H. (2020). Development of starter cultures carrier for the

production of high quality soumbala, a food condiment based on *Parkia biglobosa* seeds. *African Journal of Biotechnology*, 19, 820 – 828. <https://doi.org/10.5897/AJB2020.17244>

De Filippis, F., Parente, E., & Ercolini, D. (2017). Metagenomics insights into food fermentations. *Microbial Biotechnology*, 10, 91–102. <https://doi.org/10.1111/1751-7915.12421>.

De Vuyst, L., & Vandamme, E.J. (1994). Antimicrobial potential of lactic acid bacteria. In L. De Vuyst & E.J. Vandamme (Eds.), *Bacteriocins of Lactic Acid Bacteria* (pp. 91 -143), Blackie Academic and Professional, New York.

Diaz, M., Kellingray, L., Akinyemi, N. Adefiranye, O.O., Olaonipekun, A.B., Bayili, G.R., Ibezim, J., du Plessis, A.S., Hounghédji, M., Kanya, D., Mukisa, I.M., Mulaw, G., Josiah, S.M., Chienjo, W.O., Atter, A., Agbemafle, E., Annan, T., Ackah, N.B., Buys, E.M., Hounhouigan, D.J., Muyanja, C., Nakavuma, J., Odeny, D.A., Sawadogo-Lingani, H., Tefera, A.T., Amoa-Awua, W., Obodai, M., Mayer, M.J., Oguntoyinbo, F.A., & Narbad, A. (2019). Comparison of the microbial composition of African fermented foods using amplicon sequencing. *Scientific Reports*, 9, 13863 <https://doi.org/10.1038/s41598-019-50190-4>

Dikko, A. H., Malik, A. A., & Egena, S. S. A. (2011). Chemical and bacteriological assessment of Kindirmo (local yoghurt) put on sale in Niger state. *Journal of Agriculture, Forestry and the Social Sciences*, 9, 245-248. <https://doi.org/10.4314/JOAFSS.V9I1.28>

Dirar, H. A., Harper, D. B., & Collins, M. A. (1985). Biochemical and microbiological studies on Kawal, a meat substitute derived by fermentation of *Cassia obtusifolia* leaves. *Journal of the Science of Food and Agriculture*, 36, 881–892. <https://doi.org/10.1002/jsfa.2740360919>.

Djameh, C., Ellis, W.O., Oduro, I., Saalia, F.K., Blay, Y.M. & Komlaga, G.A. (2019). West African sorghum beer fermented with *Lactobacillus delbrueckii* and *Saccharomyces cerevisiae*: shelf-life and consumer acceptance. *Journal of the Institute of Brewing*, 125, 333 – 341. <https://doi.org/10.1002/jib.563>

Djeni, T.N., Bouatenin, K.M.J.-P., Assohoun, N.M.C., Toka, D.M., Menan, E.H., Dousset, X., & Dje, K.M. (2015). Biochemical and microbial characterization of cassava inocula from the three main attiéké production zones in Côte d'Ivoire. *Food Control*, 50, 133-140. <https://doi.org/10.1016/j.foodcont.2014.08.046>.

Djeni, T.N., Kouame, K.H., Ake, F.D.M., Amoikon, L.S.T., Dje, M.K., & Jeyaram, K. (2020). Microbial Diversity and Metabolite Profiles of Palm Wine Produced From Three Different Palm Tree Species in Côte d'Ivoire. *Scientific Reports*, 10, 1715 <https://doi.org/10.1038/s41598-020-58587-2>.

Edema, M.O., & Sanni, A.I. (2008). Functional properties of selected starter cultures for sour maize bread. *Food Microbiology*, 25, 616 - 625. <https://doi.org/10.1016/j.fm.2007.12.006>.

EFSA BIOHAZ Panel, Koutsoumanis, K., Allende, A., Alvarez-Ordóñez, A., Bolton, D., Bover-Cid, S., Chemaly, M., Davies, R., Hilbert, F., Lindqvist, R., Nauta, M., Peixe, L., Ru, G., Simmons, M., Skandamis, P., Suffredini, E., Cocconcilli, P.S., Escamez, P.S.F., Maradona, M.P., Querol, A., Suarez, J.E., Sundh, I., Vlask, J., Barizzone, F., Correia, S., & Lieve, H. (2021). The list of QPS status recommended biological agents for safety risk assessments carried out by EFSA [Data set]. Zenodo. <http://doi.org/10.5281/zenodo.4428353>

- El Marnissi, B., Bennani, L., Cohen, N., El Ouali, A., & Belkhou, R. (2013). Presence of *Listeria monocytogenes* in raw milk and traditional dairy products marketed in the north-central region of Morocco. *African Journal of Food Science*, 7, 87 – 91.
- El Sheikha, A. F., Ray, R., Montet, D., Panda, S., & Worawattanamateekul, W. (2014). African fermented fish products in scope of risks. *International Food Research Journal*, 21, 425-432.
- Eruteya, O.C., & Eze, C.O. (2016). Proximate Composition and Antibiotic Resistant *Listeria* Species in Fura-de-nunu Sold in Port Harcourt. *Journal of Advances in Biology and Biotechnology*, 11, 1 -7. <https://doi.org/10.9734/JABB/2017/30655>.
- Essid, I., Ben Ismail, H., Bel Hadj Ahmed, S., Ghedami, R., & Hassouna, M. (2007). Characterization and technological properties of *Staphylococcus xylosus* strains isolated from a Tunisian traditional salted meat. *Meat Science*, 77, 204-12. <https://doi.org/10.1016/j.meatsci.2007.03.003>.
- Essid, I., Medini, M., & Hassouna, M. (2009). Technological and safety properties of *Lactobacillus plantarum* strains isolated from a Tunisian traditional salted meat. *Meat Science*, 81, 203-208. <https://doi.org/10.1016/j.meatsci.2008.07.020>.
- Eruteya, O., & Eze, C. (2017). Proximate Composition and Antibiotic Resistant *Listeria* Species in Fura-de-nunu Sold in Port Harcourt. *Journal of Advances in Biology & Biotechnology*, 11, 1-7. <https://doi.org/10.9734/jabb%2F2017%2F30655>
- Ezeokoli, O.T., Adeleke, R.A., & Bezuidenhout, C.C. (2018). Core bacterial community of soy-daddawa: Insights from high-throughput DNA metabarcoding. *LWT*, 97, 61-66, <https://doi.org/10.1016/j.lwt.2018.06.039>.
- Fall, N. G., Tounkara, L. S., Diop, M. B., Thiaw, O. T., & Thonart, P. (2017). Chemical characteristics and microbial quality of Guedj a traditional fermented fish from Senegal. *International Journal of Sciences*, 6, 48-54. <https://doi.org/10.18483/IJSCI.1323>.
- Fasoyiro, B.S., Obatolu, V.A., Ashaye, A.O., Adegoke, G.A. & Farinde, O.E. (2010). Microbial hazards and critical control points of locally processed soy-cheese in Nigeria. *Nutrition & Food Science*, 40, 591-597. <https://doi.org/10.1108/00346651011090392>
- Food and Agriculture Organization of the United Nations. (2010). Current status and options for biotechnologies in food processing and in food safety in developing countries. FAO International Technical Conference report. FAO, Guadalajara, Mexico. Report No: ABCD-10/7.1
- Food and Agriculture Organization of the United Nations. (2020a). FAOSTAT Database. Rome, Italy: FAO. Retrieved December 30, 2020 from <http://faostat3.fao.org/home/E>
- FAO, IFAD, UNICEF, WFP and WHO. (2020b). The State of Food Security and Nutrition in the World 2020. Transforming food systems for affordable healthy diets. Rome, FAO.
- FAO/WHO (1995). Codex Standard for Edible Cassava Flour. Codex Standard 176-1989, Food and Agriculture Organisation and World Health Organisation of the United Nations, Rome.
- Ferreira, A.M., & Mendes-Faia, A. (2020) The Role of Yeasts and Lactic Acid Bacteria on the Metabolism of Organic Acids during Winemaking. *Foods*, 9, 1231; doi:10.3390/foods9091231

- Founou, L.L., Founou, R.C., & Essack, S.Y. (2016). Antibiotic Resistance in the Food Chain: A Developing Country-Perspective. *Frontiers in Microbiology*, 7,1881. <https://doi.org/10.3389/fmicb.2016.01881>
- Fowoyo, P.T., & Ogunbanwo, S.T. (2017). Antimicrobial resistance in coagulase-negative staphylococci from Nigerian traditional fermented foods. *Annals of Clinical Microbiology & Antimicrobials*, 16, 4. <https://doi.org/10.1186/s12941-017-0181-5>.
- Franz, C.M., Huch, M., Mathara, J.M., Abriouel, H., Benomar, N., Reid, G., Galvez, A., & Holzapfel, W.H. (2014). African fermented foods and probiotics. *International Journal of Food Microbiology*, 190, 84-96. <https://doi.org/10.1016/j.ijfoodmicro.2014.08.033>.
- Gagaoua, M., & Boudechicha, H-R. (2018). Ethnic meat products of the North African and Mediterranean countries: An overview. *Journal of Ethnic Foods*, 5, 83-98. <https://doi.org/10.1016/j.jef.2018.02.004>.
- Gänzle, M. (2020). Food fermentations for improved digestibility of plant foods – an essential ex-situ digestion step in agricultural societies? *Current Opinion in Food Science*, 32,124-132. <https://doi.org/10.1016/j.cofs.2020.04.002>.
- Glover, R.L.K., Abaidoo, R.C., Jakobsen, M., Jespersen, L. (2005) Biodiversity of *Saccharomyces cerevisiae* isolated from a survey of pito production sites in various parts of Ghana. *Systematic and Applied Microbiology*, 28, 755-761. <https://doi.org/10.1016/j.syapm.2005.05.003>.
- Gran, H.M., Mutukumira, A.N., Wetlesen, A., & Narvhus, J.A. (2002). Smallholder dairy processing in Zimbabwe: the production of fermented milk products with particular emphasis on sanitation and microbiological quality. *Food Control*, 13, 161-168. [https://doi.org/10.1016/S0956-7135\(01\)00094-9](https://doi.org/10.1016/S0956-7135(01)00094-9).
- Greppi, A., Rantsiou, K., Padonou, W., Hounhouigan, J., Jespersen, L., Jakobsen, M., & Cocolin, L. (2013). Determination of yeast diversity in ogi, mawè, gowé and tchoukoutou by using culture-dependent and -independent methods. *International Journal of Food Microbiology*, 165, 84-88. <https://doi.org/10.1016/j.ijfoodmicro.2013.05.005>.
- Hanchi, H., Mottawea, W., Sebei, K., & Hammami, R. (2018). The Genus *Enterococcus*: Between Probiotic Potential and Safety Concerns-An Update. *Frontiers in microbiology*, 9, 1791. <https://doi.org/10.3389/fmicb.2018.01791>
- Houngbédji, M., Johansen, P., Padonou, S.W., Akissoé, N., Arneborg, N., Nielsen, D.S., Hounhouigan, D.J., & Jespersen, L. (2018). Occurrence of lactic acid bacteria and yeasts at species and strain level during spontaneous fermentation of mawè, a cereal dough produced in West Africa. *Food Microbiology*, 76, 267-278. <https://doi.org/10.1016/j.fm.2018.06.005>.
- Jans, C., Meile, L., Kaindi, D.M.W., Kogi-Makua, W., Lamuka, P., Renault, P., Kreikemever, B., Lacroix, C., Hattendorf, J., Zinsstag, J., Schelling, E., Fokou, G., & Bonfoh, B. (2017). African fermented dairy products – Overview of predominant technologically important microorganisms focusing on African *Streptococcus infantarius* variants and potential future applications for enhanced food safety and security. *International Journal of Food Microbiology*, 250, 27 – 36. <https://doi.org/10.1016/j.ijfoodmicro.2017.03.012>.

- Johansen, P. G., Owusu-Kwarteng, J., Parkouda, C., Padonou, S. W., & Jespersen, L. (2019). Occurrence and Importance of Yeasts in Indigenous Fermented Food and Beverages Produced in Sub-Saharan Africa. *Frontiers in microbiology*, 10, 1789. <https://doi.org/10.3389/fmicb.2019.01789>.
- Kafetzopoulos, D.P., Psomas, E.L., Kafetzopoulos, P.D. (2013). Measuring the effectiveness of the HACCP food safety management system. *Food Control*, 33, 505-513.
- Karamoko, D., Djeni, N.T., N'guessan, K.F., Bouatenin, K.M.J.-P., & Dje, K.M. (2012). The biochemical and microbiological quality of palm wine samples produced at different periods during tapping and changes which occurred during their storage. *Food Control*, 26, 504-511. <https://doi.org/10.1016/j.foodcont.2012.02.018>.
- Kergourlay, G., Taminiau, B., Daube, G., & Champomier Vergès, M.C. (2015). Metagenomic insights into the dynamics of microbial communities in food. *International Journal of Food Microbiology*, 213, 31-39. <https://doi.org/10.1016/j.ijfoodmicro.2015.09.010>.
- Kimaryo, V.M., Massawe, G.A., Olasupo, N.A. and Holzapfel, W.H. (2000) The use of starter cultures in the fermentation of cassava for the production of kivunde, a traditional Tanzanian food product. *International Journal of Food Microbiology*, 56, 179-190.
- Kouamé, A.K., Djéni, T.N., N'guessan, F.K., & Dje, M.K. (2013). Postprocessing microflora of commercial attiéke (a fermented cassava product) produced in the south of Côte d'Ivoire. *Letters in Applied Microbiology*, 56, 44-50. <https://doi.org/10.1111/lam.12014>.
- Kunene, N.F., Hastings, J.W., & von Holy, A. (1999). Bacterial populations associated with a sorghum-based fermented weaning cereal. *International Journal of Food Microbiology*, 49, 75-83. [https://doi.org/10.1016/S0168-1605\(99\)00062-8](https://doi.org/10.1016/S0168-1605(99)00062-8).
- Lateef, A., & Ojo, M.O. (2015). Public health issues in the processing of cassava (*Manihot esculenta*) for the production of lafun and the application of hazard analysis control measures. *Quality Assurance and Safety of Crops & Foods*, 8, 165 – 177. <https://doi.org/10.3920/QAS2014.0476>.
- Lei, V., & Jakobsen, M. (2004). Microbiological characterization and probiotic potential of koko and koko sour water, African spontaneously fermented millet porridge and drink. *Journal of Applied Microbiology*, 96, 384-97. <https://doi.org/10.1046/j.1365-2672.2004.02162.x>.
- Lyumugabe, F., Kamaliza, G., Bajyana, E., & Thonart, P.H. (2010). Microbiological and physico-chemical characteristic of Rwandese traditional beer "Ikigage". *African Journal of Biotechnology*, 9, 4241 – 4246.
- Markowiak, P., & Śliżewska, K. (2017). Effects of Probiotics, Prebiotics, and Synbiotics on Human Health. *Nutrients*, 9, 1021. <https://doi.org/10.3390/nu9091021>
- McEwen, S. & Collignon, P. (2018). Antimicrobial Resistance: A One Health Perspective. *Microbiology Spectrum*, 6, ARBA-0009-2017. <https://doi.org/10.1128/microbiolspec.ARBA-0009-2017>.
- Miambi, E., Guyot, J.P., & Ampe, F. (2003). Identification, isolation and quantification of representative bacteria from fermented cassava dough using an integrated approach of culture-dependent and culture-independent methods. *International Journal of Food Microbiology*, 82, 111-120. [https://doi.org/10.1016/S0168-1605\(02\)00256-8](https://doi.org/10.1016/S0168-1605(02)00256-8).

- Misihairabgwi, J., & Cheikhyoussef, A. (2017). Traditional fermented foods and beverages of Namibia. *Journal of Ethnic Foods*, 4, 145 – 153. <https://doi.org/10.1016/j.jef.2017.08.001>.
- Mokoena, M.P., Mutanda, T., & Olaniran, A.O. (2016). Perspectives on the probiotic potential of lactic acid bacteria from African traditional fermented foods and beverages. *Food and Nutrition Research*, 60, <https://doi.org/10.3402/fnr.v60.29630>.
- Moodley, S.S., Dlamini, N.R., Steenkamp, L., Buys, E.M. (2019). Bacteria and yeast isolation and characterisation from a South African fermented beverage. *South African Journal of Science*, 115, <https://doi.org/10.17159/sajs.2019/5996>
- Moonga, H. B., Schoustra, S. E., van den Heuvel J., Linnemann, A. R., Samad, Md. S., Shindano, J., Smid, J. (2020) Composition and Diversity of Natural Bacterial Communities in Mabisi, a Traditionally Fermented Milk. *Frontiers in Microbiology*, 11, 1816, <https://doi.org/10.3389/fmicb.2020.01816>.
- Mourad, G., & Bettache, G. (2015). Characterization and identification of lactic acid bacteria isolated from traditional cheese (Klila) prepared from cow's milk. *African Journal of Microbiology Research*, 9, 71-77 DOI: <https://doi.org/10.5897/AJMR2014.7279>
- Mpofu, A., Linnemann, A.R., Nout, M.J.R., Zwietering, M.H., Smid, E.J., den Besten, H.M.W. (2016). Inactivation of bacterial pathogens in yoba mutandabota, a dairy product fermented with the probiotic *Lactobacillus rhamnosus* yoba. *International Journal of Food Microbiology*, 217, 42-48. <https://doi.org/10.1016/j.ijfoodmicro.2015.09.016>.
- Mugula, J.K., Nnko, S.A., Narvhus, J.A., & Sørhaug, T. (2003) Microbiological and fermentation characteristics of togwa, a Tanzanian fermented food. *International Journal of Food Microbiology*, 80,187-99. [https://doi.org/10.1016/S0168-1605\(02\)00141-1](https://doi.org/10.1016/S0168-1605(02)00141-1).
- Mukisa, I.M., Byakika, S., Meeme, R., Wacoo, A.P., Sybesma, W. & Kort, R. (2019). Adopting traditional fermented foods as carriers for probiotics: The case of Obushera and *Lactobacillus rhamnosus* yoba. *Nutrition & Food Science*, 50, 841-852. <https://doi.org/10.1108/NFS-06-2019-0188>
- Mukisa, I., Ntate, D. & Byakiya, S. (2016). Application of starter cultures in the production of Enturire - a traditional sorghum-based alcoholic beverage. *Food Science & Nutrition*, 5, 609 – 616. <https://doi.org/10.1002/fsn3.438>
- Mukisa, I.M., Porcellato, D., Byaruhanga, Y.B., Muyanja, C.M., Rudi, K., Langsrud, T., & Narvhus, J.A. (2012). The dominant microbial community associated with fermentation of Obushera (sorghum and millet beverages) determined by culture-dependent and culture-independent methods. *International Journal of Food Microbiology*, 160, 1-10. <https://doi.org/10.1016/j.ijfoodmicro.2012.09.023>.
- Mukisa, I.M., Ssendagala, G.W. & Byakika, S. (2020). Microbiological safety and physicochemical composition of Bongo, a traditional fermented milk product from Lyantonde district, Uganda. *Scientific African*, 10, e00583, <https://doi.org/10.1016/j.sciaf.2020.e00583>.
- Murtala, M., Alabi, O.S., & Ayeni, F.A. (2018). Susceptibility Profiles of Lactic and Acetic Acid Bacteria Isolated from Ogi, Yoghurt and Human Vagina in Nigeria. *Nigerian Journal of Pharmaceutical Sciences*, 17.

- Nhung, N.T., Chansiripornchai, N., & Carrique-Mas, J.J. (2017). Antimicrobial Resistance in Bacterial Poultry Pathogens: A Review. *Frontiers in Veterinary Science*, 4:126. <https://doi.org/10.3389/fvets.2017.00126>
- Nieminen, M. T., Novak-Frazer, L., Collins, R., Dawsey, S. P., Dawsey, S. M., Abnet, C. C., White, R. E., Freedman, N. D., Mwachiro, M., Bowyer, P., Salaspuro, M., & Rautemaa, R. (2013). Alcohol and acetaldehyde in African fermented milk mursik-a possible etiologic factor for high incidence of esophageal cancer in western Kenya. *Cancer Epidemiology, Biomarkers & Prevention*, 22, 69–75. <https://doi.org/10.1158/1055-9965.EPI-12-0908>.
- Nwaiwu, O., Aduba, C.C., Igbokwe, V.C., Sam, C.E., & Ukwuru, M.U. (2020). Traditional and artisanal beverages in Nigeria: Microbial diversity and safety issues. *Beverages*, 6, 1 - 22. <https://doi.org/10.3390/beverages6030053>.
- Nwaiwu, O., & Itumoh, M. (2017). Chemical contaminants associated with palm wine from Nigeria are potential food safety hazards. *Beverages*, 3, 1-12. <https://doi.org/10.3390/beverages3010016>.
- Nyambane, B., Thari, W.M., Wangoh, J., & Njage, P.M. (2014). Lactic acid bacteria and yeasts involved in the fermentation of amabere amaruranu, a Kenyan fermented milk. *Food Science & Nutrition*, 2, 692–699. <https://doi.org/10.1002/fsn3.162>.
- Nyatoti, V.M., Mtero, S.S. & Rukure, G. (1997). Pathogenic *Escherichia coli* in traditional African weaning foods. *Food Control*, 8, 51-54. [https://doi.org/10.1016/S0956-7135\(96\)00068-0](https://doi.org/10.1016/S0956-7135(96)00068-0).
- Obadina, A.O., Oyewole, O.B., Sanni, L.O., Tomlins, K.I., & Westby, A. (2008). Identification of hazards and critical control points (CCP) for cassava fufu processing in South-West Nigeria. *Food Control*, 19, 22-26. <https://doi.org/10.1016/j.foodcont.2007.01.002>.
- Obilie, E.B., Tano-Debrah, K., Amoa-Awua, W.K. (2004). Souring and breakdown of cyanogenic glucosides during the processing of cassava into akyeke. *International Journal of Food Microbiology*, 93, 115-121, <https://doi.org/10.1016/j.ijfoodmicro.2003.11.006>.
- Obioha, P.I., Ouoba, L.I.I., Anyogu, A., Awamaria, B., Atchia, S., Ojimekwe, P.C., Sutherland, J.P. & Ghoddusi, H.B. (2021). Identification and characterisation of the lactic acid bacteria associated with the traditional fermentation of a dairy fermented product. *Brazilian Journal of Microbiology*, <https://doi.org/10.1007/s42770-021-00461-y>
- Odom, T.C., Udensi, E.A., & Nwanekezi, E.C. (2012) *Microbiological Qualities of Hawked Retted Cassava Fufu in Aba Metropolis of Abia State*, *Nigerian Food Journal*, 30, 53 – 58. [https://doi.org/10.1016/S0189-7241\(15\)30013-8](https://doi.org/10.1016/S0189-7241(15)30013-8).
- Odunfa, S.A., & Oyewole, O.B. (1998). African fermented foods. In B.J.B. Wood (Ed). *Microbiology of Fermented Foods*, Vol. 2. (pp. 713–752). Blackie Academic and Professional, London.
- OECD/FAO (2016). *OECD-FAO Agricultural Outlook 2016-2025*, OECD Publishing, Paris. [http://dx.doi.org/10.1787/agr\\_outlook-2016-en](http://dx.doi.org/10.1787/agr_outlook-2016-en).
- Oguntoyinbo, F.A. (2012). Development of Hazard Analysis Critical Control Points (HACCP) and Enhancement of Microbial Safety Quality during Production of Fermented Legume Based Condiments in Nigeria. *Nigerian Food Journal*, 30, 59-66. [https://doi.org/10.1016/S0189-7241\(15\)30014-X](https://doi.org/10.1016/S0189-7241(15)30014-X).

- Oguntinyinbo, F.A. (2014) Safety challenges associated with traditional foods of West Africa. *Food Reviews International*, 30, 338 – 348. <https://doi.org/10.1080/87559129.2014.940086>
- Oguntinyinbo, F.A., Huch, M., Cho, G.S., Schillinger, U., Holzapfel, W.H., Sanni, A.I., & Franz, C.M. (2010). Diversity of bacillus species isolated from okpehe, a traditional fermented soup condiment from Nigeria. *Journal of Food Protection*, 73, 870-8. <https://doi.org/10.4315/0362-028X-73.5.870>.
- Oguntinyinbo, F.A., & Narbad, A. (2015). Multifunctional properties of *Lactobacillus plantarum* strains isolated from fermented cereal foods. *Journal of Functional Foods*, 17, 621-631. <https://doi.org/10.1016/j.jff.2015.06.022>.
- Oguntinyinbo, F.A., & Okueso, O. (2013). Prevalence, distribution, and antibiotic resistance pattern among enterococci species in two traditional fermented dairy foods. *Annals of Microbiology*, 63, 755–761. <https://doi.org/10.1007/s13213-012-0530-8>
- Oguntinyinbo, F.A., Tourlomousis, P., Gasson, M.J., & Narbad, A. (2011). Analysis of bacterial communities of traditional fermented West African cereal foods using culture independent methods. *International Journal of Food Microbiology*, 145, 205-210. <https://doi.org/10.1016/j.ijfoodmicro.2010.12.025>
- Okafor, N. (1992). Applications of Biotechnology to Fermented Foods: Report of an Ad Hoc Panel of the Board on Science and Technology for International Development. National Research Council (US) Panel on the Applications of Biotechnology to Traditional Fermented Foods. Washington (DC): National Academies Press, USA.
- Okolie, P.I., Opara, C.N., Emerenini, E.C., & Uzochukwu, S.V.A. (2013). Evaluation of Bacterial Diversity in Palm Wine by 16S rDNA Analysis of Community DNA. *Nigerian Food Journal*, 31, 83-90. [https://doi.org/10.1016/S0189-7241\(15\)30060-6](https://doi.org/10.1016/S0189-7241(15)30060-6).
- Oladipo, I.A., Sanni, A., Swarnakar, S. (2013). Phenotypic and Genomic Characterization of Enterococcus Species from Some Nigerian Fermented Foods. *Food Biotechnology*, 27, 39 - 53. <https://doi.org/10.1080/08905436.2012.755627>
- Olasupo, N.A., Odunfa, S.A., & Obayori, O.S. (2010). Ethnic African fermented foods. In J.P. Tamang, & K. Kailasapathy (Eds.), *Fermented foods and beverages of the world* (pp. 323 – 352). CRC Press, London.
- Olasupo, N.A., Okorie, C.P., & Oguntinyinbo, F.A. (2016). The Biotechnology of Ugba, a Nigerian Traditional Fermented Food Condiment. *Frontiers in Microbiology*, 7,1153. <https://doi.org/10.3389/fmicb.2016.01153>
- Oloso, N.O., Fagbo, S., Garbati, M., Olonitola, S.O., Awosanya, E.J., Aworh, M.K., Adamu, H., Odetokun, I.A., & Fasina, F.O. (2018). Antimicrobial Resistance in Food Animals and the Environment in Nigeria: A Review. *International Journal of Environmental Research & Public Health*,15,1284. <https://doi.org/10.3390/ijerph15061284>
- Omafuvbe, B. O., Adigun, A. R., Ogunsuyi, J. L., & Asunmo, A. M. (2007). Microbial diversity in ready-to-eat fufu and Lafun-fermented cassava products sold in Ile-Ife, Nigeria. *Research Journal of Microbiology*, 2, 831-837. <https://doi.org/10.3923/jm.2007.831.837>.

- Omafuvbe, B.O., Shonukan, O.O., & Abiose, S.H. (2000). Microbiological and biochemical changes in the traditional fermentation of soybean for 'soy-daddawa' — Nigerian food condiment. *Food Microbiology*, 17, 5, 469-474, <https://doi.org/10.1006/fmic.1999.0332>.
- Omarak, R.A., Hinenoya, A., Awasthi, S.P., Iguchi, A., Shima, A., Elbagory, A-R.M., & Yamasaki, S. (2016). Prevalence and pathogenic potential of *Escherichia coli* isolates from raw milk and raw milk cheese in Egypt. *International Journal of Food Microbiology*, 221, 69-76. <https://doi.org/10.1016/j.ijfoodmicro.2016.01.009>.
- Omemu, A.M., Obadina, A.O., Taiwo, G.J., & Obuotor, T.M. (2014). Microbiological assessment and prevalence of foodborne pathogens in street vended wara — Nigerian white cheese. *American Journal of Food and Nutrition* 2, 59 – 62. <https://doi.org/10.12691/ajfn-2-4-2>.
- O'Neill, J. (2016). Tackling drug-resistant infections globally: final reports and recommendations. The Review on Antimicrobial Resistance. Available from: [https://amr-review.org/sites/default/files/160525\\_Final%20paper\\_with%20cover.pdf](https://amr-review.org/sites/default/files/160525_Final%20paper_with%20cover.pdf). Accessed on 10.11.2020.
- Onyenekwe, P.C., Erhabor, G.O., & Akande, S.A. (2015). Characterisation of aroma volatiles of indigenous alcoholic beverages: burukutu and pito. *Natural Product Research*, 30, 558 – 564. <https://doi.org/10.12691/ajfn-2-4-210.1080/14786419.2015.1032279>.
- Oranusi, S., Okereke, J., Braide, W., & Okorundu, S.I. (2015). Microbiological and chemical compositions of agbarati and ogiri Igbo, popular foods of south eastern Nigerians. *Journal of Chemical, Biological and Physical Sciences*, 5,
- Osvik, R.D, Sperstad, S., Breines, E.M., Hareide, E., Godfroid, J., Zhou, Z., Ren, P., Geoghegan, C., Holzapfel, W., & Ringø, E. (2013). Bacterial diversity of aMasi, a South African fermented milk product, determined by clone library and denaturing gradient gel electrophoresis analysis. *African Journal of Microbiology Research*, 7, 4146 – 4158. <https://doi.org/10.5897/AJMR12.2317>.
- Ouoba, L.I.I., Kando, C., Parkouda, C., Sawadogo-Lingani, H., Diawara, B., & Sutherland, J.P. (2012). The microbiology of Bandji, palm wine of *Borassus akeassii* from Burkina Faso: identification and genotypic diversity of yeasts, lactic acid and acetic acid bacteria. *Journal of Applied Microbiology*, 113, 428-41. <https://doi.org/10.1111/jam.12014>.
- Ouoba, L. L. I., Diawara, B., Amoa-Awua, W. K., Traore, A. S., & Moller, P. L. (2004). Genotyping of starter cultures of *Bacillus subtilis* and *Bacillus pumilus* for fermentation of African locust bean *Parkia biglobosa*, to produce soumbala. *International Journal of Food Microbiology*, 90, 197–205. [https://doi.org/10.1016/S0168-1605\(03\)00302-7](https://doi.org/10.1016/S0168-1605(03)00302-7)
- Ouoba, L.I.I, Lei, V., & Jensen, L.B. (2008a). Resistance of potential probiotic lactic acid bacteria and bifidobacteria of African and European origin to antimicrobials: determination and transferability of the resistance genes to other bacteria. *International Journal of Food Microbiology*, 121, 217-24. <https://doi.org/10.1016/j.ijfoodmicro.2007.11.018>.
- Ouoba, L.I.I., Nielsen, D.S., Anyogu, A., Kando, C., Diawara, B., Jespersen, L. & Sutherland, J.P. (2015a). *Hanseniaspora jakobsenii* sp. nov., a novel yeast isolated from Bandji, a traditional palm wine of *Borassus akeassii* from Burkina Faso. *International Journal of Systematic and Evolutionary Microbiology*, 65, 3576-3579. <https://doi.org/10.1099/ijsem.0.000461>.

- Ouoba, L.I.I., Thorsen, L., & Varnam, A.H. (2008b). Enterotoxins and emetic toxins production by *Bacillus cereus* and other species of *Bacillus* isolated from Soumbala and Bikalga, African alkaline fermented food condiments. *International Journal of Food Microbiology*, 124, 224-230. <https://doi.org/10.1016/j.ijfoodmicro.2008.03.026>.
- Ouoba, L.I.I., Voudibo-Mbozo, A., Anyogu, A., Obioha, P., Sutherland, J.P., Jespersen, L., & Ghoddusi, H.B. (2019). Environmental heterogeneity of *Staphylococcus* species from alkaline fermented foods and associated toxins and antimicrobial resistance genetic elements. *International Journal of Food Microbiology*, 311, 108356. <https://doi.org/10.1016/j.ijfoodmicro.2019.108356>.
- Ouoba, L.I.I., Voudibo-Mbozo, A., Thorsen, L., Anyogu, A., Nielsen, D.S., Kobawila, C., & Sutherland, J.P. (2015b). *Lysinibacillus louembis* sp. nov., a novel bacterium isolated from Ntoba Mbodi, alkaline fermented leaves of cassava from Congo-Brazzaville. *International Journal of Systematic and Evolutionary Microbiology*, 65, 4256-4262. <https://doi.org/10.1099/ijsem.0.000570>.
- Owusu-Kwarteng, J., Akabanda, F., Nielsen, D.S., Tano-Debrah, K., Glover, R.L., & Jespersen, L. (2012). Identification of lactic acid bacteria isolated during traditional fura processing in Ghana. *Food Microbiology*, 32, 72-78. <https://doi.org/10.1016/j.fm.2012.04.010>.
- Owusu-Kwarteng, J., Wuni, A., Akabanda, F., & Jespersen, L. (2018). Prevalence and Characteristics of *Listeria monocytogenes* Isolates in Raw Milk, Heated Milk and Nunu, a Spontaneously Fermented Milk Beverage, in Ghana. *Beverages*, 4, 40. <https://doi.org/10.3390/beverages4020040>
- Owusu-Kwarteng, J., Wuni, A., Akabanda, F., Tano-Debrah, K., & Jespersen, L. (2017). Prevalence, virulence factor genes and antibiotic resistance of *Bacillus cereus* sensu lato isolated from dairy farms and traditional dairy products. *BMC microbiology*, 17, 65. <https://doi.org/10.1186/s12866-017-0975-9>
- Oyewole, O. B., & Odunfa, S. A. (1988). Microbiological studies on cassava fermentation for 'lafun' production. *Food Microbiology*, 5, 125-133. [https://doi.org/10.1016/0740-0020\(88\)90010-X](https://doi.org/10.1016/0740-0020(88)90010-X).
- Padonou, S.W., Nielsen, D.S., Hounhouigan, J.D., Thorsen, L., Nago, M.C., & Jakobsen, M. (2009). The microbiota of lafun, an African traditional cassava food. *International Journal of Food Microbiology*, 133, 22 – 30. <https://doi.org/10.1016/j.ijfoodmicro.2009.04.019>.
- Parker, M., Zobrist, S., Donahue, C., Edick, C., Mansen, K., Hassan, Zade., Nadjari, M., Heerikhuisen, M., Sybesma, W., Molenaar, D., Diallo, A.M., Milani, P., & Kort, R. (2018). Naturally Fermented Milk From Northern Senegal: Bacterial Community Composition and Probiotic Enrichment With *Lactobacillus rhamnosus*. *Frontiers in Microbiology*, 9:2218. <https://doi.org/10.3389/fmicb.2018.02218>.
- Parkouda, C., Nielsen, D.S, Azokpota, P., Ouoba, L.I., Amoa-Awua, W.K., Thorsen, L., Hounhouigan, J.D., Jensen, J.S., Tano-Debrah, K., Diawara, B. & Jakobsen, M. (2009). The microbiology of alkaline-fermentation of indigenous seeds used as food condiments in Africa and Asia. *Critical Reviews in Microbiology*, 35,139-56. <https://doi.org/10.1080/10408410902793056>.
- Parkouda, C., Thorsen, L., Compaoré, C.S., Nielsen, D.S., Tano-Debrah, K., Jensen, J.S., Diawara, B., & Jakobsen, M. (2010). Microorganisms associated with Maari, a Baobab seed fermented product. *International Journal of Food Microbiology*, 142, 292-301. <https://doi.org/10.1016/j.ijfoodmicro.2010.07.004>

- Paudyal, N., Anihonyi, V., Hounhouigan, J., Matsheka, M.I., Sekwati-Monang, B., Amoa-Awua, W., Atter, A., Ackah, N.B., Mbugua, S., Asagbra, A., Abdelgardi, W., Nakavuma, J., Jakobsen, M., & Fang, W. (2017). Prevalence of foodborne pathogens in food from selected African countries – a meta-analysis. *International Journal of Food Microbiology*, 249, 35 – 43. <https://doi.org/10.1016/j.ijfoodmicro.2017.03.002>.
- Rehaïem, A., Fhoula, I., Slim, A.F., Boubaker, I.B.B., Chihi, A.B., Ouzari, H.I. (2016). Prevalence, acquired antibiotic resistance and bacteriocin production of *Enterococcus* spp. isolated from tunisian fermented food products. *Food Control*, 63, 259-266. <https://doi.org/10.1016/j.foodcont.2015.11.034>.
- Reygaert, W. C. (2018). An overview of the antimicrobial resistance mechanisms of bacteria. *AIMS microbiology*, 4, 482–501. <https://doi.org/10.3934/microbiol.2018.3.482>
- Rolle, R., & Satin, M. (2002). Basic requirements for the transfer of fermentation technologies to developing countries. *International Journal of Food Microbiology*, 75, 181 – 187. [https://doi.org/10.1016/S0168-1605\(01\)00705-X](https://doi.org/10.1016/S0168-1605(01)00705-X).
- Ropkins, K., & Beck, A.J. (2000). Evaluation of worldwide approaches to the use of HACCP to control food safety. *Trends in Food Science & Technology*, 11, 10-21, [https://doi.org/10.1016/S0924-2244\(00\)00036-4](https://doi.org/10.1016/S0924-2244(00)00036-4).
- Sadek, Z.I., Fathi, F.A., & Salem, M.M.E. (2006). Incidence, survival and biocontrol of psychrotrophic *Bacillus cereus* and its potential for toxin production in milk and tallaga cheese. *Polish Journal of Food and Science Technology*, 56, 419-425.
- Samet-Bali, O., Felfoul, I., Lajnaf, R., Attia, H., & Ayadi, M.A. (2016). Hygienic quality of “rayeb”, a traditional Tunisian fermented cow’s milk. *International Food Research Journal*, 23, 366 – 369.
- Sanni, A.I., Asiedu, M. & Ayernor, G.S. (2002). Microflora and Chemical Composition of Momoni, a Ghanaian Fermented Fish Condiment. *Journal of Food Composition and Analysis*, 15, 577 – 583. <https://doi.org/10.1006/jfca.2002.1063>.
- Sanni, A.I., & Lönnér, C. (1993). Identification of yeasts isolated from Nigerian traditional alcoholic beverages. *Food Microbiology*, 10, 517-523. <https://doi.org/10.1006/fmic.1993.1057>.
- Sawadogo-Lingani, H., Lei, V., Diawara, B., Nielsen, D.S., Møller, P.L., Traoré, A.S., & Jakobsen, M. (2007). The biodiversity of predominant lactic acid bacteria in dolo and pito wort for the production of sorghum beer. *Journal of Applied Microbiology*, 103, 765-77. <https://doi.org/10.1111/j.1365-2672.2007.03306.x>.
- Schoder, D., Maichin, A., Lema, B., & Laffa, J. (2013). Microbiological Quality of Milk in Tanzania: From Maasai Stable to African Consumer Table. *Journal of Food Protection*, 76, 1908–1915. <https://doi.org/10.4315/0362-028X.JFP-13-101>.
- Schoustra, S.E., Kasase, C., Toarta, C., Kassen, R., Poulain, A.J. (2013). Microbial community structure of three traditional zambian fermented products: mabisi, chibwantu and munkoyo. *PLoS One*, 8, e63948. <https://doi.org/10.1371/journal.pone.0063948>.

- Setta, M.C., Matemu, A., & Mbega, E.R. (2020) Potential of probiotics from fermented cereal-based beverages in improving health of poor people in Africa. *Journal of Food Science & Technology*, 57, 3935-3946. <https://doi.org/10.1007/s13197-020-04432-3>.
- Sivamaruthi, B.S., Kesika, P., & Chaiyasut, C. (2019). Toxins in Fermented Foods: Prevalence and Preventions—A Mini Review. *Toxins*, 11, 4. <https://doi.org/10.3390/toxins11010004>.
- Soni, S., & Dey, G. (2014). Perspectives on global fermented foods. *British Food Journal*, 116, 1767-1787. <https://doi.org/10.1108/BFJ-01-2014-0032>.
- Soro-Yao, A.A., Brou, K., Amani, G., Thonart, P., & Dje, K.M. (2014). The use of lactic acid bacteria starter cultures during the processing of fermented cereal-based foods in West Africa: A Review. *Tropical Life Sciences Research*, 25, 81 – 100.
- Spellberg, B. (2014). The future of antibiotics. *Critical Care*, 18, 228. <https://doi.org/10.1186/cc13948>.
- Sudi, I.Y., De, Nandita, De, Dunkrah, U. A. (2011). Nigerian indigenous yoghurt (kindirmo) production using *Lactobacillus bulgaricus* and *Streptococcus thermophilus* mutants as starter culture. *African Journal of Biotechnology*, 10, 1651-1654. <https://doi.org/10.5897/AJB10.1829>.
- Suri DJ, Tano-Debrah K, Ghosh SA. (2014). Optimization of the nutrient content and protein quality of cereal legume blends for use as complementary foods in Ghana. *Food and Nutrition Bulletin*, 35, 372-381.
- Szyniszewska, A.M. (2020). CassavaMap, a fine-resolution disaggregation of cassava production and harvested area in Africa in 2014. *Scientific Data*, 7, 159, <https://doi.org/10.1038/s41597-020-0501-z>
- Tadesse, B.T., Abera, A.B., Tefera, A.T., Muleta, D., Alemu, Z.T., & Wessel, G. (2019). Molecular Characterization of Fermenting Yeast Species from Fermented Teff Dough during Preparation of Injera Using ITS DNA Sequence. *International Journal of Food Science*, Article ID 1291863, <https://doi.org/10.1155/2019/1291863>.
- Tadesse, B.T., Ashley, E.A., Ongarello, S., Havumaki, J., Wijegoonewardena, M., González, I.J., & Dittrich, S. (2017). Antimicrobial resistance in Africa: a systematic review. *BMC Infectious Diseases*, 17, 616. <https://doi.org/10.1186/s12879-017-2713-1>.
- Tamang, J.T., Watanabe, K. & Holzapfel, W. (2016) Review: Diversity of microorganisms in global fermented foods and beverages. *Frontiers in Microbiology*, <https://doi.org/10.3389/fmicb.2016.00377>.
- Tano-Debrah, K., Saalia, F. K., Ghosh, S., & Hara, M. (2019). Development and Sensory Shelf-Life Testing of KOKO Plus: A Food Supplement for Improving the Nutritional Profiles of Traditional Complementary Foods. *Food and Nutrition Bulletin*, 40, 340–356. <https://doi.org/10.1177/0379572119848290>
- Thorsen, L., Kando, C.K., Sawadogo, H., Larsen, N., Diawara, B., Ouédraogo, G.A., Hendriksen, N.B., & Jespersen, L. (2015). Characteristics and phylogeny of *Bacillus cereus* strains isolated from Maari, a traditional West African food condiment. *International Journal of Food Microbiology*, 196, 70-80. <https://doi.org/10.1016/j.ijfoodmicro.2014.11.026>.

- Usman, U. B., Kwaga, J. K. P., KABIR, J., & Olonitola, O.S. (2016). Isolation and Antimicrobial Susceptibility of *Listeria monocytogenes* from Raw Milk and Milk Products in Northern Kaduna State, Nigeria. *Journal of Applied & Environmental Microbiology*, 4, 46-54. <https://doi.org/10.12691/JAEM-4-3-1>.
- United Nations. (2020). Population. Available from: <https://www.un.org/en/sections/issues-depth/population/> Accessed 28.12.2020
- Van, T.T.H., Yidana, Z., Smooker, P.M., & Coloe, P.J. (2020). Antibiotic use in food animals worldwide, with a focus on Africa: Pluses and minuses. *Journal of Global Antimicrobial Resistance*, 20, 170-177. <https://doi.org/10.1016/j.jgar.2019.07.031>
- Valenzuela, A.S., Omar, N.B., Abriouel, H., Lopez, R.L., Ortega, E., Canamero, M.M., & Gálvez, A. (2008). Risk factors in enterococci isolated from foods in Morocco: determination of antimicrobial resistance and incidence of virulence traits. *Food and Chemical Toxicology*, 46, 2648-2652. <https://doi.org/10.1016/j.fct.2008.04.021>.
- Van Boeckel, T.P., Gandra, S., Ashok, A., Caudronm Q., Grenfell, B.T., & Levin, S.A. (2014). Global antibiotic consumption 2000 to 2010: an analysis of national pharmaceutical sales data. *Lancet Infectious Diseases*, 14, 742 – 750.
- Vieira-Dalodé, G., Jespersen, L., Hounhouigan, J., Moller, P.L., Nago, C.M., & Jakobsen, M. (2007). Lactic acid bacteria and yeasts associated with gowé production from sorghum in Bénin. *Journal of Applied Microbiology*, 103, 342-349. <https://doi.org/10.1111/j.1365-2672.2006.03252.x>.
- Voudibio Mbozo, A.B., Kobawilaa, S.C., Anyogu, A., Awamaria, B., Louembe, D., Sutherland, J.P., & Ouoba, L.I.I. (2017). Investigation of the diversity and safety of the predominant *Bacillus pumilus* sensu lato and other *Bacillus* species involved in the alkaline fermentation of cassava leaves for the production of Ntoba Mbodi. *Food Control*, 82, 154-162. <https://doi.org/10.1016/j.foodcont.2017.06.018>.
- Wacoo, A.P., Mukisa, I.M., Meeme, R., Byakika, S., Wendi, D., Sybesma, W. & Kort, R. (2019). Probiotic Enrichment and Reduction of Aflatoxins in a Traditional African Maize-Based Fermented Food. *Nutrients*, 11, 265. <https://doi.org/10.3390/nu11020265>
- Walsh, A.M., Crispie, F., Daari, K., O'Sullivan, O., Martin, J.C., Arthur, C.T., Claesson, M.J., Scott, K.P., & Cotter, P.D. (2017). Strain-Level metagenomic analysis of the fermented dairy beverage nunu highlights potential food safety risks. *Applied and Environmental Microbiology*, 83, Article number: e01144 -17. <https://doi.org/10.1128/AEM.01144-17>.
- Walton, R.N., Clemens, A., Chung, J., Moore, S., Wharton, D., Haydu, L., de Villa, E., Sanders, G., Bussey, J., Richardson, D., & Austin, J. W. (2014). Outbreak of type E foodborne botulism linked to traditionally prepared salted fish in Ontario, Canada. *Foodborne Pathogens and Disease*, 11, 830 – 834. <https://doi.org/10.1089/fpd.2014.1783>.
- World Health Organisation (2015). WHO estimates of the global burden of foodborne diseases: foodborne disease burden epidemiology reference group 2007-2015. WHO Press, Geneva.
- World Health Organisation (2017a). Antimicrobial resistance in the food chain. Available from: [https://www.who.int/foodsafety/areas\\_work/antimicrobial-resistance/amrfoodchain/en/](https://www.who.int/foodsafety/areas_work/antimicrobial-resistance/amrfoodchain/en/) Accessed 15.12.2019

- World Health Organisation. (2017b). Food Safety. Available from: <https://www.who.int/news-room/fact-sheets/detail/food-safety> Accessed 15.12.2019
- World Health Organisation. (2017c). Food Safety. Available from: <https://www.who.int/news-room/q-a-detail/one-health> Accessed 18.10.2020
- World Health Organisation. (2019). Critically Important Antimicrobials for Human Medicine. 6<sup>th</sup> Revision. WHO Press, Geneva.
- Yousif, N.M., Huch, M., Schuster, T., Cho, G.S., Dirar, H.A., Holzapfel, W.H., & Franz, C.M. (2010). Diversity of lactic acid bacteria from Hussuwa, a traditional African fermented sorghum food. *Food Microbiology*, 27, 757-68. <https://doi.org/10.1016/j.fm.2010.03.012>.
- Zakpaa, H.D., Imbeah, C.M., & Mak-Mensah, E.E. (2009). Microbial characterization of fermented meat products in some selected markets in the Kumasi metropolis, Ghana. *African Journal of Food Science*, 3, 340-346. <https://doi.org/10.5897/AJFS.9000183>.