

Obstetric-neonatal care during birth and postpartum in symptomatic and asymptomatic women infected with SARS-COV-2: a retrospective multicenter study

Rafael Vila-Candel ^{1,2}; Víctor M González-Chordá ^{3*}; Francisco Javier Soriano-Vidal ^{2,4}; Enrique Castro-Sánchez ^{5,6}; Noelia Rodríguez-Blanco ^{7,8}; Ana Gómez-Seguí ⁹; Laura Andreu-Pejó ³; Cristina Martínez-Porcar ¹⁰; Carmen Rodríguez González ¹¹; Patricia Torrent-Ramos ^{12,13}; Nieves Asensio-Tomás ¹⁴; Yolanda Herraiz-Soler ^{15,16}; Ramon Escuriet ^{17,18}; Desirée Mena-Tudela ³

- ¹ Department of Nursing, Faculty of Nursing and Podiatry, Universitat de València, València, Spain
 - ² Department of Obstetrics and Gynaecology, Hospital Universitario de la Ribera, Alcira, Spain. Foundation for the Promotion of Health and Biomedical Research in the Valencian Region (FISABIO), Valencia, Spain; vila_rafcán@gva.es
 - ³ GIENF-281 Nursing Research Group. Nursing Department, Univesitat Jaume I, Castellón, Spain vchorda@uji.es ; pejo@uji.es ; dmena@uji.es
 - ⁴ Department of Obstetrics and Gynaecology, Hospital Lluís Alcanyís, Xativa, Foundation for the Promotion of Health and Biomedical Research in the Valencian Region (FISABIO), Valencia, Spain; soriano_fraivid@gva.es
 - ⁵ College of Nursing, Midwifery and Healthcare, University of West London, Brentford, Middlesex, UK
 - ⁶ Health Protection Research Unit in Healthcare-Associated Infections and Antimicrobial Resistance at Imperial College London, London, UK enrique.castrosanchez@uwl.ac.uk
 - ⁷ Department of Nursing Universidad CEU Cardenal Herrera, Plaza Reyes Católicos, 19, 03204 Elche, Spain
 - ⁸ Department of Obstetrics and Gynaecology, Hospital Marina Baixa, Spain noelia.2ww@gmail.com
 - ⁹ Department of Obstetrics and Gynaecology, Hospital Universitario y Politécnico La Fe, Valencia, Spain anagomezsegui@gmail.com
 - ¹⁰ Department of Paediatrics, Hospital Universitario de la Ribera cmartinezporcar@gmail.com
 - ¹¹ Department of Obstetrics and Gynaecology, Hospital Universitario de Vinalopó, Spain matrona.haptonomia@gmail.com
 - ¹² Preventive Medicine Service, Hospital General de Castellón, 12071 Castellón, Spain
 - ¹³ Nursing Department, Univesitat Jaume I, Castellón, Spain ptorrent@uji.es
 - ¹⁴ Department of Obstetrics and Gynaecology, Hospital Universitario y Politécnico La Fe, Valencia, Spain niasto@hotmail.com
 - ¹⁵ Department of Obstetrics and Gynaecology, Consorcio Hospital General Universitario Valencia.
 - ¹⁶ Facultat d'Infermeria i Podologia, Universitat de València, Valencia, Spain yolandaherraiz@hotmail.com
 - ¹⁷ Ghenders research group. School of Health Sciences Blanquerna, Universitat Ramon Lull, Carrer Padilla 326, 08025 Barcelona, Spain
 - ¹⁸ Catalan Health Service, Government of Barcelona, Travessera de les Corts 131, 08028 Barcelona, Spain rescuriet@gencat.cat
- * Correspondence: vchorda@uji.es

Abstract: This study analyzes the obstetric-neonatal outcomes of women in labour with symptomatic and asymptomatic COVID-19. A retrospective, multicenter, observational study was carried out between 1 March 2020 and 28 February 2021 in eight public hospitals in the Valencian Community (Spain). Chi-square test compared the obstetric-neonatal outcomes and general care for symptomatic and asymptomatic women. 11,883 births were assisted in participating centers, with 10.9‰ maternities (n = 130) infected with SARS-CoV-2. The 20.8% were symptomatic and had more complications both upon admission (p = .042) and during puerperium (p = .042), as well as transfer to intensive care unit (ICU). Percentage admission to Neonatal Intensive Care Unit (NICU) was greater among offspring of symptomatic women compared to infants born of asymptomatic women (p = .005). Compared with asymptomatic women, those with symptoms underwent less labour companion (p = .028), less early skin-to-skin contact (p = .029) and greater mother-infant separation (p = .005). The overall maternal mortality rate was 0.8%. No vertical transmission was recorded. In conclusion, symptomatic infected women are at increased risk lack of labour companion, mother-infant separation, admission to ICU and to have preterm births and NICU admissions.

Keywords: COVID-19; SARS-CoV-2; Obstetric nursing; Neonatal nursing; Delivery; Obstetric

1. Introduction

Disease due to severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) infection had caused over 4.7 million deaths worldwide by 2021 [1]. Although most countries adopted measures to contain the pandemic, including lockdowns and preventive hygiene protocols [2] as well as vaccination programs wherever vaccines were available [3], SARS-CoV-2 infection remains a significant global health threat.

With regard to coronavirus disease-2019 (COVID-19), pregnant women do not seem to be more susceptible to infection than the general population [4]. However, in the event of disease symptoms appear to be more severe, particularly during the third trimester of pregnancy, with more frequent admission to the Intensive Care Unit (ICU) and a greater risk of poorer maternal and neonatal health outcomes (such as preterm birth, caesarean sections, and low birth weight) [5–8].

Routine evidence-based clinical practices of benefit for maternal and childbirth care (labour companion, early skin-to-skin contact, breastfeeding and rooming-in in the maternity) have been modified or interrupted during the pandemic, which could reduce the quality of birth care models [9,10]. Furthermore, vertical transmission seems possible [11], although the mechanisms of such transmission remain unclear [12,13]. Whilst severe events for newborn infants seem rare [13], perhaps thanks to the passive transmission of anti-SARS-CoV-2 antibodies via the transplacental route and in breast milk, there is growing concern about such effects [14,15].

The primary aim of this study was to analyze the obstetric-neonatal outcomes of women in labour with symptomatic and asymptomatic SARS-CoV-2 infection. The study also aimed to describe and contrast the routine care received by these groups of women and their newborns.

2. Materials and Methods

2.1. Study population and sampling criteria

A retrospective, multicenter, observational study was carried out based on the review of the clinical records of pregnant women assisted during labour and birth, with positive real-time polymerase chain reaction (RT-PCR) test result for SARS-CoV-2 in nasopharyngeal exudate at the time of admission. The study was carried out in eight state-funded hospitals of the Valencian Community in Spain. These hospitals (four in Valencia, one in Castellón, three in Alicante) were all reference centers for their province, and cared for at least 1,000 deliveries per year, or were located in rural areas with large catchment populations. Overall, the participating hospitals served one million people and attended ~12,000 deliveries in the previous year. The information about the study was disseminated via the regional research network and suitable health care organizations approached by the researchers. The study period was from 1 March 2020 to 28 February 2021.

The study population comprised women giving birth in any of the participating hospitals. The inclusion criteria were: (a) women with a positive RT-PCR test for SARS-CoV-2 RNA in nasopharyngeal exudate performed on hospital admission for labour and birth; and (b) infants born of infected mothers, with RT-PCR testing for SARS-CoV-2 in nasopharyngeal exudate during hospital admission (<48h, not extracted from the placenta or amniotic fluid).

Pregnant women infected with SARS-CoV-2 and admitted to the hospital for medical reasons other than birth were excluded from the study.

2.2. Measurements

The research staff at each participating center reviewed the obstetric history, neonatal and postpartum outcomes, and general labour care records of all the positive patients assisted during the study period. The variables related to birth and postpartum were collected from the Orion Logis® electronic health records, while the data for mothers and newborns related to follow-up during the first six weeks after birth were obtained from the Abucasis II® health database. Any readmission via the emergency room of mothers or newborns registered at the same hospitals were identified during the 6 weeks from the Abucasis II® database. Both electronic medical records are routinely used by all the health facilities in the Valencian Community, including all the centers participating in the study. The following variables were collected:

-
- Demographic variables: maternal age, country of origin, and the hospital where the birth occurred. 99
100
 - Obstetric-neonatal variables: Gestational age at the time of birth, parity (primiparous/multiparous), previous maternal history of health problems (diabetes mellitus/hypertension/cardiac diseases/neurological diseases/thrombotic diseases/thyroid diseases/drugs misuse/ COVID-not related infection diseases), gestational disease (preeclampsia/eclampsia/gestational hypertension/gestational diabetes/hypothyroidism/hyperthyroidism/ COVID-not related infectious diseases), fetal alterations identified by health care providers (preterm birth, small for gestational age/large for gestational age/fetal growth restriction/congenital abnormality), start of labour (spontaneous/induced/elective caesarean section), type of birth (eutocic, instrumental birth/caesarean section [CS]), cause of CS, maternal complications prior labour and during puerperium (respiratory, cardiac, neurologic, thrombotic manifestations, and COVID-related coagulopathy), maternal admission to the intensive care unit (ICU) prior to birth and/or puerperium, cause of maternal admission to the ICU, weight of the newborn, Apgar score at one and five minutes, and umbilical artery pH at birth. Mother to child transmission can occur in different stages, including in utero, intrapartum or postnatal. Our study defined vertical transmission in the early postnatal period (<48h) as a positive test for SARS-CoV-2 maternal infection at admission, coupled with a confirmed positive test for the newborn [16], infant admission to the neonatal ICU (NICU), and cause of admission to the NICU. 101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
 - Clinical variables: Maternal and newborn symptoms compatible with SARS-CoV-2 infection (fever, cough, shortness of breath, fatigue, body aches, headache, anosmia, ageusia, nausea or vomiting, diarrhea), time of result of the RT-PCR SARS-CoV-2 test for the mother (ante-, intra- or postpartum), the result of the RT-PCR SARS-CoV-2 test for the newborn infant within 48h (positive or negative), and follow-up of complications and readmissions (mother/newborn infant, with reasons) during the first six weeks after birth. Maternal or newborn death (up to 28 days) if COVID-related. 117
118
119
120
121
122
 - Obstetric and neonatal general labour care: late clamping (clamping and cutting the cord at least one minute from birth, or when the umbilical cord stopped pulsating [17]), early skin-to-skin contact [SSC] (defined as prone placing of the naked infant on the mother's bare chest at birth, in the first minute after birth, or very soon afterwards [18]), labour companion, mother-infant separation during hospital admission, the reason for separation. 123
124
125
126
127
 - Breastfeeding-related variables: Type of feeding at discharge and 6 weeks postpartum (exclusive breastfeeding, formula feeding, mixed feeding). Exclusive breastfeeding (EBF) was defined as offering only breast milk and excluding all other food or fluids, including water. This case definition did, however, allow the infant to receive oral rehydration salts, drops and syrups (vitamins, minerals and medicines) [19]. Mixed feeding was defined as the combination of breastfeeding and formula feed. 128
129
130
131
132
Obstetric and neonatal outcomes were compared between symptomatic and asymptomatic SARS-CoV-2 infected women. 133
134
 - The obstetric outcomes were: Preterm birth < 37 weeks; Fetal growth restriction (birth weight percentile <5); Induced hypertension in pregnancy; Gestational diabetes; CS; Operative birth; ICU admission; Maternal complications prior labour/during puerperium COVID-related: cardiac, neurologic, thrombotic, respiratory manifestations; Maternal death 135
136
137
138
The neonatal outcomes were: Small for gestational age (birth weight percentile <10 for babies of the same gestational age); Large for gestational age (birth weight p>90 for babies of the same gestational age); Apgar <7 at 5 minutes; NICU admission. 139
140
141
- The obstetric and neonatal general care were compared between symptomatic (participants with SARS-Cov-2 infection and who presented compatible clinical manifestations such as fever, cough or loss of smell, among other signs and symptoms) and asymptomatic (participants with SARS-Cov-2 infection but who did not develop compatible clinical manifestations) women SARS-CoV-2 infected were: Labour companion; Late umbilical cord clamping; Early SSC; Mother-infant separation; EBF (at discharge/6 weeks). 142
143
144
145
146
147

2.3. Statistical analysis

148

Basic descriptive statistics were expressed as the mean \pm standard deviation (SD) in the case of continuous variables and as frequencies and percentages in the case of categorical variables. Normal data distribution was assessed using the Kolmogorov-Smirnov test.

The incidence rate was determined using the total number of new positive SARS-CoV-2 cases per 1,000 maternities and divided by the number of births during the study period. The chi-square test was used to compare the obstetric-neonatal general care rates according to the different qualitative variables, corrected by the Fisher exact test. Likewise, the chi-square test and crude odds ratio (OR) with 95% confidence intervals (CIs) were used to compare the different study variables between asymptomatic SARS-CoV-2 positive women and women who developed COVID-19 (symptomatic cases). Statistical significance was considered for $p < .05$. Data analysis was performed using the R statistical package (version 4.0.5).

2.4. Ethical considerations

Patients were not involved in the development of the research questions, the study's design, or the recruitment of participants. Because of the retrospective study design, patient informed consent was not required. All patient data were handled anonymously. The local Ethics Committees approved the study protocols in the eight centers. The ethical principles of medical research that current Spanish legislation contemplated have been considered, and the study was conducted following the Declaration of Helsinki.

3. Results

11,883 births were assisted during the study period in the participating hospitals. We included all women with SARS-CoV-2 infection admitted to the participating hospitals during the study period (130, 10.9 per 1,000 maternities). The case distribution according to hospitals is shown in Table 1. Hospital H4 had an incidence of 17.8 per 1,000 maternities, and hospital H8 assisted the largest number of positive cases. The majority of women were born in Spain (53.8%), with a mean age of 32 ± 5.1 years. There were not statistically significant differences between symptomatic and asymptomatic women for any demographic variables (Table 2). There was no loss of any participant.

Table 1. Births and incidence of women with SARS-CoV-2 infection during the study.

Hospital	Total births	Births in women infected with SARS-CoV-2	Incidence
H1	1288	18	14.0
H2	1090	7	6.4
H3	1308	7	5.4
H4	1182	21	17.8
H5	1256	14	11.1
H6	679	11	16.2
H7	920	8	8.7
H8	4160	44	10.6
TOTAL	11,883	130	11.3

Table 2. Demographic characteristics of women infected with SARS-CoV-2 in the study.

182

Study variables	Women infected with SARS-CoV-2		p*
	Asymptomatic (n=103; 79.2%)	Symptomatic (n=27; 20.8%)	
	Mean (SD)	Mean (SD)	
Age	31.5 (4.9)	33.9 (5.5)	.05
	n (%)	n (%)	
Country of origin			
Spain	64 (81.0)	6 (29.0)	.287
Central and South America	14 (70.0)	15 (30.0)	
Rest of EU countries	12 (80.0)	3 (20.0)	
Africa	9 (81.8)	2 (18.2)	
Asia	4 (80.0)	1 (20.0)	
Hospital of birth			
H8	34 (97.1)	1 (2.9)	.11
H1	18 (100.0)	0 (0.0)	
H4	15 (71.4)	6 (28.6)	
H5	10 (71.4)	4 (28.6)	
H6	8 (72.7)	3 (27.3)	
H3	7 (70.0)	3 (30.0)	
H2	6 (37.5)	10 (62.5)	

*Chi-square; severe acute respiratory syndrome coronavirus-2 = SARS-CoV-2; EU = European Union; H = Hospital.

183
184

Regarding the women's characteristics, 50% were primiparous, with a mean gestational age of 39.2 ± 1.58 weeks, and most had no previous maternal, gestational or fetal disease conditions, without significant differences between symptomatic and asymptomatic women ($P < 0.05$). Positivity for SARS-CoV-2 was confirmed before or during labour in 91.5% of cases and 8.5% after birth. 56.2% had laboured spontaneously, with an overall induction rate of 39.2%. Induction of labour was higher, but without significant differences, among symptomatic women (44.4% vs 37.9%; $P = .09$). The 4.6% of women had an elective-CS, without significant differences between symptomatic and asymptomatic women ($P = 0.09$). There were no statistically significant differences between symptomatic and asymptomatic women in terms of maternal ($P = .084$), gestational ($P = .089$) or fetal disease conditions ($P = .719$).

185
186
187
188
189
190
191
192
193

In terms of the clinical parameters, 20.8% of women had symptoms upon admission being fever and cough (22.2% each) and headache (18.5%), the most frequent manifestations. Other symptoms were 11.1% anosmia, 11.1% dyspnea, 3.7% vomiting, 3.7% ageusia and, 7.4% body aches. Prior to birth, 7.4% of symptomatic women were admitted to the ICU with COVID-related respiratory distress versus none of the asymptomatic women ($P = .042$); of these women, 3.7% required mechanical ventilation due to dyspnea and COVID-related thrombotic stroke. 7.4% of symptomatic women were admitted to the ICU during the postpartum period compared with 1.9% asymptomatic women ($P = .042$). Mechanical ventilation was not required for any asymptomatic women admitted to the ICU. The admissions were not COVID-related (1 preeclampsia, and 1 postpartum hemorrhage). However, both symptomatic women required mechanical ventilation. The mortality rate was 0.8%; among all participants, death was recorded in a woman with symptoms (specific mortality rate = 3.7%) (Table 3).

194
195
196
197
198
199
200
201
202
203
204

205

206

207

208

Table 3. Obstetric-neonatal characteristics of the study sample.

Variable	Women infected with SARS-CoV-2 (n / %)		p*
	Asymptomatic (103 / 79.2%)	Symptomatic (27 / 20.8%)	
	Mean (SD)	Mean (SD)	
Gestational age (weeks)	39.4 (1.4)	38.8 (2.2)	.595
Apgar 1 minute	9.63 (0.8)	9.56 (1.0)	.694
Apgar 5 minutes	9.93 (0.4)	9.93 (0.4)	.825
	n (%)	n (%)	
Parity			
Multiparous	52 (80.0)	13 (20.0)	.829
Primiparous	51 (78.5)	14 (21.5)	
Previous maternal disease			
No	94 (81.7)	21 (18.3)	.084
Yes	9 (60.0)	6 (40.0)	
Gestational disease			
No	88 (82.2)	19 (17.8)	.089
Yes	15 (65.2)	8 (34.8)	
Fetal disease			
No	94 (79.7)	24 (20.3)	.713
Yes	9 (75.0)	3 (25.0)	
Start of labour			
Elective caesarean section	5 (83.3)	1 (16.7)	.09
Spontaneous	59 (57.3)	14 (51.9)	
Induced	39 (37.9)	12 (44.4)	
COVID-related Maternal complications prior to labour			
Asymptomatic	103 (100.0)	23 (85.2)	.002
Respiratory distress	0 (0.0)	3 (100.0)	
Thrombotic stroke	0 (0.0)	1 (100.0)	
Type of birth			
Elective caesarean section	5 (83.3)	1 (16.7)	.269
Urgent caesarean section	11 (64.7)	6 (35.3)	
Eutocic	75 (83.3)	15 (16.7)	
Instrumental birth	12 (70.6)	5 (29.4)	
Cause of caesarean section (n=23)			
Other	10 (62.5)	6 (37.5)	.366
NRFHR	7 (87.5)	1 (12.5)	
Maternal complications COVID-related before discharge			
No complications	101 (81.4)	23 (18.6)	.002
Dyspnea	0 (0.0)	1 (100.0)	
Respiratory distress	0 (0.0)	2 (100.0)	
Others	2 (100.0)	0 (0.0)	
Death	0 (0.0)	1 (100.0)	
RT-PCR testing of newborn infant on day 1 of life			
Positive	0 (0.0)	0 (0.0)	1
Negative	103 (79.2)	27 (20.8)	
Mother required ICU admission prior to labour			
No	103 (80.5)	25 (19.5)	.042

Yes	0 (0.0)	2 (100.0)	
Mother required ICU admission before discharge			
No	101 (80.2)	25 (19.8)	.042
Yes	2 (50.0)	2 (50.0)	
Newborn infant required admission to NICU before discharge			
No	92 (85.2)	16 (14.8)	.005
Yes	14 (56.0)	11 (44.0)	
Mother requiring emergency readmission after discharge in first 6 weeks			
No	102 (79.1)	27 (20.9)	1
Yes	1 (100.0)	0 (0.0)	
Newborn infant requiring emergency readmission after discharge in first 6 weeks			
No	100 (79.3)	26 (20.7)	1
Yes	3 (75.0)	1 (25.0)	
Reason for maternal readmission after discharge in first 6 weeks			
Puerperal fever	1 (100.0)	0 (0.0)	1
No	102 (79.1)	27 (20.9)	
Reason for newborn infant readmission after discharge in first 6 weeks			
Choking	1 (100.0)	0 (0.0)	1
Non-COVID-19 respiratory infection	0 (0.0)	1 (100.0)	
SARS-CoV-2 + (hospital admission)	1 (100.0)	0 (0.0)	
SARS-CoV-2 + (emergency room care)	1 (100.0)	0 (0.0)	

*chi-square; severe acute respiratory syndrome coronavirus-2= SARS-CoV-2; NRFHR= non-reassuring fetal heart pattern; RT-PCR= real-time polymerase chain reaction; ICU = Intensive Care Unit; NICU = Neonatal Intensive Care Unit.

None of the newborn infants tested positive for SARS-CoV-2 infection during hospital admission. Organizational issues caused the reasons for admission of newborns to the NICU before discharge (19.2%), and none of them concerned a disease associated with maternal COVID-19. The reasons were separation following protocol active on the date (26.9%), prematurity (23.1%), stabilization to maladaptation after birth (15.4%), hyperbilirubinemia (11.5%), social services/adoption (7.6%), 3.8% sepsis (3.8%), observation following cerebral-vascular event (3.8%), and maternal drug misuse (3.8%).

The obstetric and neonatal general care is presented in Table 4. Early SSC occurred in 75.4% of deliveries, and late clamping was performed in 55.6%. Mother-infant separation occurred in 19.2% of the cases, with no clinical indication in 5.4%, and with a maternal accompaniment rate of 53.8%. On the other hand, asymptomatic women had a greater percentage of early SSC (79.6% vs 59.3%; $P = .029$), a greater labour companion rate (64.1% vs 40.7%; $P = .028$), and less mother-infant separation (13.6% vs 40.7%; $P < .001$). There were no statistically significant differences in late umbilical cord clamping (asymptomatic = 57.3%; symptomatic = 40.7%; $P = .125$).

210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232

Table 4. Obstetric-neonatal general care of the study participants.

Variable	Women infected with SARS-CoV-2 (n / %)		*P
	Asymptomatic (103 / 79.2%)	Symptomatic (27 / 20.8%)	
Labour companion			
No	37 (69.8)	16 (30.2)	.028
Yes	66 (85.7)	11 (14.3)	
Late umbilical cord clamping			
No	44 (73.3)	16 (26.7)	.125
Yes	59 (84.3)	11 (15.7)	
Early skin-to-skin contact			
No	21 (65.6)	11 (34.4)	.029
Yes	82 (83.7)	16 (16.3)	
Mother-infant separation during hospital admission			
No	89 (84.8)	16 (15.2)	<.001
Yes	14 (56.0)	11 (44.0)	
Reason for mother-infant separation			
No separation	89 (84.8)	16 (15.2)	<.001
Monitoring and control	2 (50.0)	2 (50.0)	
Other	5 (40.0)	9 (60.0)	
Protocol to date	7 (100.0)	0 (0.0)	
Feeding at discharge			
Formula feeding (maternal decision)	12 (70.6)	5 (29.4)	.450
Formula feeding (medical recommendation)	1 (25.0)	3 (75.0)	
Exclusive breastfeeding	76 (82.6)	16 (17.4)	
Mixed feeding (maternal decision)	6 (75.0)	2 (25.0)	
Mixed feeding (medical recommendation)	8 (88.9)	1 (11.1)	
Feeding at six weeks			
Formula feeding	26 (74.3)	9 (25.7)	.408
Exclusive breastfeeding	55 (79.7)	14 (20.3)	
Mixed feeding	22 (84.6)	4 (15.4)	

* chi-square; SARS-CoV-2 = severe acute respiratory syndrome coronavirus-2.

Concerning the EBF, the percentage at the time of hospital discharge was 70.8%. The EBF rate at six weeks after birth was 53.1%. We recorded no statistically significant differences in the type of feeding at hospital discharge or the type of feeding at six weeks after birth between women with and without COVID-19 symptoms.

235
236
237
238

Table 5 presents the obstetric-neonatal results and care received during labour, birth, and puerperium of women infected with SARS-CoV-2. There were differences between symptomatic and asymptomatic participants. The symptomatic women were 9 times more likely to have a preterm birth (95%CI: 2.20-41.15), eight times more likely to be admitted to ICU (95%CI:1.52-50.89), and infants were five times more likely to be admitted to NICU (95%CI:1.96-13.83).

239
240
241
242
243

Regarding general care for symptomatic women, they were more likely to be unaccompanied during labour (95%CI: 1.07-4.19), also markedly likely to experience mother-child separation (95%CI: 1.57-10.36) and twice as likely to experience non-performance of early SSC (95%CI: 1.09-4.06) compared to asymptomatic women.

244
245
246
247

CS was significantly associated to preterm birth (OR = 12.23; 95%CI: 2.79-53.62), admission to ICU (OR = 11.05; 95%CI: 1.88-64.65), and admission to NICU (OR = 5.56; 95%CI: 2.03-15.24). Additionally, mother with preterm birth was associated to ICU admission (OR = 19.66; 95%CI: 3.25-118.77).

248
249
250

Table 5. Obstetric-neonatal outcomes and general care of the participants.

		Women infected with SARS-CoV-2				OR	95% CI	p*
		Asymptomatic		Symptomatic				
		(103/ 79.2%)		(27/ 20.8%)				
		n	%	n	%			
Obstetric outcomes								
Pregnancy week	<37w	3	33.3	6	66.7	9.52	2.20-41.15	<.001
	>37w	100	82.6	21	17.4			
ICU admission	No	101	81.5	23	18.5	8.78	1.52-50.89	.005
	Yes	2	33.3	4	66.7			
Neonatal outcomes								
NICU admission	No	91	85.0	16	15.0	5.21	1.96-13.83	<.001
	Yes	12	52.2	11	47.8			
General care								
Mother-infant separation during hospital admission	No	89	83.8	16	16.2	4.03	1.57-10.36	.005
	Yes	14	56.0	11	44.0			
Early skin-to-skin-contact	No	21	65.6	11	34.4	2.11	1.09-4.06	.029
	Yes	82	83.7	16	16.3			
Labour companion	No	37	69.8	16	30.2	2.11	1.07-4.19	.028
	Yes	66	85.7	11	14.3			

*chi-square; OR = crude odds ratio; CI = confidence interval; SARS-CoV-2 = severe acute respiratory syndrome coronavirus-2; EBF = exclusive breastfeeding; ICU = intensive care unit; NICU = neonatal intensive care unit.

4. Discussion

This study identified an increased risk of adverse obstetric-neonatal outcomes (preterm birth and ICU/NICU admission) and poorer obstetric-neonatal general care (mother-infant separation, not early SSC, lack of labour companion) among symptomatic SARS-CoV-2 infected pregnant women versus those without symptoms of the infection.

In our study, the incidence of infection among women in labour was low (11.3 per 1,000 maternities). In addition, only 20.8% of women in the study developed symptoms consistent with COVID-19, in line with observations by other authors [10,20]. The most common symptoms reported were fever, headache and cough, again in common with other studies [8,20–24].

4.1. Obstetrical-neonatal outcomes

In our series, the induction rate was significantly higher among symptomatic women. This was possibly related to patient care adjustments made by the clinical teams at the labour wards of the different hospitals based on available human and material resources [10].

With an observed increase in the probability of adverse outcomes in symptomatic versus asymptomatic women, the recorded incidence of CS and preterm births was lower than reported elsewhere [24–27]. This low incidence of CS in our series compared to other studies [2,8,28] may be explained by other factors. Healthcare professionals may have adopted a more expectant attitude in childbirth, driven by a lack of knowledge of this new disease, the risk and concern about infection among the professionals [29–31]. Nevertheless, we found an association between CS and admission to ICU and NICU [32,33]. This may reflect an association towards the indirect impact of SARS-CoV-2 on maternity care and perinatal morbidity [9,24,34].

Even though the percentage of complications was low, as per other studies [8] the risk of an adverse outcome in the prepartum and/or postpartum period increased in symptomatic patients, especially those with severe disease. In addition, as in other studies have shown [5,20,24,30], the maternal mortality rate attributed to COVID-19 (0.8%) was higher. Our study identified no association between maternal background disease and the manifestation of COVID-19 symptoms. However, even with a

lower reported ICU admission rate [5,12,35], maternal complications associated with the infection and required admission to the ICU were only observed in symptomatic women, with pneumonia and thrombotic stroke being the most frequent conditions.

4.2. *Obstetric-neonatal outcomes and general care.*

Significant differences were observed in maternal labour companion, SSC, and mother-infant separation after birth, with lower rates among the symptomatic women. The benefits of labour companionship are already known, with increasing spontaneous vaginal birth, decreased caesarean birth and instrumental birth, and improving general satisfaction with childbirth [36,37]. Even though it may have a negative impact on our study outcomes, it was out of our sphere of analysis. Initially, separation was carried out in all cases driven by the Spanish Society of Neonatology recommendations at the time of the study [9,38]. Both practices, the limited labour companionship and mother-infant separation, may have been influenced how the practice was modified as recommended by infection prevention and control (IPC) guidelines [39,40], driven by the fear of infection and the particularities of the healthcare system [41]. As reported by Coxon et al. [10], such measures may have been adopted in a context of uncertainty, even some medical paternalism [42] regarding the risks for the newborn infants or the professionals; in this regard, it seems that mother-infant separation occurred more often in the early days of the pandemic [10].

Although the results do not show statistically significant differences between groups of women, there is variation in the percentage of breastfeeding between groups. It is observed that symptomatic mothers have a lower percentage of follow-up, less SSC and a higher percentage of mother-infant separation. All this can influence the results of breastfeeding [43].

In terms of neonatal outcomes, over 95% of infants born to SARS-CoV-2 positive mothers were in good general condition at birth, as with previous reports [5,44]. However, admission to the NICU was significantly more frequent among infants born to symptomatic mothers. Thus, although the vertical transmission of SARS-CoV-2 is possible, current data suggests that it remains uncommon with uncertainty about this event's degree and timing of this event [42,45–47]. Only 1.6% of newborn infants were infected in the first month of life, suggesting a lack of vertical transmission after birth in all the infants. Further, this observation would support the data from different studies indicating that horizontal transmission generally occurs in the home environment [44,47,48].

EBF at discharge was the most frequent type of feeding in both groups and persisted in all the symptomatic women that chose EBF after birth. The possibility of SARS-CoV-2 transmission through breast milk is unclear, and there is no evidence that the virus is viable and transmitted through breast milk [47]. Furthermore, although breastfeeding enables the transmission of immune mechanisms from the mother to the infant [49,50], and thus benefits of breastfeeding may outweigh the potential risk of transmission [13,44], considerations were also required for women to maintain any recommended SARS-CoV-2 infection prevention measures as well as hygienic measures when breastfeeding [48,51,52].

4.3. *Strengths and limitations*

We are aware that our study has some limitations. Firstly, not all the hospitals of the Valencian Community were included in the study. Even though eight of the 26 hospitals with the most significant number of annual childbirths in our region were represented, the sample size may be small, though we included all positive cases assisted in the participating centers. The magnitude of the OR obtained should be considered in light of the limited sample size, corresponding to obstetrical outcomes and general care. This could have led to overestimating the likelihood of these outcomes. Finally, the study was based on data obtained from electronic care records, with the usual potential for information bias and the quality of the information included in the records. This bias may have been mitigated as recorded variables were common to all participating centers and the clinical records referred to general birth care. In addition, the use of registry data limited access to variables of interest such as body mass index, ethnicity or economical estatus.

Despite these limitations, our study reflects the scant evidence on the possible differences in obstetric and neonatal outcomes and general care between symptomatic and asymptomatic women infected with SARS-CoV-2. Our study took place in the first year of the COVID-19 pandemic whether the changes that occurred by the Spanish Ministry of Health and Spanish Medical Societies were introduced simultaneously in all the centers, and they could have influenced the results obtained [9]. The findings may inform recommendations for the general care of women in a future health crisis.

5. Conclusions

The symptomatic infected women are at increased risk of admission to ICU and to have preterm births and NICU admissions of their newborns compared with asymptomatic women. Further, symptomatic infected women are increased risk of mother-infant separation, not early SSC and lack of labour companion compared with asymptomatic women. The incidence of SARS-CoV-2 infection among women giving birth was low, and no vertical transmission was recorded in any infants after birth.

Author Contributions: Conceptualization and methodology, R.V.; formal analysis, D.M.T. R.V.C. V.G.C.; data curation, R.V.C., F.S.V., A.G.S., Y.H.S., N.A.T., C.R.G., N.R.B., P.T.R., C.M.P.; writing—original draft preparation, ECS RVC FSV DMT VGC.; writing—review and editing, ECS RVC DMT FSV VGC NRB LAP AGS YHS CMP. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by FISABIO grant number UGP-20-245. ECS is affiliated with the National Institute for Health Research Health Protection Research Unit (NIHR HPRU) in Healthcare Associated Infections and Antimicrobial Resistance at Imperial College London in partnership with Public Health England (PHE).

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki, and approved by the Research Ethics Committee of Hospital Universitario La Ribera (HULR2020_27) and by the corresponding Committees of the rest of the participating centers.

Informed Consent Statement: Patient consent was waived due to this is a retrospective study based on recorded data.

Data Availability Statement: Data are available upon reasonable request. All necessary data are supplied and available in the manuscript; however, the corresponding author will provide the dataset upon request. All data relevant to the study are included in the article or uploaded as supplementary information.

Acknowledgments: Thanks are due to all the healthcare staff participating directly or indirectly in the care of the patients during the pandemic. We thank Francisca Corpas-Burgos (Department of Statistical Studies of Fundación FISABIO) for her support in carrying out the statistical analysis. And finally, thanks to WHO for the participation and data sharing in the project “A prospective cohort study investigating maternal, pregnancy and neonatal outcomes for women and neonates infected with SARS-CoV-2” (hrp_covid19pregnancycohort@who.int).

Conflicts of Interest: The authors declare no conflict of interest.

329
330
331
332
333
334

335
336
337
338
339
340

341
342
343
344

345
346
347

348
349
350

351
352

353
354
355

356
357
358
359
360

361

362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380

References

1. World Health Organization. WHO Coronavirus (COVID-19) Dashboard With Vaccination Data. World Health Organization. Published 2021. Available online: <https://covid19.who.int/> (accessed September 27, 2021)
2. Zhu, N., Zhang, D., Wang, W., et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med.* 2020, 382, 727-733. doi:10.1056/nejmoa2001017
3. Mathieu, E., Ritchie, H., Ortiz-Ospina, E., et al. A global database of COVID-19 vaccinations. *Nat Hum Behav.* 2021, 5(7), 947-953. doi:10.1038/s41562-021-01122-8
4. Docherty, A.B., Harrison, E.M., Green, C.A., et al. Features of 20 133 UK patients in hospital with covid-19 using the ISARIC WHO Clinical Characterisation Protocol: Prospective observational cohort study. *BMJ.* 2020, 369, m1985. doi:10.1136/bmj.m1985
5. Allotey, J., Stallings, E., Bonet, M., et al. Clinical manifestations, risk factors, and maternal and perinatal outcomes of coronavirus disease 2019 in pregnancy: Living systematic review and meta-analysis. *BMJ.* 2020, 370, 3320. doi:10.1136/bmj.m3320
6. Musa, S.S., Bello, U.M., Zhao, S., Abdullahi, Z.U., Lawan, M.A., He, D. Vertical Transmission of SARS-CoV-2: A Systematic Review of Systematic Reviews. *Viruses.* 2021, 13(9), 1877. doi:10.3390/V13091877
7. Rodríguez-Blanco, N., Vegara-Lopez, I., Aleo-Giner, L., Tuells, J. Scoping review of coronavirus case series (SARS-CoV, MERS-CoV and SARS-CoV-2) and their obstetric and neonatal results. *Rev Esp Quimioter.* 2020, 33(5), 313-326. doi:10.37201/req/064.2020
8. Villar, J., Ariff, S., Gunier, R.B., et al. Maternal and Neonatal Morbidity and Mortality Among Pregnant Women With and Without COVID-19 Infection: The INTERCOVID Multinational Cohort Study. *JAMA Pediatr.* 2021, 175(8), 817-826. doi:10.1001/JAMAPEDIATRICS.2021.1050
9. Lalaguna Mallada, P., Díaz-Gómez, N.M., Costa Romero, M., et al. Covid-19 Transmission Recommendations During. *Rev Esp Salud Pública.* 2020, 94, 1-7.
10. Coxon, K., Turienzo, C.F., Kweekel, L., et al. The impact of the coronavirus (COVID-19) pandemic on maternity care in Europe. *Midwifery.* 2020, 88, 102779. doi:10.1016/j.midw.2020.102779
11. Birkhead, M., Glass, A.J., Allan-Gould, H., Goossens, C., Wright, C.A. Ultrastructural evidence for vertical transmission of SARS-CoV-2. *Int J Infect Dis.* 2021, 111, 10-11. doi:10.1016/J.IJID.2021.08.020
12. Thompson, J.L., Nguyen, L.M., Noble, K.N., Aronoff, D.M.. COVID-19-related disease severity in pregnancy. *Am J Reprod Immunol.* 2020, 84(5), e13339. doi:10.1111/aji.13339
13. Wastnedge, E.A.N., Reynolds, R.M., van Boeckel, S.R., et al. Pregnancy and COVID-19. *Physiol Rev.* 2021, 101(1), 303-318. doi:10.1152/physrev.00024.2020
14. Moore, K.M., Suthar, M.S. Comprehensive analysis of COVID-19 during pregnancy. *Biochem Biophys Res Commun.* 2021, 538, 180-186. doi:10.1016/j.bbrc.2020.12.064
15. Lechosa-Muñiz, C., Paz-Zulueta, M., Mendez-Legaza, J.M., et al. Induction of SARS-CoV-2-Specific IgG and IgA in Serum and Milk with Different SARS-CoV-2 Vaccines in Breastfeeding Women: A Cross-Sectional Study in Northern Spain. *Int j environ res public Heal.* 2021, 18(16). doi:10.3390/IJERPH18168831
16. Sexual and Reproductive Health and Research WH (HQ). Definition and categorization of the timing of mother-to-child transmission of SARS-CoV-2. Published 2021. Available online: <https://www.who.int/publications/i/item/WHO-2019-nCoV-mother-to-child-transmission-2021.1> (accessed February 24, 2022)
17. World Health Organization. Guideline: Delayed umbilical cord clamping for improved maternal and infant health and nutrition outcomes. Published 2014. Available online: https://www.who.int/nutrition/publications/guidelines/cord_clamping/en (accessed February 24, 2022)
18. Vila-Candel, R., Duke, K., Soriano-Vidal, F.J., Castro-Sánchez, E. Affect of Early Skin-to-Skin Mother-Infant Contact in the Maintenance of Exclusive Breastfeeding: Experience in a Health Department in Spain. *J Hum Lact.* 2018, 34(2), 304-312. doi:10.1177/0890334416676469
19. WHO. Breastfeeding. Published 2015. Available online: <https://www.who.int/news-room/questions-and-answers/item/breastfeeding> (accessed February 24, 2022)
20. Berghella, V., Brenna, L. COVID-19: Labor, birth, and postpartum issues and care - UpToDate. Published 2021. Available on line: <https://www.uptodate.com/contents/covid-19-labor-birth-and-postpartum-issues-and-care> (accessed February 24, 2022).
21. Zambrano, L.D., Ellington, S., Strid, P., et al. Update: Characteristics of Symptomatic Women of Reproductive Age with Laboratory-Confirmed SARS-CoV-2 Infection by Pregnancy Status — United States, January 22–October 3, 2020. *Morb Mortal Wkly Rep.* 2020, 69(44), 1641-1647. doi:10.15585/mmwr.mm6944e3
22. Martínez-Pérez, O., Prats Rodríguez, P., Muner Hernández, M., et al. The association between SARS-CoV-2 infection and preterm delivery: a prospective study with a multivariable analysis. *BMC Pregnancy Childbirth.* 2021, 21(1). doi:10.1186/s12884-021-03742-4
23. Jafari, M., Pormohammad, A., Sheikh Neshin, S.A., et al. Clinical characteristics and outcomes of pregnant women with COVID-19 and comparison with control patients: A systematic review and meta-analysis. *Rev Med Virol.* 2021, 31(5), 1-16. doi:10.1002/rmv.2208

-
24. Khoury, R., Bernstein, P.S., Debolt, C., et al. Characteristics and outcomes of 241 births to women with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection at Five New York City Medical Centers. *Obstet Gynecol.* 2020, 136(2), 273-282. doi:10.1097/AOG.0000000000004025 440-442
25. Royal College of Obstetricians and Gynaecologists. Coronavirus (COVID-19) infection and pregnancy. Published 2021. Available on line: <https://www.rcog.org.uk/coronavirus-pregnancy> (accessed September 27) 443-444
26. World Health Organization. Clinical Management of Severe Acute Respiratory Infection (SARI) When COVID-19 Disease Is Suspected: Interim Guidance, 13 March 2020. World Health Organization. Available online: <https://apps.who.int/iris/handle/10665/331446> (accessed February 24, 2022). 445-447
27. Donati, S., Corsi, E., Salvatore, M.A., et al. Childbirth care among sars-cov-2 positive women in Italy. *Int J Environ Res Public Health.* 2021, 18(8). doi:10.3390/ijerph18084244 448-449
28. Rasmussen, S.A., Smulian, J.C., Lednický, J.A., Wen, T.S., Jamieson, D.J. Coronavirus Disease 2019 (COVID-19) and pregnancy: what obstetricians need to know. *Am J Obstet Gynecol.* 2020, 222(5), 415-426. doi:10.1016/j.ajog.2020.02.017 450-452
29. Ferrazzi, E., Frigerio, L., Savasi, V., et al. Vaginal delivery in SARS-CoV-2-infected pregnant women in Northern Italy: a retrospective analysis. *An Int J Obstet Gynaecol.* 2020, 127(9), 1116-1121. doi:10.1111/1471-0528.16278 453-455
30. Sadler, M., Leiva, G., Olza, I. COVID-19 as a risk factor for obstetric violence. *Sex Reprod Heal Matters.* 2020, 28(1), 1-3. doi:10.1080/26410397.2020.1785379 456-457
31. Tolcher, M.C., McKinney, J.R., Eppes, C.S., et al. Prone Positioning for Pregnant Women With Hypoxemia Due to Coronavirus Disease 2019 (COVID-19). *Obstet Gynecol.* 2020, 136(2), 259-261. doi:10.1097/AOG.0000000000004012 458-459
32. Giesbers, S., Goh, E., Kew, T., et al. Treatment of COVID-19 in pregnant women: A systematic review and meta-analysis. *Eur J Obstet Gynecol Reprod Biol.* 2021, 267, 120-128. doi:10.1016/j.ejogrb.2021.10.007 460-462
33. Allotey, J., Stallings, E., Bonet, M., et al. Clinical manifestations, risk factors, and maternal and perinatal outcomes of coronavirus disease 2019 in pregnancy: Living systematic review and meta-analysis. *BMJ.* 2020, 370. doi:10.1136/bmj.m3320 463-465
34. Vousden, N., Bunch, K., Morris, E., et al. The incidence, characteristics and outcomes of pregnant women hospitalized with symptomatic and asymptomatic SARS-CoV-2 infection in the UK from March to September 2020: A national cohort study using the UK Obstetric Surveillance System (UKOSS). *PLoS One.* 2021, 16, 1-19. doi:10.1371/journal.pone.0251123 466-469
35. World Health Organization. Definition and categorization of the timing of mother-to-child transmission of SARS-CoV-2. Published 2021. Available online: <https://www.who.int/publications/i/item/WHO-2019-nCoV-mother-to-child-transmission-2021.1> (accessed February 24, 2022) 470-472
36. Bohren, M.A., Hofmeyr, G.J., Sakala, C., Fukuzawa, R.K., Cuthbert, A. Continuous support for women during childbirth. *Cochrane database Syst Rev.* 2017, 7(7), CD003766. doi:10.1002/14651858.CD003766.pub6 473-474
37. Bohren, M.A., Munthe-Kaas, H., Berger, B.O., Allanson, E.E., Tunçalp, Ö. Perceptions and experiences of labour companionship: a qualitative evidence synthesis. *Cochrane Database Syst Rev.* 2016, 2016(12), CD012449. doi:10.1002/14651858.CD012449 475-477
38. SeNeo Sociedad Española de Neonatología. Recomendaciones para el manejo del recién nacido en relación con la infección por SARS-CoV-2. Published 2020. Available online: https://www.seneo.es/images/site/noticias/home/Recomendaciones_SENeo_SARS-CoV-2Version_2.pdf (accessed February 25, 2022). 478-480
39. Dong, L., Tian, J., He, S., et al. Possible Vertical Transmission of SARS-CoV-2 from an Infected Mother to Her Newborn. *J Am Med Assoc.* 2020, 12. doi:10.1001/jama.2020.4621 481-482
40. Alonso Díaz, C., López Maestro, M., Moral Pumarega, M.T., Flores Antón, B., Pallás Alonso, C.R. First case of neonatal infection due to SARS-CoV-2 in Spain. *An Pediatr.* 2020, 92(4), 237-238. doi:10.1016/j.anpedi.2020.03.002 483-484
41. Dotters-Katz, S.K., Hughes, B.L. Considerations for Obstetric Care during the COVID-19 Pandemic. *Am J Perinatol.* 2020, 37, 773-779. doi:10.1055/s-0040-1710051 485-486
42. Houghton, C., Meskell, P., Delaney, H., et al. Barriers and facilitators to healthcare workers' adherence with infection prevention and control (IPC) guidelines for respiratory infectious diseases: A rapid qualitative evidence synthesis. *Cochrane Database Syst Rev.* 2020, 4(4), 1-55. doi:10.1002/14651858.CD013582 487-489
43. Császár-Nagy, N., Bókkon, I. Mother-newborn separation at birth in hospitals: A possible risk for neurodevelopmental disorders? *Neurosci Biobehav Rev.* 2018, 84, 337-351. 490-491
44. Marín Gabriel, M.A., Cuadrado, I., Álvarez Fernández, B., et al. Multicentre Spanish study found no incidences of viral transmission in infants born to mothers with COVID-19. *Acta Paediatr Int J Paediatr.* 2020, 109(11), 2302-2308. doi:10.1111/apa.15474 492-494
45. Bohren, M.A., Berger, B.O., Munthe-Kaas, H., Tunçalp, Ö. Perceptions and experiences of labour companionship: A qualitative evidence synthesis. *Cochrane Database Syst Rev.* 2019, 2019(3). doi:10.1002/14651858.CD012449.pub2 495-497

46. World Health Organization. Breastfeeding and COVID-19. Available online: https://www.who.int/publications/i/item/WHO-2019-nCoV-Sci_Brief-Breastfeeding-2020.1 (accessed February 24, 2022). 498
499

47. Allotey, J., Chatterjee, S., Kew, T., et al. SARS-CoV-2 positivity in offspring and timing of mother-to-child transmission: living systematic review and meta-analysis. *BMJ*. 2022, 376. doi:10.1136/bmj-2021-067696 500
501

48. Vila-Candel, R., Mena-Tudela, D., Gómez-Seguí, A., Asensio-Tomás, N., Cervera-Gasch, A., Herraiz-Soler, Y. Management of labour, puerperium, and lactation in SARS-CoV-2 positive women. Multicentric study in the Valencian Community. *Enf Clín*. 2021, 31(3), 184-188. doi:10.1016/J.ENFCLE.2021.01.002 502
503
504

49. Fox, A., Marino, J., Amanat, F., et al. Robust and Specific Secretory IgA Against SARSCoV-2 Detected in Human Milk. *iScience*. 2020, 23(11). doi:10.1016/j.isci.2020.101735 505
506

50. Selma-Royo, M., Bäuerl, C., Mena-Tudela, D. et al. Anti-SARS-CoV-2 IgA and IgG in human milk after vaccination is dependent on vaccine type and previous SARS-CoV-2 exposure: a longitudinal study. *Genome Med*. 2022, 14, 42. doi: 10.1186/s13073-022-01043-9 507
508
509

51. Centers for Disease Control and Prevention. Breastfeeding and Caring for Newborns if You Have COVID-19. Available online: <https://www.cdc.gov/coronavirus/2019-ncov/if-you-are-sick/pregnancy-breastfeeding.html#breastfeeding%20this%20page%20may%20be%20also%20useful%20for%20you:%20https://www.e-lactancia.org/breastfeeding/maternal-sars-cov-2/synonym/> (accessed April 22, 2022). 510
511
512
513

52. World Health Organization. Home care for patients with suspected or confirmed COVID-19 and management of their contacts. Available online: [https://www.who.int/publications/i/item/home-care-for-patients-with-suspected-novel-coronavirus-\(ncov\)-infection-presenting-with-mild-symptoms-and-management-of-contacts](https://www.who.int/publications/i/item/home-care-for-patients-with-suspected-novel-coronavirus-(ncov)-infection-presenting-with-mild-symptoms-and-management-of-contacts) (accessed April 22, 2022) 514
515
516
517