



## **UWL REPOSITORY**

**repository.uwl.ac.uk**

Evaluation of concordance in estimation of excess mortality due to Covid-19 pandemic

Verma, Vivek, Nath, Dilip C. and Khan, Hafiz T.A. ORCID logo ORCID: <https://orcid.org/0000-0002-1817-3730> (2023) Evaluation of concordance in estimation of excess mortality due to Covid-19 pandemic. *Journal of Evaluation in Clinical Practice*, 29 (6). ISSN 1356-1294

<http://dx.doi.org/10.1111/jep.13866>

**This is the Accepted Version of the final output.**

**UWL repository link:** <https://repository.uwl.ac.uk/id/eprint/9949/>

**Alternative formats:** If you require this document in an alternative format, please contact: [open.research@uwl.ac.uk](mailto:open.research@uwl.ac.uk)

### **Copyright:**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy:** If you believe that this document breaches copyright, please contact us at [open.research@uwl.ac.uk](mailto:open.research@uwl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

### **Rights Retention Statement:**

# Evaluation of Concordance in Estimation of Excess Mortality due to Covid-19 Pandemic

## Short title: Estimation of Excess Covid-19 Mortality

Vivek Verma<sup>1\*</sup>, Dilip C. Nath<sup>2</sup>, Hafiz T.A. Khan<sup>3</sup>

<sup>1</sup> **Corresponding Author:** Department of Statistics, Assam University, Silchar, Assam 788011, India  
Email: [viv\\_verma456@yahoo.com](mailto:viv_verma456@yahoo.com)

<sup>2</sup> School of Applied and Pure Sciences, Royal Global University, Guwahati, Assam 781035, India.  
Email: [dilipc.nath@gmail.com](mailto:dilipc.nath@gmail.com)

<sup>3</sup> Public Health Group, College of Nursing, Midwifery and Healthcare, University of West London,  
Boston Manor Road, Brentford TW8 9GB, United Kingdom.  
Email: [hafiz.khan@uwl.ac.uk](mailto:hafiz.khan@uwl.ac.uk)

### Abstract

**Background:** The World Health Organization (WHO) kept track of Covid-19 data at country level on a daily basis during the pandemic that included the number of tests, infected cases, and fatalities. This daily record was susceptible to change depending on the time and place and impacted by underreporting. In addition to reporting cases of excess COVID-19-related deaths, the WHO also provided estimates of excess mortality based on mathematical models.

**Objective:** To evaluate the WHO reported and model-based estimate of excess deaths in order to determine the degree of agreement and universality.

**Methodology:** Epidemiological data gathered from nine different countries between April 2020 to December 2021 are used in this study. These countries are India, Indonesia, Italy, Russia, UK, Mexico, USA, Brazil and Peru and each of them recorded more than 1.5 million deaths from Covid-19 during these months. Statistical tools including correlation, linear regression, intra-class correlation, and Bland-Altman plots are used to assess the degree of agreement between reported and model-based estimates of excess deaths.

**Results:** The WHO derived mathematical model for estimating excess deaths due to Covid-19 was found to be appropriate for only four of the nine chosen countries, namely Italy, UK, USA, and Brazil. The other countries showed proportional biases and significantly high regression coefficients.

**Conclusion:** The study revealed that, for some of the chosen nations, the mathematical model proposed by the WHO is practical and capable of estimating the number of excess deaths brought on by COVID-19. However, the derived approach cannot be applied globally.

**Keywords:** Excess deaths, World Health Organization, intra-class correlation, Bland-Altman plots

## 1           **1. Introduction**

2   Throughout the course of the pandemic, the World Health Organization (WHO) was  
3   monitoring the effects of COVID-19. Countries reported the total number of Covid-19 cases  
4   and fatalities in their areas to the WHO, and this data has been made available. The Covid-19  
5   indices that are reported, however, fluctuate with time and place and are influenced or  
6   underreported<sup>1</sup> by established biases<sup>2</sup>. Recent studies,<sup>3-7</sup> including those from the WHO,<sup>8-10</sup>  
7   have noted that the data given does not give a clear picture of the health burden attributed to  
8   Covid-19 or the number of lives lost, both directly and indirectly, as a result of the pandemic.  
9   As a result of no testing being carried out before death, some Covid-19-related deaths were not  
10   recognised as such<sup>6,11</sup>.

11   Additionally, there have been modifications in the Covid-19<sup>7,10,12-13</sup> death certification laws  
12   that some nations have implemented. Excess mortality is seen as a more objective and  
13   comparable (cross-country) estimate of the influence of Covid-19 on mortality because using  
14   reported Covid-19 data presents certain challenges<sup>14-16</sup>. Covid-19 caused mortality in addition  
15   to other causes of death, known as “excess deaths”, that occurred in a country during a specific  
16   time period.

17   The ability to effectively implement public health programmes is aided by knowledge of excess  
18   mortality, and also helps to clarify the impact of the pandemic on the numbers. Several  
19   methodologies<sup>7-8,17-19</sup> have been developed to identify the excess mortality attributable to  
20   Covid-19 that take into consideration various assumptions regarding deaths from all causes and  
21   those related to Covid-19.

22   The question of concern is whether the mathematical model-based estimates can be followed  
23   universally and will align with reported excess deaths due to Covid-19 where both are subject  
24   to variance due to time and location and then further affected due to under-reporting. This  
25   study's objective **is** to assess WHO reported and model-based estimates of excess deaths in  
26   order to gauge levels of agreement and universality at nation level. Nine **countries**, including  
27   India, Indonesia, Italy, Russia, UK, Mexico, USA, Brazil, and Peru, recorded more than 1.5  
28   million deaths as a result of COVID-19. The degree of agreement between reported and model-  
29   based estimates of excess fatalities has been evaluated using statistical approaches such as  
30   correlation, linear regression, intra-class correlation, and Bland-Altman plots.

## 31           **2. Methodology**

1 **Data:** The “Our World in Data” website (<https://ourworldindata.org/coronavirus-source-data>;  
2 assessed on May, 30, 2022) provided data from dependable sources such as the WHO and the  
3 European Centre for Disease Prevention and Control and was used to gather country-by-  
4 country reported daily Covid-19 cases and deaths for the current study. This database was used  
5 to assess and investigate the excess mortality trend associated with the Covid-19 pandemic  
6 from March 2020 (since the WHO proclaimed Covid-19 a pandemic on March 11, 2020)  
7 through to December 2021. The WHO website ([https://www.who.int/data/sets/global-excess-  
8 deaths-associated-with-covid-19-modelled-estimates](https://www.who.int/data/sets/global-excess-deaths-associated-with-covid-19-modelled-estimates); assessed on May, 30, 2022) has been  
9 used to acquire the mathematical model-based estimates<sup>9</sup> of excess mortality resulting from the  
10 Covid-19 pandemic. The top nine countries that have reported more than 1.5 million Covid-19  
11 deaths including India and Indonesia (from Asia), Italy, Russia, and UK (from Europe), Mexico  
12 and USA (from North America), Brazil, and Peru (from South America), were taken into  
13 account in this study.

#### 14 **Statistical Analysis:**

15 Estimates of the additional mortality brought on by COVID-19 were supplied by the WHO and  
16 based on reported data and mathematical models. In this study, three distinct approaches have  
17 been used to examine assessments of increased mortality caused by COVID-19 that accord  
18 reasonably well. The first step was to determine the correlation ( $r$ ) between the difference and  
19 mean (of reported and estimated excess fatalities) and intra-class correlation (ICC) to see  
20 whether there was a pattern of similarity between them on a month-by-month basis. Also to  
21 determine whether a correlation coefficient had a high enough value to be considered  
22 acceptable in order to measure the degree of linear relationship between two variables. The  
23 second step was to look for a linear relationship between reported and model-based estimates  
24 of additional deaths caused by COVID-19 as follows:

$$25 \quad \textit{Expected}(E) = \beta_0 + \beta_1 \textit{Reported}(R) + \varepsilon$$

26 where  $\beta_0$  and  $\beta_1$  denotes the regression coefficient and  $\varepsilon$  denotes the error. In a case of good  
27 linear agreement, the values  $\beta_0$  would need to be 0 and  $\beta_1$ , closer to 1, that is, the regression  
28 model is without a constant, and both the variables are equal to zero at the origin. Thirdly, we  
29 measured the change in reported and model-based estimates of the additional fatalities caused  
30 by Covid-19 using the Bland-Altman plot. This offers a visual representation of the relationship  
31 and agreement between two paired observations gathered on the same measurement scale<sup>20-25</sup>.  
32 Suppose  $x_i$  denotes WHO reported and  $y_i$  denotes WHO model-based estimates of excess  
33 deaths due to Covid-19 at  $i^{\text{th}}$  point of time, for all  $i = 1, 2, \dots, m$ , say months, during a specified  
34 period. The Bland-Altman plot is formed by plotting the differences  $d_{ij}(= y_{ij} - x_{ij})$  in the  $j^{\text{th}}$

1 month by the  $i^{th}$  country on the vertical axis versus the averages,  $a_{ij} \left( = \frac{y_{ij} + x_{ij}}{2} \right)$  on the  
 2 horizontal axis. The mean difference is referred to as the bias ( $\bar{d}_i$ ) that indicates the mean  
 3 direction of the deviation from reported excess mortality due to Covid-19 of the  $i^{th}$  country.  
 4 The 95% reference range of the mean difference illustrates the magnitude of the systematic  
 5 difference between reported and mathematical model-based estimates of excess deaths due to  
 6 covid-19 and is labelled as limits of agreement<sup>26</sup>, which is obtained through  $(\bar{d}_i \pm 1.96 S_{\bar{d}_i})$   
 7 where

$$10 \quad S_{\bar{d}_i} = \sqrt{\frac{\sum_{j=1}^m (d_{ij} - \bar{d}_i)^2}{m - 1}}$$

8 The variance is obtained  $V(\bar{d}_i) = \frac{S_{\bar{d}_i}}{m}$ , then the 95% confidence interval (CI) has been given  
 9 by

$$11 \quad \left( \bar{d}_i \pm t_{(1-\frac{\alpha}{2}, m-1)} \sqrt{V(\bar{d}_i)} \right)$$

12 The 95% confidence intervals for the limits of agreement shows error bars, which is obtained  
 13 as

$$14 \quad \left( (\bar{d}_i \pm 1.96 S_{\bar{d}_i}) \pm t_{(1-\frac{\alpha}{2}, m-1)} \sqrt{V((\bar{d}_i \pm 1.96 S_{\bar{d}_i}))} \right)$$

15 To test the hypothesis ( $H_0$ ) that there is no difference among WHO reported and mathematical  
 16 model-based estimates of excess mortality, the Kolmogorov-Smirnov test for comparing  
 17 distributions was carried out. If this test confirmed distributional homogeneity, high positive  
 18 values of the correlation coefficient ( $r$ ), and ICC, and the value of the regression coefficient  
 19  $\beta_1$  was closer to 1 and free from proportional bias, then the agreement between WHO reported  
 20 and expected values of excess mortalities due to Covid-19 for any country could be established.  
 21 The statistical analysis was conducted using R-software version 4.1.3 and SAS university  
 22 edition.

### 23 **3. Results**

24 Table 1 displays a summary of statistics (mean of standard deviation - SD) and their 95%  
 25 confidence intervals of WHO reported and estimated Covid-19 related excess deaths and their  
 26 differences in the selected nine countries. The correlation coefficient ( $r$ ), ICC and linear  
 27 regression coefficient ( $\beta_1$ ) are displayed and their 95% tolerance limits, corresponding to each  
 28 of the 95% limits of confidence intervals known as level of agreements, are also shown. The  
 29 measures considered, analysing the accord between WHO reported and mathematical model-

1 based estimates, are statistically significant for their differences or biases to values of  $r$ , ICC  
2 and  $\beta_1$ , respectively.

3 The results showed that differences ( $d_{ij}$ ) or biases in WHO reported and estimated excess  
4 deaths, and the associated 95% lower and upper confidence intervals and tolerance limits, were  
5 significantly different and higher in four out of the nine countries. These countries were India  
6 (259927±324884), and Indonesia (50279±34653) in Asia; among European countries, only in  
7 Russia (34770± 24249), and in Mexico from North America (17426±18425). The data suggest  
8 that, on average, the estimated excess deaths measures in India, Indonesia, Russia, and Mexico  
9 were significantly higher (ranges from 17.5 thousand to 2.6 lakhs) than the reported excess  
10 death counts, which shows that there is consistent bias. On the other hand, in the remaining  
11 five countries of Italy, UK, USA, Brazil and Peru, the difference in reported and estimated  
12 Covid-19 related excess deaths were statistically insignificant, and, on average, the estimated  
13 excess deaths measures were closer (ranging from one thousand to six thousand) than the  
14 reported excess death counts.

15  
16 A high positive correlation ( $r>0.80$ ) among mean and differences of reported and estimated  
17 Covid-19 related excess deaths were observed in eight of the nine countries. Italy, UK, USA,  
18 Brazil and Peru provided excellent consistency ICC( $\geq 0.70$ ) among values of reported and  
19 estimated Covid-19 related excess deaths. Single score intra-class correlation between both the  
20 reported and estimated mortality counts showed excellent consistency between both values in  
21 Italy (0.75), UK (0.81), USA (0.76), Peru (0.88) and Brazil (0.93). To measure the accord  
22 among reported and estimated mortality patterns due to Covid-19, the value of the regression  
23 slope coefficient ( $\beta_1$ ) must be closer to one and was only found in Italy (1.18), UK(1.10),  
24 USA(1.12) and Brazil(1.1).

25  
26 Figure 1 provides a graphic demonstration of the Bland-Altman plot to show the agreement  
27 between WHO reported and expected values of excess mortalities due to Covid-19 for all the  
28 selected nine countries. The Bland-Altman plot was used to calculate the mean and difference  
29 (deviation) of estimated and reported excess deaths due to covid-19, as measures of agreement.  
30 The 95% confidence interval for difference limits included '0' in Italy (-460, 4298), UK(-  
31 2708,4434), USA(-3146,13528) and Brazil(-1120,6283), with no proportional bias.

#### 32 4. Discussion

1 The COVID-19 pandemic, one of the most significant global catastrophes in centuries, has had  
2 significant and far-reaching effects on health systems, economies, and civilizations<sup>27</sup>. Different  
3 populations have been impacted by COVID-19 and governmental responses to it in different  
4 ways. Government decisions frequently exacerbated pre-existing structural disparities in  
5 income and poverty as well as socioeconomic inequalities in education and skill levels, and in  
6 intergenerational inequalities<sup>28</sup>. Social constraints caused by the pandemic<sup>29</sup> compelled people  
7 to acclimatize themselves for solitude, that then led to increases in the prevalence of familial  
8 violence<sup>30</sup>, depression<sup>31-32</sup>, anxiety<sup>31-32</sup>, and post-traumatic stress disorder<sup>33-34</sup>. COVID-19  
9 deaths acted as an indicator for policymakers and medical professionals to take necessary  
10 decisions and proper infrastructural developments to try and control the pandemic<sup>35-37</sup>.

11 The excess mortality rate, due to Covid-19, created additional burdens on countries in terms of  
12 healthcare, finance, and life expectancy<sup>38-39</sup>. However, due to differences in time zones,  
13 geographical composition, implementation and duration of lockdowns<sup>40-41</sup>, along with socio-  
14 economic conditions in different countries, exact reporting of cases and deaths due to Covid-  
15 19 was not possible throughout the globe.<sup>42</sup> Considering the different methods employed in  
16 reporting Covid-19 figures, some mathematical models have been formulated to approximate  
17 excess mortality.<sup>9-10,12-15</sup>

18  
19 The present study is an attempt to analyse the extent, accuracy, credibility, and uniformity of  
20 agreement among the WHO reported and WHO estimated<sup>9</sup> excess mortality rates using the top  
21 nine countries that reported more than 1.5 million Covid-19 deaths. Statistically, significant  
22 proportional bias was found in five countries out of the nine, that is, India, Indonesia, Russia,  
23 Mexico and Peru. There was no proportional bias with closure (ranges from one to six  
24 thousand) with estimated values of excess deaths found in the other four countries of Italy, UK,  
25 USA, and Brazil. High positive correlation among mean and differences of reported and  
26 estimated Covid-19 related excess deaths were observed in all of the selected countries, except  
27 Indonesia, but it could not be confirmed<sup>43</sup> whether this was due to a better alignment among  
28 estimated and reported values. High ICC values in Italy, UK, USA, Brazil and Peru showed  
29 similarities among reported and estimated mortality counts. The value of regression coefficient  
30 for the null intercept regression model was found to closure to one for Italy, UK, USA, and  
31 Brazil, that defined a more reliable and significant agreement of excess mortality due to Covid-  
32 19 in terms of reported and estimated mortalities in these countries only. The Bland-Altman  
33 plot revealed that for India, Indonesia, Russia, Mexico and Peru the distributional pattern of  
34 mean and difference of estimated and reported Covid-19 related excess deaths showed

1 proportional bias, as the values increased in proportion to the average values. The Bland-  
2 Altman plot confirmed that the reported and estimated mortality counts in Italy, UK, USA and  
3 Brazil are free from proportional bias.

4 According to the study, only Italy, UK, USA, and Brazil demonstrated agreement between  
5 WHO reported and expected values of excess mortalities resulting from Covid-19. The  
6 Kolmogorov-Smirnov test confirmed their distributional homogeneity, having high positive  
7 values of correlation coefficient ( $r$ ) and ICC, value of regression coefficient ( $\beta_1$ ) (closer to 1),  
8 and free from proportional bias.

9 This study shows that proper reporting and estimation are both necessary in order to develop  
10 and/or update policy measures in order to address any pandemics in the future. The  
11 mathematical models that were developed to estimate the extra mortality must be validated and  
12 take into account the geographic makeup, pattern of incidences, healthcare infrastructures,  
13 governmental initiatives, and the socioeconomic conditions of different countries.

## 14 **5. Conclusion**

15 The WHO published estimated excess mortality rates attributable to COVID-19, as well as  
16 reported excess mortality rates, and it recommended using the estimated figures to describe the  
17 pandemic's state in various countries. The aim of this study is to describe the consistency and  
18 assumption of universality of the model-based estimates of excess mortality in the top nine  
19 countries that have reported more than 1.5 million Covid-19 deaths. To do this, excess mortality  
20 estimates from the WHO were compared with those from our model, in order to determine how  
21 well they both aligned. Results obtained indicate that only four of the nine countries, that is,  
22 Italy, UK, USA, and Brazil, were appropriate for using the WHO's mathematical approach for  
23 predicting excess mortality caused by COVID-19. The study comes to the further conclusion  
24 that larger correlation values or ICC, as well as linear associations among reported and  
25 estimated excess mortality attributable to COVID-19, do not always explain the applicability  
26 of any approach. Before claiming to be a universally accepted model and making the  
27 corresponding remark about its implications for policy, any mathematical models created to  
28 estimate the extra deaths must be validated.

## 29 **Acknowledgments**

30 Authors would like to thank all of the anonymous reviewers and editorial board for their  
31 valuable comments and suggestions to improve the quality of this manuscript.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43

**Disclosure Statement:** No potential conflict of interest was reported by the authors.

**Conflict of Interest Statements:** No potential conflict of interest was reported by the authors.

**Ethics Approval:** The database considered for the present study does not contain any individual identifiable information. Due to the unidentified nature of the dataset and no human subject were directly involved in the present study; therefore, this study was exempt from any ethical or Institutional Review Board clearance.

**Funding:** This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

**References:**

1. Kofanov, D., Kozlov, V., Libman, A., & Zakharov, N. (2022). Encouraged to Cheat? Federal Incentives and Career Concerns at the Sub-national Level as Determinants of Under-Reporting of COVID-19 Mortality in Russia. *British Journal of Political Science*, 1-26.
2. Kobak, D. (2022). Underdispersion: A statistical anomaly in reported Covid data. *Significance*, 19(2), 10-13.
3. Besson, E. S. K., Norris, A., Ghouth, A. S. B., Freemantle, T., Alhaffar, M., Vazquez, Y., ... & Checchi, F. (2021). Excess mortality during the COVID-19 pandemic: a geospatial and statistical analysis in Aden governorate, Yemen. *BMJ global health*, 6(3), e004564.
4. Biswas, R. K., Afiaz, A., & Huq, S. (2020). Underreporting COVID-19: the curious case of the Indian subcontinent. *Epidemiology & Infection*, 148.
5. Hanifi, S. M. A., Alam, S. S., Shuma, S. S., & Reidpath, D. D. (2021). Insights Into Excess Mortality During the First Months of the COVID-19 Pandemic From a Rural, Demographic Surveillance Site in Bangladesh. *Frontiers in Public Health*, 1016.
6. Lau, H., Khosrawipour, T., Kocbach, P., Ichii, H., Bania, J., & Khosrawipour, V. (2021). Evaluating the massive underreporting and undertesting of COVID-19 cases in multiple global epicenters. *Pulmonology*, 27(2), 110-115.
7. Karlinsky, A., & Kobak, D. (2021). Tracking excess mortality across countries during the COVID-19 pandemic with the World Mortality Dataset. *Elife*, 10.
8. WHO, A. (2020). WHO Methods and Data Sources for Life Tables 1990–2019.
9. WHO. (2021) The true death toll of COVID-19: estimating global excessmortality. June 29, 2021. [https://www.who.int/data/stories/the-true-death-toll-of-covid-19-estimating-global-excess-mortality (accessed June 29, 2021).]

10. Haklai, Z., Aburbah, M., Goldberger, N., & Gordon, E. S. (2021). Excess mortality during the COVID-19 pandemic in Israel, March–November 2020: when, where, and for whom?. *Israel Journal of Health Policy Research*, 10(1), 1-7.
11. Ioannidis, J. P. (2021). Over-and under-estimation of COVID-19 deaths. *European journal of epidemiology*, 36, 581-588.
12. Stokes, A. C., Lundberg, D. J., Elo, I. T., Hempstead, K., Bor, J., & Preston, S. H. (2021). COVID-19 and excess mortality in the United States: A county-level analysis. *PLoS medicine*, 18(5), e1003571.
13. Vandoros, S. (2020). Excess mortality during the Covid-19 pandemic: Early evidence from England and Wales. *Social Science & Medicine*, 258, 113101.
14. Kobak, D. (2021). Excess mortality reveals Covid's true toll in Russia. *Significance (Oxford, England)*, 18(1), 16.
15. Kupek, E. (2021). How many more? Under-reporting of the COVID-19 deaths in Brazil in 2020. *Tropical Medicine & International Health*, 26(9), 1019-1028.
16. Levitt, M., Zonta, F., & Ioannidis, J. P. (2022). Comparison of pandemic excess mortality in 2020–2021 across different empirical calculations. *Environmental Research*, 213, 113754.
17. Nogueira, P. J., de Araújo Nobre, M., Nicola, P. J., Furtado, C., & Carneiro, A. V. (2020). Excess mortality estimation during the COVID-19 pandemic: preliminary data from Portugal. *Acta Médica Portuguesa*, 33(6), 376-383.
18. Friedman, J., Liu, P., Troeger, C. E., Carter, A., Reiner, R. C., Barber, R. M., ... & Gakidou, E. (2021). Predictive performance of international COVID-19 mortality forecasting models. *Nature communications*, 12(1), 1-13.
19. Knutson, V., Aleshin-Guendel, S., Karlinsky, A., Msemburi, W., & Wakefield, J. (2022). Estimating Global and Country-Specific Excess Mortality During the COVID-19 Pandemic. *arXiv preprint arXiv:2205.09081*.
20. Bland, J. M., & Altman, D. (1986). Statistical methods for assessing agreement between two methods of clinical measurement. *The lancet*, 327(8476), 307-310.
21. Bland, J. M., & Altman, D. G. (1990). A note on the use of the intraclass correlation coefficient in the evaluation of agreement between two methods of measurement. *Computers in biology and medicine*, 20(5), 337-340.
22. Flegal, K. M., Graubard, B., & Ioannidis, J. P. (2020). Use and reporting of Bland–Altman analyses in studies of self-reported versus measured weight and height. *International Journal of Obesity*, 44(6), 1311-1318.
23. Giavarina, D. (2015). Understanding bland altman analysis. *Biochemia medica*, 25(2), 141-151.
24. Fukuoka, Y., Haskell, W., & Vittinghoff, E. (2016). New insights into discrepancies between self-reported and accelerometer-measured moderate to vigorous physical activity among women—the mPED trial. *BMC public health*, 16(1), 1-11.

- 1 25. Sharif Nia, H., Chong, P. P., Yiong Huak, C., Gorgulu, O., Taghipour, B., Sivarajan Froelicher,  
2 E., ... & Rahmatpour, P. (2022). Clinical accuracy and agreement between tympanic and  
3 forehead body temperature measurements for screening of patients with COVID-19. *Journal of*  
4 *Clinical Nursing*, 31(21-22), 3272-3285.
- 5
- 6 26. Ludbrook, J. (2010), Confidence in Altman–Bland plots: A critical review of the method of  
7 differences. *Clinical and Experimental Pharmacology and Physiology*, 37: 143-149.
- 8
- 9 27. Ali, M. J., Bhuiyan, A. B., Zulkifli, N., & Hassan, M. K. (2022). The COVID-19 pandemic:  
10 Conceptual framework for the global economic impacts and recovery. *Towards a Post-Covid*  
11 *Global Financial System*, 225-242.
- 12
- 13 28. Blundell, R., Cribb, J., McNally, S., Warwick, R., & Xu, X. (2021). Inequalities in education,  
14 skills, and incomes in the UK: The implications of the COVID-19 pandemic. *Institute for Fiscal*  
15 *Studies*, 1-42.
- 16
- 17 29. Hosseinzadeh, P., Zareipour, M., Baljani, E., & Moradali, M. R. (2022). Social Consequences  
18 of the COVID-19 Pandemic. A Systematic Review. *Investigación y Educación en Enfermería*,  
19 40(1).
- 20
- 21 30. Boxall, H., Morgan, A., & Brown, R. (2020). The prevalence of domestic violence  
22 among women during the COVID-19 pandemic. *Australasian Policing*, 12(3), 38-46.
- 23
- 24 31. Ettman, C. K., Abdalla, S. M., Cohen, G. H., Sampson, L., Vivier, P. M., & Galea, S.  
25 (2020). Prevalence of depression symptoms in US adults before and during the  
26 COVID-19 pandemic. *JAMA network open*, 3(9), e2019686-e2019686.
- 27
- 28 32. Lakhan, R., Agrawal, A., & Sharma, M. (2020). Prevalence of depression, anxiety,  
29 and stress during COVID-19 pandemic. *Journal of neurosciences in rural practice*,  
30 11(04), 519-525.
- 31
- 32 33. Alshehri, F. S., Alatawi, Y., Alghamdi, B. S., Alhifany, A. A., & Alharbi, A. (2020).  
33 Prevalence of post-traumatic stress disorder during the COVID-19 pandemic in Saudi  
34 Arabia. *Saudi Pharmaceutical Journal*, 28(12), 1666-1673.
- 35
- 36 34. Karbasi, Z., & Eslami, P. (2022). Prevalence of post-traumatic stress disorder during  
37 the COVID-19 pandemic in children: A review and suggested solutions. *Middle East*  
38 *Current Psychiatry*, 29(1), 1-9.
- 39
- 40 35. Ozili, P. K. (2021). Covid-19 pandemic and economic crisis: The Nigerian experience and  
41 structural causes. *Journal of Economic and Administrative Sciences*, 37(4), 401-418.
- 42
- 43 36. Nicodemo, C., Barzin, S., Lasserson, D., Moscone, F., Redding, S., & Shaikh, M. (2020).  
44 Measuring geographical disparities in England at the time of COVID-19: results using a  
45 composite indicator of population vulnerability. *BMJ open*, 10(9), e039749.
- 46
- 47 37. Marzouk, M., Elshaboury, N., Azab, S., Megahed, A., & Metawie, M. (2022). Assessment of  
48 COVID-19 impact on sustainable development goals indicators in Egypt using fuzzy analytic  
49 hierarchy process. *International Journal of Disaster Risk Reduction*, 82, 103319.
- 50
- 51 38. Moolla, I., & Hiilamo, H. (2023). Health system characteristics and COVID-19 performance in  
52 high-income countries. *BMC Health Services Research*, 23(1), 1-14.
- 53

- 1 39. Msemburi, W., Karlinsky, A., Knutson, V., Aleshin-Guendel, S., Chatterji, S., & Wakefield, J.  
2 (2023). The WHO estimates of excess mortality associated with the COVID-19 pandemic.  
3 *Nature*, 613(7942), 130-137.  
4
- 5 40. Verma, B. K., Verma, M., Verma, V. K., Abdullah, R. B., Nath, D. C., Khan, H. T., ... & Verma,  
6 V. (2020). Global lockdown: An effective safeguard in responding to the threat of COVID-19.  
7 *Journal of Evaluation in Clinical Practice*, 26(6), 1592-1598.  
8
- 9 41. Lancet, T. (2020). India under COVID-19 lockdown. *Lancet (London, England)*, 395(10233),  
10 1  
11
- 12 42. Willems, S. J., Castells, M. C., & Baptist, A. P. (2022). The magnification of health disparities  
13 during the COVID-19 pandemic. *The Journal of Allergy and Clinical Immunology: In Practice*,  
14 10(4), 903-908.  
15
- 16 43. Dogan, N. Ö. (2018). Bland-Altman analysis: A paradigm to understand correlation and  
17 agreement. *Turkish journal of emergency medicine*, 18(4), 139-141.  
18





Table 1: Descriptive statistics for WHO reported and expected values of excess mortalities due to Covid-19





Parameters	Asia		Europe			North America		South America	
	India	Indonesia	Italy	Russia	UK	Mexico	USA	Brazil	Peru
Months	18	19	20	22	16	20	22	22	22
Expected (E) (M±SD)	285449± 350303	57778± 36536	8261± 8432	48528± 32023	9738± 11971	31769± 24486	42659± 31289	30732± 23056	12999± 10383
95% CI	111247, 459650	40169, 75388	4315, 12207	34330, 62726	3359, 16117	20309, 43229	28786, 56532	20510, 40955	8395, 17602
Reported (R) (M±SD)	25521± 28020	7499± 10794	6342± 6433	13758± 10532	8875± 9198	14343± 7606	37468± 23340	28151± 21234	9245± 7812
95% CI	11587, 39455	2296, 12702	3331, 9353	9088, 18428	3974, 13776	10783, 17902	27119, 47817	18736, 37566	5781, 12709
Diff (E-R) or Bias(Diff) (M±SD)	259927± 324884	50279± 34653	1919± 5083	34770± 24249	863± 6702	17426± 18425	5191± 18803	2581± 8349	3753± 2770
95% CI	98366, 421488	33577, 66981	-460, 4298	24019, 45522	-2708, 4434	8803, 26049	-3146, 13528	-1120, 6283	2525, 4982
p-value#	<b>0.000013</b>	<b>0.000156</b>	0.174533	<b>0.000674</b>	0.716412	<b>0.012299</b>	0.871679	0.990057	0.218368
r	0.91	0.32	0.80	0.82	0.83	0.85	0.80	0.93	0.99
ICC (95% CI)	0.10 (0.0, 0.45)	0.06 (0.0,0.32)	0.75 (0.48,0.90)	0.24 (0.0, 0.59)	0.81 (0.54,0.93)	0.34 (0.0,0.67)	0.76 (0.52,0.89)	0.93 (0.83,0.97)	0.88 (0.11,0.97)
$\beta_1^{**}$	11.3153	3.3127	1.1761	3.1487	1.0893	2.3273	1.1211	1.0637	1.7240
Lower Level of Agreement (M±SD)	-376845± 133378	-17641± 13833	-8045± 1976	-12758± 8973	-12274± 2925	-18687± 7162	-31662± 6958	-13782± 3089	-1676± 1025
95% CI	-658248, -95442	-46703, 11421	-12181, -3909	-31418, 5902	-18510, -6037	-33678, 3697	-46131, -17193	-20207, 7358	-3807, 455
Upper Level of Agreement (M±SD)	896699± 133378	118199± 13833	11882± 1975	82298± 8973	14000± 2925	53540± 7162	42044± 6957	18945± 3089	9183± 1025
95% CI	615296, 1178102	89138, 147261	7746, 16018	63638, 100959	7764, 20236	38549, 68531	27575, 56513	12520, 25370	7051, 11315


\* Shapiro-Wilk Test of Normality of Differences;

\*\* Regression equation:  $Expected(E) = \beta_1 Reported(R) + \epsilon$

# pvalue- Based on Kolmogorov-Smirnov test for comparing distributions







**Figure 1 :** Bland-Altman plot showing agreement between WHO reported and expected values of excess mortalities due to Covid-19 of the country (A) India, (B) Indonesia (C) Italy, (D) Russia, (E) United Kingdom, (F) Mexico, (G) The United States of America, (H) Brazil, and (I) Peru