



UWL REPOSITORY

repository.uwl.ac.uk

Systematic literature review of the prevalence, pattern, and determinant of multimorbidity among older adults in Nigeria

Ahmed, Abdulsalam, Khan, Hafiz T.A. ORCID: <https://orcid.org/0000-0002-1817-3730> and Lawal, Muili ORCID: <https://orcid.org/0000-0002-1502-3884> (2023) Systematic literature review of the prevalence, pattern, and determinant of multimorbidity among older adults in Nigeria. Health Services Research & Managerial Epidemiology. ISSN 2333-3928

<http://dx.doi.org/10.1177/23333928231178774>

This is the Accepted Version of the final output.

UWL repository link: <https://repository.uwl.ac.uk/id/eprint/9979/>

Alternative formats: If you require this document in an alternative format, please contact: open.research@uwl.ac.uk

Copyright:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy: If you believe that this document breaches copyright, please contact us at open.research@uwl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Abstract

Introduction

Multimorbidity is a rising health issue globally and it is likely to become challenging in developing countries like Nigeria as they experience economic, demographic, and epidemiology transition. Yet, evidence of prevalence and patterns of multimorbidity, and their determinants, are scarce. This study aims to systematically review studies of the prevalence, patterns, and determinants of multimorbidity in Nigeria.

Methods

Studies were identified by searching five electronic databases (PubMed, Web of Science, CINAHL, PsycINFO, Africa Index Medicus/Global Index Medicus). Multimorbidity as well as other versions of it was used to search. The prevalence and determinants were also searched. According to pre-established inclusion criteria, and using different search strategies, 6 articles were included. The quality and risk of bias were assessed using Joanna Briggs Institute (JBI) appraisal tool for prevalence studies. Two researchers assessed the eligibility of studies for inclusion. The Protocol was registered on PROSPERO Ref no. CRD42021273222. The overall prevalence, pattern, and determinants were analyzed.

Results

We identified 6 eligible publications describing studies that included a total of 3332 (men 47.5%, women 52.5%) patients from 4 states plus the federal capital territory Abuja. The multimorbidity prevalence ranges from 27% to 74 % among elderly Nigerians. Cardiovascular together with metabolic and/or musculoskeletal conditions were the frequent patterns of multimorbidity. A positive association was observed between age and multimorbidity in most studies. Other factors associated with multimorbidity were female gender, low education status, poor monthly income/unemployment, hospitalization, medical visits, and emergency services.

Conclusion

There has been a growing need for more applied health services research to understand better and manage multimorbidity in developed countries. The scarcity of studies in our review reveals that multimorbidity is not a priority area of research in Nigeria and this will continue to hinder policy development in that area.

Keywords: Multimorbidity, Prevalence, Pattern, Determinant, Older Adults, Nigeria

Introduction

The increase in average life span and transformation in global age structure towards a rapidly ageing society is a major success for medical and public health systems. Though lengthening lifespans is a global trend, the impact is not spread equally across the world (Khan, 2019). A huge difference still exists in life expectancy across the globe, it is said to be less than 60 years in sub-Saharan Africa and it is higher in first-world countries and even higher than 80 years in Japan (Roser et al., 2013). They also reported that global life expectancy increased from an average of 29 to 73 years in 2019. This success epitomizes new challenges for public health policy to ensure that healthy life expectancy is increased rather than just life expectancy (Brayne, 2007). This is because an ageing population presents many challenges and ignoring them could undermine the potential benefits and opportunities that living for longer can bring (Khan, 2019).

Ageing represents the greatest risk factor for disease and brings with it the chronic uncontrol of multiple organ systems (Fabbri et al., 2015). In other words, the uncompromising form of reality is that chronic diseases rarely occur in isolation, and as life expectancy increases, people acquire a growing number of illnesses (Barnett et al., 2012a). The number of people affected by multiple chronic diseases a condition termed multimorbidity is increasing dramatically around the world and caring for them has placed substantial stress on many health systems (Navickas et al., 2016). Although this rising burden of chronic diseases has attracted the attention of public health researchers and policymakers worldwide, studies have shown that evidence on the epidemiology of multimorbidity in Low- and Middle-Income Countries (LMICs) is limited even though the region bears 80% of the global burden of non-communicable diseases (NCDs) (Hunter & Reddy, 2013). Healthcare utilization and cost have surged in LMICs as a result of the prevalence of multimorbidity which places strain on the health system (Frølich et al., 2019; Sum et al., 2019). Multimorbidity management requires a lot of resources that are hard work for both the patients and practitioners, especially when deepened with socioeconomic deprivation (O'Brien et al., 2011). Likewise, multimorbid patients are prone to frequent hospitalization, polypharmacy, treatment burden, and mortality (Duerden et al., 2013; Palmer et al., 2018).

Recent studies reported that only 5% of multimorbidity research studies originated in LMICs, out of which were confined to only six middle-income countries (Brazil, China, South Africa, India, Mexico, and Iran) (Xu et al., 2017). Similarly, most of the recognized studies on multimorbidity

extrapolated from the global population through the largest systematic review of the prevalence of multimorbidity conducted up to date for over 25 years (1992-2017), by Nguyen et al. (2019) were largely skewed to the other region of the world excluding Africa. This skewed distribution of multimorbidity studies demonstrates that there is a lack of attention on studying the phenomenon in other LMICs where it is likely to be more prevalent.

It has been estimated that the number of people experiencing multimorbidity is projected to rise along with population ageing by >1% per annum until 2030 (Yoon et al., 2014). Therefore, there is a need for greater insight and an up-to-date understanding of the prevalence and patterns of multimorbidity, especially among the older population to inform preventive strategies in LMICs like Nigeria. There is a dearth of knowledge on the prevalence, pattern, and determinants of multimorbidity in Nigeria to our knowledge. And hence, this systematic review of literature was conducted to determine (I) the prevalence of multimorbidity in older adults aged 60 years and above in Nigeria. (II) The common multimorbidity disease clusters in Nigeria and (III) the determinant of multimorbidity disease in Nigeria.

Methods

We conducted a systematic review that was preceded by a designed priori protocol (S1 File); following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist (S2 File) (Moher, 2009) and the PRISMA Protocols statement (S3 File) (Shamseer et al., 2015) respectively. The Protocol was registered on PROSPERO Ref no. CRD42021273222 and available at https://www.crd.york.ac.uk/prospero/display_record.php?RecordID=273222.

Inclusion and exclusion criteria

Included articles were peer-reviewed original articles with abstracts in English. However, conference presentations, opinion articles, books, and dissertations are excluded. The articles are observational cross-sectional articles on multimorbidity with a well-defined population 60 years and above conducted in Nigeria. The setting of the study was either community-based or health facility-based involving either or both inpatient and outpatient. Any other studies besides cross-sectional studies like cohort studies, and experimental studies were excluded. In addition, papers with single-review morbidity studies and without a clear description of the population were also excluded. Articles about the prevalence, pattern, and determinants of multimorbidity in Nigeria

were included. Where multimorbidity has not been clearly defined, articles documenting two or more chronic medical conditions were considered in this (SR) even if they did not mention multimorbidity. Papers with single-review morbidity and studies with suboptimal methodology were excluded. No limitations were placed on the years of publication.

Outcomes

Recent studies centered on defined multimorbidity as the simultaneous presence of more than one chronic medical condition in the same individual (Cucinotta & Vanelli, 2020; Javanmardi et al., 2020; Johnston et al., 2019; Sinclair & Abdelhafiz, 2020; van den Akker et al., 1996) and multimorbidity patterns as the most frequent combination of specific disease pairs and the groups of health conditions with the highest degree of association using the corresponding statistical analyses of either cluster and factor analysis (Violan et al., 2014).

Search strategy and study selection

We conducted an online literature search on PubMed, Web of Science, CINAHL, PsycINFO, Africa Index Medicus/Global Index Medicus electronic databases, from inception up to 16 of August 2021. Additionally, a corresponding internet search was done in Google Scholar, Google, and an online search from Africa Journal Online (AJOL) applying the same algorithm used in the bibliographic database search. However, where necessary the search was altered in line with the search engines and databases used. Screening of the reference list of included articles for the likelihood of relevant articles was conducted. The search terms included multimorbidity and other versions like ‘multimorbidities’, ‘multi-morbidities’, ‘multi morbidity’, and multiple morbidities. We excluded ‘comorbidity’ and other synonyms in our search strategy. And prevalence or epidemiology AND (pattern) AND determinants were used. This was done by using the ‘AND’ and ‘OR’ Boolean operators where appropriate (online supplement). These terms will be further restricted by location ‘Nigeria (Abia OR Adamawa OR Akwa Ibom OR Anambra or Bauchi or Bayelsa OR Benue OR Borno OR Cross River OR Delta OR Ebonyi OR Edo State OR Ekiti OR Enugu OR Gombe OR Imo OR Jigawa OR Kaduna OR Kano OR Katsina OR Kebbi OR Kogi OR Kwara OR Lagos OR Nasarawa OR Niger OR Ogun OR Ondo OR Osun OR Oyo OR Plateau OR Rivers OR Sokoto OR Taraba OR Yobe OR Zamfara OR ABUJA OR FCT)’. The details of the search in specific databases are given in the study protocol (S1 File).

We downloaded and exported all identified citations to RefWorks referencing software manager. Duplicates were excluded using the Mendeley reference manager deduplication function. Afterward, the citations were exported from the reference manager into Rayyan systematic review software (Ouzzani et al., 2016) for the screening process. Title/abstract and full-text screening were carried out on the Rayyan software. The first reviewer (AA) screened the title and abstract of all initial articles. To ensure that appropriate studies were not unexploited a random 10% of all references were tested by the second reviewer (ML). (AA) and (ML) screened the full text independently and any area of variance was fixed between the two reviewers and also with input from the third reviewer (HK). The most suitable data was included from multiple studies from the equivalent dataset. A PRISMA flow diagram is attached to show the detail of the study selection decisions made.

Data extraction

The extraction of data was conducted simultaneously with full-text searching. The relevant information was extracted from each article included and recorded immediately in the data extraction file (MS Excel). (AA) and (ML) independently extract the data. The extracted data are citation details: authors, title, journal, and year, details of the study: study setting (community or health facility) study design, period of data collection, location of the study, size of the sample, case definition, characteristics of the participants, description of main results: percentage prevalence of multimorbidity and the most common disease clusters in the study sample.

Study quality assessment

Assessing the study characteristics and the risk of bias was done by two reviewers. The Joanna Briggs Institute (JBI) critical appraisal tool for prevalence studies was used (Joanna Briggs Institute, 2017). The JBI uses 9 items. There are four possible answers to the nine items (yes, no, not clear, and not applicable). The threshold for the conversion of the JBI is as follows. Any item with a yes gets a score of 1, for no and unclear the score is zero, and for not applicable items are not included in the % calculation. A score of 60% and above was regarded as good quality. Based on the JBI data quality assessment system, all 6 studies were rated good quality. 4 studies had a score of 100%, one study have 88% and one study had 62.5%. The results from the two researchers were compared and differences were discussed between them.

Ethics and dissemination


This review received ethics approval as part of a larger project from the College of Nursing, Midwifery, and Healthcare, Research Ethics Panel (ethical approval number 1055). This research provided information on the prevalence of multimorbidity and other studied outcomes in Nigeria. The findings of this SR will be disseminated in a peer-reviewed journal article.

Results

Study overview

The processes of screening articles are shown on the PRISMA flowchart in Figure 1. The preliminary search yielded 738 titles; this was reduced to 581 after deduplication in the RefWorks citation manager software. In the title and abstract screening process 567 were excluded leaving only 14 articles for assessment and only 6 were included in the narrative synthesis (S1Table).

Figure 1 shows the Prisma flow chart



Study characteristics

Although the studies are relatively few, available ones are spread to cover the northern and southern parts of the country. Two studies each were conducted in north-central of Nigeria (Abdulraheem et al., 2017; Adams & Abubakar, 2019). Two studies were also conducted in Kano state northwestern Nigeria (Abdulazeez et al., 2021; Olawumi et al., 2021). One study each was conducted in the western and eastern parts of Nigeria respectively (Faronbi et al., 2020; Nwani & Isah, 2016a). The total number of participants across the six studies was 3332 (men: 47.5%, women: 52.5%) (Abdulazeez et al., 2021; Abdulraheem et al., 2017; Adams & Abubakar, 2019; Faronbi et al., 2020; Nwani & Isah, 2016a; Olawumi et al., 2021). The sample sizes of the included studies range from 333 to 1650 participants see Table 1. All included studies were published after 2013 and the majority were in the last 6 years. Four studies conducted primary cross-sectional descriptive using either or a combination of clinical evaluation and administration of a questionnaire, interviews, and review of medical records (Abdulazeez et al., 2021; Abdulraheem et al., 2017; Faronbi et al., 2020; Olawumi et al., 2021). One study conducted a cross-sectional retrospective study over 12 months from January to December 2018 (Adams & Abubakar, 2019), and the other study longitudinal prospective study both using medical health records (Nwani & Isah, 2016b). Two studies were carried out at the family medicine/ outpatient department of Aminu Kano teaching hospital Kano, Nigeria respectively (Abdulazeez et al., 2021; Olawumi et al., 2021). One study each was conducted at Nnamdi Azikiwe University Teaching Hospital (NAUTH), Nnewi Anambra state Nigeria (Nwani & Isah, 2016a), and General Out-Patient Clinic of the UATH Gwagwalada, Abuja, Nigeria (Adams & Abubakar, 2019), Osogbo, and Osun State, Nigeria respectively (Faronbi et al., 2020). Four studies were hospital-based (Abdulazeez et al., 2021; Adams & Abubakar, 2019; Nwani & Isah, 2016b; Olawumi et al., 2021) while two were conducted in the community (Abdulraheem et al., 2017; Faronbi et al., 2020).

Study quality assessment

The result of the study quality assessment is provided in S2 **Table**. Of the included studies, 66% were high quality, and the remaining studies were medium quality. There was no study with low quality. The threshold for the conversion of the JBI is as follows. Any item with a yes gets a score of 1, for no and unclear the score is zero, and for not applicable items are not included in the % calculation. A score of 60% and above was regarded as good quality (high and medium).

Based on the JBI data quality assessment system, all 6 studies were rated good quality. 4 studies had a score of 100%, one study have 88% and one study had 62.5%. The results from the two researchers were compared and differences were discussed between them.

Outcomes

Primary outcome

The definition of multimorbidity used, the number of disease conditions included in the study, and how the disease conditions were measured are all known facts that influence the prevalence of multimorbidity. However, all included studies used a ‘count’ of the number of diseases to define multimorbidity, and multimorbidity was defined by having two or more diseases in an individual. All the included studies specified they were only focused on chronic conditions. One study qualified chronic diseases as compiled and counted by the world health organization (WHO) (Nwani & Isah, 2016b). They stated that WHO defines chronic diseases as health problems that require ongoing management over a period of years or decades. Four studies draw and classify the chronic health problems of interest from the 147 International Classification of Primary Care, second edition (ICPC-2) rubrics list gathered by the Family Medicine Research Centre, University of Sydney (Abdulazeez et al., 2021; Abdulraheem et al., 2017; Nwani & Isah, 2016a; Olawumi et al., 2021). Two studies do not clearly measure (Adams & Abubakar, 2019; Faronbi et al., 2020).

Table 1 shows study characteristics

Study (state of study)	Study setting	Study design	Data collection period	Data source	Sample size	Age of participants
Nwani and Isah, 2016 (Anambra state)	Nnamdi Azikiwe University Teaching Hospital (NAUTH), Nnewi Anambra State Nigeria	prospective study	January 1, 2009, to December 31, 2009	Not reported	345 patients	patients aged 65 years
Adams and Abubakar, 2018 (Abuja)	General Out-Patient Clinic of the UATH Gwagwalada, Abuja.	Cross-sectional retrospective study	12 months, January 2018- December 2018	Medical health record	333 patients	60 and above years

Olawumi et al., 2021 (Kano)	conducted in the family medicine clinic (FMC) of Aminu kano teaching hospital kano	descriptive cross-sectional study	5th October 2020 to 28th December 2020.	Clinical evaluation and administration of a questionnaire	348 patients	60 and above years
Abdulraheem et al., 2017 (Niger)	The study was carried out in Niger State, Nigeria	A descriptive cross-sectional study	August 2014 to February 2015	Data were collected by questionnaire, interviews, review of medical records and clinical examination	conducted among 1650 rural elderly populations attending primary health care centers	age 60 years and above
Faronbi, Ajadi and Gobbens, 2020 (Osun state)	Osogbo, Osun State, Nigeria.	A cross-sectional study	Data collection took six weeks (between October and November 2015)	Data were collected by questionnaire	400	60 years of age and above
Abdulazeez et al., 2021 (Kano)	General Outpatient Clinic of Aminu Kano Teaching Hospital (AKTH) Kano	A descriptive cross-sectional study	May to June 2018.	interviewer-administered questionnaire	384	60 years and above

228

229

230 **Prevalence of multimorbidity**

231 All the included studies measured the prevalence of multimorbidity out of which 5 was among
232 60 years and above (Abdulazeez et al., 2021; Abdulraheem et al., 2017; Adams & Abubakar, 2019;
233 Faronbi et al., 2020; Olawumi et al., 2021) and one study uses 65 years and above (Nwani & Isah,
234 2016a). One study conducted in Oshogbo in Osun state reported the lowest prevalence of 27%
235 (Faronbi et al., 2020) while another conducted in Kano, kano state-reported prevalence of 74%
236 (Olawumi et al., 2021). The overall estimate ranges from 27% to 74.4% among elderly Nigerians.
237 The most common study design observed is cross-sectional (Abdulazeez et al., 2021; Abdulraheem
238 et al., 2017; Adams & Abubakar, 2019; Faronbi et al., 2020; Olawumi et al., 2021). In addition to the
239 common cut-off point of 2 chronic diseases used in 5 studies, one study also investigated the

240 prevalence estimate when multimorbidity was defined as ‘the co-occurrence of three or more
 241 chronic diseases’ (Abdulazeez et al., 2021). The higher the number of diseases in the operational
 242 definition, the lower the prevalence. See table 2

243

Table 2 shows the prevalence and pattern of multimorbidity among elderly Nigerians

Study (state of study)	Prevalence (%)	Patterns of multimorbidity
Nwani and Isah, 2016 (Anambra state)	The overall prevalence of multimorbidity among the elderly population studied was 49%. Two chronic diseases were present in 39.4% (n = 136), whereas three or more chronic diseases were present in 9.6%.	No pattern was reported. However, a percentage of single morbidity was reported
Adams and Abubakar, 2018 (Abuja)	The majority 236 (71%) of the study participants had multiple morbidities.	No morbidity pattern was reported. The cardiovascular system was the most affected system with 227 of the study population followed by the musculoskeletal system 90 and 84 had metabolic derangement. The least affected system was the Ears Nose and Throat with 13 persons.
Olawumi <i>et al.</i> , 2021 (Kano)	The prevalence of multimorbidity in this study is 74.4%.	No multimorbidity pattern was reported. CV diseases were the most prevalent morbidity (88.5%) among the respondents, followed by the diseases of the musculoskeletal system (42%). Hypertension was the most prevalent (50%) among all the CV diseases.
Abdulraheem et al., 2017 (Niger)	The percentage of participants with multimorbidity was 68.4% for 2 or more and 57.3 for 3 or more morbidities.	The most prevalent dyads of morbidities were hypertension and diabetes (31.4%), and hypertension and heart disease (25.6%). For triads of morbidities, the highest prevalence was found in HBP, diabetes and heart problem (10.3), and HBP, heart problem and Osteoarthritis (9.8%).
Faronbi, Ajadi and Gobbens, 2020 (Osun state)	This study also showed that multimorbidity is prevalent (27.0%) among the older adults in Nigeria.	No pattern was reported. However, percentage of single morbidity was reported
Abdulazeez <i>et al.</i> , 2021 (Kano)	More than half 190 (68.2%) of the participants had a 201(72.0%) had two or more chronic diseases (multimorbidity).	The commonest multimorbidity pattern based on system cluster were cardio-metabolic-mechanical conditions 42(15.1%). The top three frequent patterns of multimorbidity involving two clusters (dyad) of chronic diseases were hypertension-diabetes 7(17%), followed by hypertension overweight 5(12.1%) and hypertension osteoarthritis 4(9.7%). The commonest triads of chronic

		diseases were hypertension-diabetes-osteoarthritis 15(22.1%). The commonest quartet of chronic disease was hypertension diabetes-osteoarthritis-depression 3(5.9%). The commonest quintet of chronic diseases was hypertension diabetes-osteoarthritis-visual impairment-obesity/overweight 3(7.3%).
--	--	--

244

245 **Secondary outcomes**

246 **Pattern of multimorbidity**

247 The results of the studies were difficult to compare due to how data were reported. Only two
248 studies reported the most frequent common pairs (Abdulazeez et al., 2021; Abdulraheem et al.,
249 2017), See table 2. The study by Abdulraheem et al., (2018) reported the most prevalent dyads of
250 morbidities were hypertension and diabetes (31.4%), and hypertension and heart disease
251 (25.6%). For triads of morbidities, the highest prevalence was found in HBP, diabetes, and heart
252 problem (10.3), and HBP, heart problems, and Osteoarthritis (9.8%) (Abdulraheem et al., 2017).
253 While Abdulazeez et al., (2021) reported the commonest multimorbidity pattern based on system
254 clusters was cardio-metabolic-mechanical conditions at 15.1% (Abdulazeez et al., 2021). The top
255 three frequent patterns of multimorbidity involving two clusters (dyad) of chronic disease were
256 hypertension-diabetes (17%) followed by hypertension-overweight (12.1%) and hypertension-
257 osteoarthritis (9.7). The commonest triad of chronic disease was hypertension-diabetes-
258 osteoarthritis (22.1%). The commonest quartet of chronic diseases was hypertension-diabetes-
259 osteoarthritis- depression (5.9%) (Abdulazeez et al., 2021). The commonest quintet of chronic
260 disease was hypertension-diabetes-osteoarthritis-visual impairment/overweight (7.3%). The
261 remaining four studies were not structured to display the pairs but the pattern reported that
262 hypertension was the commonest in all the studies (Abdulazeez et al., 2021).

263

264 **Determinant of multimorbidity**

265 Age was the frequently studied determinant of multimorbidity (Barnett et al., 2012b; Marengoni et
266 al., 2008; Minas et al., 2010; Prados-Torres et al., 2012; Van den Akker et al., 1998). Although the
267 determinant of multimorbidity was not assessed in all the included studies, 3 studies reported the
268 determinants of multimorbidity (Abdulazeez et al., 2021; Abdulraheem et al., 2017; Adams &

Abubakar, 2019). One study reported that multi-morbidity occurred more in males (73.4% of males; OR = 1.062; CI = 0.926-1.219), aged 70 – 79 years (72.2%), unskilled workers (73.8%), and urban dwellers (73% of patients living in urban areas) (Adams & Abubakar, 2019). Although the p values show that these associations were not statistically significant however the relatively small sample size could account for this. However, a larger study by (Abdulraheem et al., 2017) showed that age and sex are independent risks factors for multi-morbidity (Abdulraheem et al., 2017). They further stated that apart from age, factors most strongly and independently associated with multi-morbidity were female gender, low education status, poor monthly income/ unemployment, hospitalization, medical visits, and emergency services. One study reported that apart from age, factors most strongly and independently associated with multi-morbidity were female gender, low education status, poor monthly income/ unemployment, hospitalization, medical visits and emergency services (Abdulraheem et al., 2017).

Another study reported that Participants with formal education were more than 30% less likely to have multimorbidity than those without formal education (Abdulazeez et al., 2021). Similarly, participants that were employed were almost 40% less likely to have multimorbidity than those that were unemployed (Abdulazeez et al., 2021). The same study reported that participants that were overweight/obese had higher chances of developing multimorbidity when compared with individuals with normal BMI. And that participants that were functionally dependent were 20 times more likely to have multimorbidity than functionally independent elderly participants (Abdulazeez *et al.*, 2021).

Discussion

This systematic review provides a well-informed and comprehensive analysis of multimorbidity prevalence, pattern, and determinants in Nigeria. We identified 6 articles across 5 states in Nigeria. Earlier authors of systematic literature reviews of multimorbidity also noted the limited representation of developing countries in multimorbidity research (Khanam et al., 2011). In essence, multimorbidity is not a priority area of research in Nigeria and this will continue to hinder policy development in that area. Our review shows that prevalence estimates varied markedly according to age, gender, marital status, marital setting, tribe, educational levels, adequate income, living conditions, family support, and operational definitions of multimorbidity

(Abdulazeez et al., 2021; Abdulraheem et al., 2017; Adams & Abubakar, 2019; Nwani & Isah, 2016b). However, the findings from our studies were consistent with other previous studies and systematic reviews (Marengoni et al., 2011; Nguyen et al., 2019; Violan et al., 2014). While the prevalence estimates varied between and within age groups, most studies in our sample indicated multimorbidity as a common phenomenon in individuals 60 years and above. Where prevalence estimates by gender were reported, it showed variation, two studies reported higher prevalence among men (Adams & Abubakar, 2019; Nwani & Isah, 2016a). However, in one study females appeared to have a higher multimorbidity prevalence than males in studies (Abdulraheem et al., 2017). This is suggestive of an association between sex and multimorbidity, evidence of which was provided in multiple studies (Alimohammadian et al., 2017; Munn et al., 2014; Nguyen et al., 2019). The higher the number of diseases in the operational definition, the lower the prevalence. This was observed in our review, the percentage of participants with multimorbidity was 68.4% for 2 or more and 57.3 for 3 or more morbidities (Abdulraheem et al., 2017). This finding supported an observation where it was found that from 44% when multimorbidity was defined as 2 diseases, the prevalence reduced to 27% for 3 diseases, 15% for 4 diseases, 7% for 5 diseases, and only 3% for 6 diseases (Harrison et al., 2014). The highest prevalence estimates in our sample were reported in studies that used 2 diseases to define multimorbidity 71% (Adams & Abubakar, 2019) and 74% (Olawumi et al., 2021). The combination of diseases may make multimorbidity prevalence differ significantly (Fortin et al., 2012; Harrison et al., 2014). In the existing literature, a range of different combinations has been proposed from a list of 16 chronic diseases (Ferrer et al., 2017) to a list of 291 diseases (Quiñones et al., 2016) and anything in between (Van den Bussche et al., 2011). Ferrer et al. (Ferrer et al., 2017) argued that an open list of diagnoses should be used since it gave the highest prevalence estimate. For our studies, most of the chronic health problems of interest were drawn from the 147 International Classification of Primary Care, second edition (ICPC-2) rubrics list gathered by the Family Medicine Research Centre, the University of Sydney (WHO, 2022). The known fact is that there were no specific criteria for disease inclusion in these studies because of the lack of a standardized list and they were often determined by the author's expertise and experience. However, the most common conditions included were those that have the highest prevalence of clinical relevance.

While hypertension and diabetes mellitus appear to be common findings in this study, this is not completely different from other parts of the world. For example, in Korean adult men, diabetes mellitus and hypertension yielded the highest probability of multimorbidity, and for women polyarthritis and hypertension (Lee et al., 2020). In the United state, the most prevalent dyads of multimorbidity in men after 65 years of age is arthritis and hypertension followed by diabetes mellitus and hypertension in both sexes (Goodman et al., 2016). The most prevalent cluster in all four strata of multimorbidity included hypertension diseases and metabolic disorders (Foguet-Boreu et al., 2015).

Strengths and limitations

Our study selection, screening processes, search strategy, inclusion criteria, and quality assessment were wide-ranging and exhaustive. Our review included studies in both the hospital and communities and the first of its kind to conduct SLR of prevalence, pattern, and determinant of multimorbidity in Nigeria. This review, however, was not without limitations. Evaluations of prevalence, determinants, and patterns in our study are limited by the methods used in the primary studies. Similarly, all the measures of multimorbidity used in these studies were mostly disease count and disease count are only one of the 20 measures to date.

Conclusion

The overall prevalence estimates range from 27% to 74.4% among elderly Nigerians. Although, the disease pair varies among studies, hypertension, and diabetes appear to be the commonest dyads. Determinants of multimorbidity in our study include age, gender, and lower socioeconomic status. The scarcity of studies in our review reveals that multimorbidity is not a priority area of research in Nigeria and this will continue to hinder policy development in that area.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

No funding support was received for this research.

References

Abdulazeez, Z. U., Grema, B. A., Michael, G. C., & Abdulkadir, Z. (2021). Multimorbidity and Functional Status of the Elderly in a Primary Care Setting of Northern Nigeria: A Cross-Sectional Study. *West African Journal of Medicine*, 38(7), 620-628.

Abdulraheem, I. S., Amodu, M., Salami, S. K., Adegboye, A., Fatiregun, A., & Tobin-West, C. (2017). Prevalence and Pattern of Multi-Morbidity among Elderly People in Rural Nigeria: Implications for Health Care System, Research and Medical Education. *Journal of Community Medicine & Health Care*, 2(3), 1019. <https://austinpublishinggroup.com/community-medicine/fulltext/jcmhc-v2-id1019.pdf>

Adams, O., & Abubakar, I. (2019). Morbidity Patterns of Elderly Patients Attending the General Out-Patient Clinic of a Tertiary Centre in North-Central – Nigeria. *Texila International Journal of Public Health*, , 1-7. 10.21522/TIJPH.2013.08.03.Art034

Alimohammadian, M., Majidi, A., Yaseri, M., Ahmadi, B., Islami, F., Derakhshan, M., Delavari, A., Amani, M., Feyz-Sani, A., & Poustchi, H. (2017). Multimorbidity as an important issue among women: results of a gender difference investigation in a large population-based cross-sectional study in West Asia. *BMJ Open*, 7(5), e013548.

374 Barnett, K., Mercer, S. W., Norbury, M., Watt, G., Wyke, S., & Guthrie, B. (2012a).
 375 Epidemiology of multimorbidity and implications for health care, research, and medical
 376 education: a cross-sectional study. *The Lancet*, 380(9836), 37-43.

377 Barnett, K., Mercer, S. W., Norbury, M., Watt, G., Wyke, S., & Guthrie, B. (2012b).
 378 Epidemiology of multimorbidity and implications for health care, research, and medical
 379 education: a cross-sectional study. *The Lancet*, 380(9836), 37-43.

380 Brayne, C. (2007). The elephant in the room—healthy brains in later life, epidemiology and
 381 public health. *Nature Reviews Neuroscience*, 8(3), 233-239.

382 Cucinotta, D., & Vanelli, M. (2020). WHO declares COVID-19 a pandemic. *Acta Bio Medica:*
 383 *Atenei Parmensis*, 91(1), 157.

384 Duerden, M., Avery, T., & Payne, R. (2013). Polypharmacy and medicines optimisation. *Making*
 385 *it Safe and Sound*. London: The King's Fund,

386 Fabbri, E., Zoli, M., Gonzalez-Freire, M., Salive, M. E., Studenski, S. A., & Ferrucci, L. (2015).
 387 Aging and multimorbidity: new tasks, priorities, and frontiers for integrated gerontological
 388 and clinical research. *Journal of the American Medical Directors Association*, 16(8), 640-
 389 647.

390 Faronbi, J., Ajadi, A., & Gobbens, R. (2020). Associations of chronic illnesses and socio-
 391 demographic factors with health-related quality of life of older adults in Nigeria: A cross-
 392 sectional study. *Ghana Medical Journal*, 54(3), 164-172.

393 Ferrer, A., Formiga, F., Sanz, H., Almeda, J., & Padrós, G. (2017). Multimorbidity as specific
 394 disease combinations, an important predictor factor for mortality in octogenarians: the
 395 Octabaix study. *Clinical Interventions in Aging*, 12, 223.

396 Fortin, M., Bravo, G., Hudon, C., Vanasse, A., & Lapointe, L. (2005). Prevalence of
 397 multimorbidity among adults seen in family practice. *The Annals of Family Medicine*, 3(3),
 398 223-228.

399 Fortin, M., Stewart, M., Poitras, M., Almirall, J., & Maddocks, H. (2012). A systematic review
 400 of prevalence studies on multimorbidity: toward a more uniform methodology. *The Annals*
 401 *of Family Medicine*, 10(2), 142-151.

402 Frølich, A., Ghith, N., Schiøtz, M., Jacobsen, R., & Stockmarr, A. (2019). Multimorbidity,
 403 healthcare utilization and socioeconomic status: a register-based study in Denmark. *PloS*
 404 *One*, 14(8), e0214183.

405 Harrison, C., Britt, H., Miller, G., & Henderson, J. (2014). Examining different measures of
 406 multimorbidity, using a large prospective cross-sectional study in Australian general
 407 practice. *BMJ Open*, 4(7)

408 Hunter, D. J., & Reddy, K. S. (2013). Noncommunicable diseases. *New England Journal of*
 409 *Medicine*, 369(14), 1336-1343.

410 Javanmardi, F., Keshavarzi, A., Akbari, A., Emami, A., & Pirbonyeh, N. (2020). Prevalence of
 411 underlying diseases in died cases of COVID-19: A systematic review and meta-analysis.
 412 *PloS One*, 15(10), e0241265.

413 Joanna Briggs Institute. (2017). The Joanna Briggs Institute critical appraisal tools for use in JBI
 414 systematic reviews: checklist for prevalence studies. *Crit Apprais Checkl Preval Stud*, 7, 65-
 415 72.

416 Johnston, M. C., Crilly, M., Black, C., Prescott, G. J., & Mercer, S. W. (2019). Defining and
 417 measuring multimorbidity: a systematic review of systematic reviews. *European Journal of*
 418 *Public Health*, 29(1), 182-189.

419 Khan, H. T. (2019). Population ageing in a globalized world: Risks and dilemmas? *Journal of*
 420 *Evaluation in Clinical Practice*, 25(5), 754-760.

421 Marengoni, A., Angleman, S., Melis, R., Mangialasche, F., Karp, A., Garmen, A., Meinow, B.,
 422 & Fratiglioni, L. (2011). Aging with multimorbidity: a systematic review of the literature.
 423 *Ageing Research Reviews*, 10(4), 430-439.

424 Marengoni, A., Winblad, B., Karp, A., & Fratiglioni, L. (2008). Prevalence of chronic diseases
 425 and multimorbidity among the elderly population in Sweden. *American Journal of Public*
 426 *Health*, 98(7), 1198-1200.

427 Minas, M., Koukousias, N., Zintzaras, E., Kostikas, K., & Gourgoulialis, K. I. (2010). Prevalence
 428 of chronic diseases and morbidity in primary health care in central Greece: an
 429 epidemiological study. *BMC Health Services Research*, 10(1), 1-8.

430 Moher, D. (2009). Liberati A, Tetzlaff J, Altman DG, Group TP. *Preferred Reporting Items for*
 431 *Systematic Reviews and Meta-Analyses: The PRISMA Statement*. *PLoS Med*, 6(7), 1000097.

432 Munn, Z., Moola, S., Riitano, D., & Lis, K. (2014). The development of a critical appraisal tool
 433 for use in systematic reviews addressing questions of prevalence. *International Journal of*
 434 *Health Policy and Management*, 3(3), 123.

435 Navickas, R., Petric, V., Feigl, A. B., & Seychell, M. (2016). Multimorbidity: what do we know?
 436 What should we do? *Journal of Comorbidity*, 6(1), 4-11.

437 Nguyen, H., Manolova, G., Daskalopoulou, C., Vitoratou, S., Prince, M., & Prina, A. M. (2019).
 438 Prevalence of multimorbidity in community settings: A systematic review and meta-analysis
 439 of observational studies. *Journal of Comorbidity*, 9, 2235042X19870934.

440 Nwani, P. O., & Isah, A. O. (2016a). Chronic diseases and multimorbidity among elderly
 441 patients admitted in the medical wards of a Nigerian tertiary hospital. *Journal of Clinical*
 442 *Gerontology and Geriatrics*, 7(3), 83-86.

443 Nwani, P. O., & Isah, A. O. (2016b). Chronic diseases and multimorbidity among elderly
 444 patients admitted in the medical wards of a Nigerian tertiary hospital. *Journal of Clinical*
 445 *Gerontology and Geriatrics*, 7(3), 83-86.

446 O'Brien, R., Wyke, S., Guthrie, B., Watt, G., & Mercer, S. (2011). An 'endless struggle': a
 447 qualitative study of general practitioners' and practice nurses' experiences of managing
 448 multimorbidity in socio-economically deprived areas of Scotland. *Chronic Illness*, 7(1), 45-
 449 59.

450 Olawumi, A. L., Grema, B. A., Suleiman, A. K., Omeiza, Y. S., Michael, G. C., & Shuaibu, A.
 451 (2021). Nutritional status and morbidity patterns of the elderly in a Northwestern Nigerian
 452 hospital: A cross-sectional study. *Nigerian Postgraduate Medical Journal*, 28(3), 160.

453 Ouzzani, M., Hammady, H., Fedorowicz, Z., & Elmagarmid, A. (2016). Rayyan—a web and
 454 mobile app for systematic reviews. *Systematic Reviews*, 5(1), 1-10.

455 Palmer, K., Marengoni, A., Forjaz, M. J., Jureviciene, E., Laatikainen, T., Mammarella, F.,
 456 Muth, C., Navickas, R., Prados-Torres, A., & Rijken, M. (2018). Multimorbidity care
 457 model: Recommendations from the consensus meeting of the Joint Action on Chronic
 458 Diseases and Promoting Healthy Ageing across the Life Cycle (JA-CHRODIS). *Health*
 459 *Policy*, 122(1), 4-11.

460 Prados-Torres, A., Poblador-Plou, B., Calderón-Larrañaga, A., Gimeno-Feliu, L. A., González-
 461 Rubio, F., Poncel-Falcó, A., Sicras-Mainar, A., & Alcalá-Nalvaiz, J. T. (2012).
 462 Multimorbidity patterns in primary care: interactions among chronic diseases using factor
 463 analysis. *PloS One*, 7(2), e32190.

464 Quiñones, A. R., Markwardt, S., & Botoseneanu, A. (2016). Multimorbidity combinations and
 465 disability in older adults. *Journals of Gerontology Series A: Biomedical Sciences and*
 466 *Medical Sciences*, 71(6), 823-830.

467 Shamseer, L., Moher, D., Clarke, M., Gherzi, D., & Liberati, A. (2015). Research methods &
 468 reporting. *Bmj*, , 1-25.

469 Sinclair, A. J., & Abdelhafiz, A. H. (2020). Age, frailty and diabetes—triple jeopardy for
 470 vulnerability to COVID-19 infection. *EClinicalMedicine*, 22

471 Sum, G., Salisbury, C., Koh, G. C., Atun, R., Oldenburg, B., McPake, B., Vellakkal, S., & Lee,
 472 J. T. (2019). Implications of multimorbidity patterns on health care utilisation and quality of
 473 life in middle-income countries: cross-sectional analysis. *Journal of Global Health*, 9(2)

474 van den Akker, M., Buntinx, F., & Knottnerus, J. A. (1996). Comorbidity or multimorbidity:
 475 what's in a name? A review of literature. *The European Journal of General Practice*, 2(2),
 476 65-70.

477 Van den Akker, M., Buntinx, F., Metsemakers, J. F., Roos, S., & Knottnerus, J. A. (1998).
 478 Multimorbidity in general practice: prevalence, incidence, and determinants of co-occurring
 479 chronic and recurrent diseases. *Journal of Clinical Epidemiology*, 51(5), 367-375.

480 Van den Bussche, H., Koller, D., Kolonko, T., Hansen, H., Wegscheider, K., Glaeske, G., von
 481 Leitner, E., Schäfer, I., & Schön, G. (2011). Which chronic diseases and disease
 482 combinations are specific to multimorbidity in the elderly? Results of a claims data based
 483 cross-sectional study in Germany. *BMC Public Health*, 11(1), 1-9.

484 Violan, C., Foguet-Boreu, Q., Flores-Mateo, G., Salisbury, C., Blom, J., Freitag, M., Glynn, L.,
 485 Muth, C., & Valderas, J. M. (2014). Prevalence, determinants and patterns of
 486 multimorbidity in primary care: a systematic review of observational studies. *PloS One*,
 487 9(7), e102149.

WHO. (2022). *International Statistical Classification of Diseases and Related Health Problems (ICD) 11th Revision*. World health organization website.

<https://www.who.int/standards/classifications/classification-of-diseases>

Xu, X., Mishra, G. D., & Jones, M. (2017). Mapping the global research landscape and knowledge gaps on multimorbidity: a bibliometric study. *Journal of Global Health*, 7(1)

Yoon, J., Zulman, D., Scott, J. Y., & Maciejewski, M. L. (2014). Costs associated with multimorbidity among VA patients. *Medical Care*, , S31-S36.

References

Abdulazeez, Z. U., Grema, B. A., Michael, G. C., & Abdulkadir, Z. (2021). Multimorbidity and Functional Status of the Elderly in a Primary Care Setting of Northern Nigeria: A Cross-Sectional Study. *West African Journal of Medicine*, 38(7), 620-628.

Abdulraheem, I. S., Amodu, M., Salami, S. K., Adegboye, A., Fatiregun, A., & Tobin-West, C. (2017). Prevalence and Pattern of Multi-Morbidity among Elderly People in Rural Nigeria: Implications for Health Care System, Research and Medical Education. *Journal of Community Medicine & Health Care*, 2(3), 1019. <https://austinpublishinggroup.com/community-medicine/fulltext/jcmhc-v2-id1019.pdf>

Adams, O., & Abubakar, I. (2019). Morbidity Patterns of Elderly Patients Attending the General Out-Patient

507 Clinic of a Tertiary Centre in North-Central – Nigeria. *Texila International Journal of*
508 *Public Health*, , 1-7. 10.21522/TIJPH.2013.08.03.Art034

509 Alimohammadian, M., Majidi, A., Yaseri, M., Ahmadi, B., Islami, F., Derakhshan, M., Delavari,
510 A., Amani, M., Feyz-Sani, A., & Poustchi, H. (2017). Multimorbidity as an important issue
511 among women: results of a gender difference investigation in a large population-based
512 cross-sectional study in West Asia. *BMJ Open*, 7(5), e013548.

513 Barnett, K., Mercer, S. W., Norbury, M., Watt, G., Wyke, S., & Guthrie, B. (2012a).
514 Epidemiology of multimorbidity and implications for health care, research, and medical
515 education: a cross-sectional study. *The Lancet*, 380(9836), 37-43.

516 Barnett, K., Mercer, S. W., Norbury, M., Watt, G., Wyke, S., & Guthrie, B. (2012b).
517 Epidemiology of multimorbidity and implications for health care, research, and medical
518 education: a cross-sectional study. *The Lancet*, 380(9836), 37-43.

519 Brayne, C. (2007). The elephant in the room—healthy brains in later life, epidemiology and
520 public health. *Nature Reviews Neuroscience*, 8(3), 233-239.

521 Cucinotta, D., & Vanelli, M. (2020). WHO declares COVID-19 a pandemic. *Acta Bio Medica:*
522 *Atenei Parmensis*, 91(1), 157.

523 Duerden, M., Avery, T., & Payne, R. (2013). Polypharmacy and medicines optimisation. *Making*
524 *it Safe and Sound*. London: The King's Fund,

525 Fabbri, E., Zoli, M., Gonzalez-Freire, M., Salive, M. E., Studenski, S. A., & Ferrucci, L. (2015).
526 Aging and multimorbidity: new tasks, priorities, and frontiers for integrated gerontological

527 and clinical research. *Journal of the American Medical Directors Association*, 16(8), 640-
528 647.

529 Faronbi, J., Ajadi, A., & Gobbens, R. (2020). Associations of chronic illnesses and socio-
530 demographic factors with health-related quality of life of older adults in Nigeria: A cross-
531 sectional study. *Ghana Medical Journal*, 54(3), 164-172.

532 Ferrer, A., Formiga, F., Sanz, H., Almeda, J., & Padrós, G. (2017). Multimorbidity as specific
533 disease combinations, an important predictor factor for mortality in octogenarians: the
534 Octabaix study. *Clinical Interventions in Aging*, 12, 223.

535 Foguet-Boreu, Q., Violán, C., Rodriguez-Blanco, T., Roso-Llorach, A., Pons-Vigués, M., Pujol-
536 Ribera, E., Cossio Gil, Y., & Valderas, J. M. (2015). Multimorbidity patterns in elderly
537 primary health care patients in a South Mediterranean European region: a cluster analysis.
538 *PLoS One*, 10(11), e0141155.

539 Fortin, M., Bravo, G., Hudon, C., Vanasse, A., & Lapointe, L. (2005). Prevalence of
540 multimorbidity among adults seen in family practice. *The Annals of Family Medicine*, 3(3),
541 223-228.

542 Fortin, M., Stewart, M., Poitras, M., Almirall, J., & Maddocks, H. (2012). A systematic review
543 of prevalence studies on multimorbidity: toward a more uniform methodology. *The Annals*
544 *of Family Medicine*, 10(2), 142-151.

545 Frølich, A., Ghith, N., Schiøtz, M., Jacobsen, R., & Stockmarr, A. (2019). Multimorbidity,
546 healthcare utilization and socioeconomic status: a register-based study in Denmark. *PloS*
547 *One*, 14(8), e0214183.

548 Goodman, R. A., Ling, S. M., Briss, P. A., Parrish, R. G., Salive, M. E., & Finke, B. S. (2016).
549 Multimorbidity patterns in the United States: implications for research and clinical practice.
550 *Journals of Gerontology Series A: Biomedical Sciences and Medical Sciences*, 71(2), 215-
551 220.

552 Harrison, C., Britt, H., Miller, G., & Henderson, J. (2014). Examining different measures of
553 multimorbidity, using a large prospective cross-sectional study in Australian general
554 practice. *BMJ Open*, 4(7)

555 Hunter, D. J., & Reddy, K. S. (2013). Noncommunicable diseases. *New England Journal of*
556 *Medicine*, 369(14), 1336-1343.

557 Javanmardi, F., Keshavarzi, A., Akbari, A., Emami, A., & Pirbonyeh, N. (2020). Prevalence of
558 underlying diseases in died cases of COVID-19: A systematic review and meta-analysis.
559 *PloS One*, 15(10), e0241265.

560 Joanna Briggs Institute. (2017). The Joanna Briggs Institute critical appraisal tools for use in JBI
561 systematic reviews: checklist for prevalence studies. *Crit Apprais Checkl Preval Stud*, 7, 65-
562 72.

563 Johnston, M. C., Crilly, M., Black, C., Prescott, G. J., & Mercer, S. W. (2019). Defining and
 564 measuring multimorbidity: a systematic review of systematic reviews. *European Journal of*
 565 *Public Health*, 29(1), 182-189.

566 Khan, H. T. (2019). Population ageing in a globalized world: Risks and dilemmas? *Journal of*
 567 *Evaluation in Clinical Practice*, 25(5), 754-760.

568 Khanam, M. A., Streatfield, P. K., Kabir, Z. N., Qiu, C., Cornelius, C., & Wahlin, Å. (2011).
 569 Prevalence and patterns of multimorbidity among elderly people in rural Bangladesh: a
 570 cross-sectional study. *Journal of Health, Population, and Nutrition*, 29(4), 406.

571 Lee, Y., Kim, H., Jeong, H., & Noh, Y. (2020). Patterns of multimorbidity in adults: an
 572 association rules analysis using the Korea health panel. *International Journal of*
 573 *Environmental Research and Public Health*, 17(08), 2618.

574 Marengoni, A., Angleman, S., Melis, R., Mangialasche, F., Karp, A., Garmen, A., Meinow, B.,
 575 & Fratiglioni, L. (2011). Aging with multimorbidity: a systematic review of the literature.
 576 *Ageing Research Reviews*, 10(4), 430-439.

577 Marengoni, A., Winblad, B., Karp, A., & Fratiglioni, L. (2008). Prevalence of chronic diseases
 578 and multimorbidity among the elderly population in Sweden. *American Journal of Public*
 579 *Health*, 98(7), 1198-1200.

580 Minas, M., Koukousias, N., Zintzaras, E., Kostikas, K., & Gourgoulisanis, K. I. (2010). Prevalence
 581 of chronic diseases and morbidity in primary health care in central Greece: an
 582 epidemiological study. *BMC Health Services Research*, 10(1), 1-8.

583 Moher, D. (2009). Liberati A, Tetzlaff J, Altman DG, Group TP. *Preferred Reporting Items for*
584 *Systematic Reviews and Meta-Analyses: The PRISMA Statement*. *PLoS Med*, 6(7), 1000097.

585 Munn, Z., Moola, S., Riitano, D., & Lisy, K. (2014). The development of a critical appraisal tool
586 for use in systematic reviews addressing questions of prevalence. *International Journal of*
587 *Health Policy and Management*, 3(3), 123.

588 Navickas, R., Petric, V., Feigl, A. B., & Seychell, M. (2016). Multimorbidity: what do we know?
589 What should we do? *Journal of Comorbidity*, 6(1), 4-11.

590 Nguyen, H., Manolova, G., Daskalopoulou, C., Vitoratou, S., Prince, M., & Prina, A. M. (2019).
591 Prevalence of multimorbidity in community settings: A systematic review and meta-analysis
592 of observational studies. *Journal of Comorbidity*, 9, 2235042X19870934.

593 Nwani, P. O., & Isah, A. O. (2016a). Chronic diseases and multimorbidity among elderly
594 patients admitted in the medical wards of a Nigerian tertiary hospital. *Journal of Clinical*
595 *Gerontology and Geriatrics*, 7(3), 83-86.

596 Nwani, P. O., & Isah, A. O. (2016b). Chronic diseases and multimorbidity among elderly
597 patients admitted in the medical wards of a Nigerian tertiary hospital. *Journal of Clinical*
598 *Gerontology and Geriatrics*, 7(3), 83-86.

599 O'Brien, R., Wyke, S., Guthrie, B., Watt, G., & Mercer, S. (2011). An 'endless struggle': a
600 qualitative study of general practitioners' and practice nurses' experiences of managing
601 multimorbidity in socio-economically deprived areas of Scotland. *Chronic Illness*, 7(1), 45-
602 59.

603 Olawumi, A. L., Grema, B. A., Suleiman, A. K., Omeiza, Y. S., Michael, G. C., & Shuaibu, A.
604 (2021). Nutritional status and morbidity patterns of the elderly in a Northwestern Nigerian
605 hospital: A cross-sectional study. *Nigerian Postgraduate Medical Journal*, 28(3), 160.

606 Ouzzani, M., Hammady, H., Fedorowicz, Z., & Elmagarmid, A. (2016). Rayyan—a web and
607 mobile app for systematic reviews. *Systematic Reviews*, 5(1), 1-10.

608 Palmer, K., Marengoni, A., Forjaz, M. J., Jureviciene, E., Laatikainen, T., Mammarella, F.,
609 Muth, C., Navickas, R., Prados-Torres, A., & Rijken, M. (2018). Multimorbidity care
610 model: Recommendations from the consensus meeting of the Joint Action on Chronic
611 Diseases and Promoting Healthy Ageing across the Life Cycle (JA-CHRODIS). *Health*
612 *Policy*, 122(1), 4-11.

613 Prados-Torres, A., Poblador-Plou, B., Calderón-Larrañaga, A., Gimeno-Feliu, L. A., González-
614 Rubio, F., Poncel-Falcó, A., Sicras-Mainar, A., & Alcalá-Nalvaiz, J. T. (2012).
615 Multimorbidity patterns in primary care: interactions among chronic diseases using factor
616 analysis. *PloS One*, 7(2), e32190.

617 Quiñones, A. R., Markwardt, S., & Botoseneanu, A. (2016). Multimorbidity combinations and
618 disability in older adults. *Journals of Gerontology Series A: Biomedical Sciences and*
619 *Medical Sciences*, 71(6), 823-830.

620 Roser, M., Ortiz-Ospina, E., & Ritchie, H. (2013). Life expectancy. *Our World in Data*,

621 Shamseer, L., Moher, D., Clarke, M., Gherzi, D., & Liberati, A. (2015). Research methods &
622 reporting. *Bmj*, , 1-25.

623 Sinclair, A. J., & Abdelhafiz, A. H. (2020). Age, frailty and diabetes—triple jeopardy for
624 vulnerability to COVID-19 infection. *EClinicalMedicine*, 22

625 Sum, G., Salisbury, C., Koh, G. C., Atun, R., Oldenburg, B., McPake, B., Vellakkal, S., & Lee,
626 J. T. (2019). Implications of multimorbidity patterns on health care utilisation and quality of
627 life in middle-income countries: cross-sectional analysis. *Journal of Global Health*, 9(2)

628 van den Akker, M., Buntinx, F., & Knottnerus, J. A. (1996). Comorbidity or multimorbidity:
629 what's in a name? A review of literature. *The European Journal of General Practice*, 2(2),
630 65-70.

631 Van den Akker, M., Buntinx, F., Metsemakers, J. F., Roos, S., & Knottnerus, J. A. (1998).
632 Multimorbidity in general practice: prevalence, incidence, and determinants of co-occurring
633 chronic and recurrent diseases. *Journal of Clinical Epidemiology*, 51(5), 367-375.

634 Van den Bussche, H., Koller, D., Kolonko, T., Hansen, H., Wegscheider, K., Glaeske, G., von
635 Leitner, E., Schäfer, I., & Schön, G. (2011). Which chronic diseases and disease
636 combinations are specific to multimorbidity in the elderly? Results of a claims data based
637 cross-sectional study in Germany. *BMC Public Health*, 11(1), 1-9.

638 Violan, C., Foguet-Boreu, Q., Flores-Mateo, G., Salisbury, C., Blom, J., Freitag, M., Glynn, L.,
639 Muth, C., & Valderas, J. M. (2014). Prevalence, determinants and patterns of
640 multimorbidity in primary care: a systematic review of observational studies. *PloS One*,
641 9(7), e102149.

642 WHO. (2022). *International Statistical Classification of Diseases and Related Health Problems*
643 *(ICD) 11th Revision*. World health organization website. Retrieved 29/12/2021, from
644 <https://www.who.int/standards/classifications/classification-of-diseases>

645 Xu, X., Mishra, G. D., & Jones, M. (2017). Mapping the global research landscape and
646 knowledge gaps on multimorbidity: a bibliometric study. *Journal of Global Health*, 7(1)

647 Yoon, J., Zulman, D., Scott, J. Y., & Maciejewski, M. L. (2014). Costs associated with
648 multimorbidity among VA patients. *Medical Care*, , S31-S36.

649